



Relationship between vitamin D status and the vaginal microbiome during pregnancy

Kimberly K. Jefferson¹ · Hardik I. Parikh^{1,2} · Erin M. Garcia¹ · David J. Edwards^{2,3} · Myrna G. Serrano¹ · Martin Hewison⁴ · Judith R. Shary⁵ · Anna M. Powell⁶ · Bruce W. Hollis⁵ · Jennifer M. Fettweis^{1,7} · Jerome F. Strauss III^{2,7} · Gregory A. Buck¹ · Carol L. Wagner⁵

Received: 19 July 2018 / Revised: 22 January 2019 / Accepted: 30 January 2019
© Springer Nature America, Inc. 2019

Abstract

Objective Evidence supports an inverse association between vitamin D and bacterial vaginosis (BV) during pregnancy. Furthermore, both the vaginal microbiome and vitamin D status correlate with pregnancy outcome. Women of African ancestry are more likely to experience BV, to be vitamin D deficient, and to have certain pregnancy complications. We investigated the association between vitamin D status and the vaginal microbiome.

Study design Subjects were assigned to a treatment (4400 IU) or a control group (400 IU vitamin D daily), sampled three times during pregnancy, and vaginal 16S rRNA gene taxonomic profiles and plasma 25-hydroxyvitamin D [25(OH)D] concentrations were examined.

Result Gestational age and ethnicity were significantly associated with the microbiome. *Megasphaera* correlated negatively ($p = 0.0187$) with 25(OH)D among women of African ancestry. Among controls, women of European ancestry exhibited a positive correlation between plasma 25(OH)D and *L. crispatus* abundance.

Conclusion Certain vaginal bacteria are associated with plasma 25(OH)D concentration.

Introduction

The composition of the vaginal microflora can significantly impact both reproductive and neonatal health. Vaginal lactobacilli, through the production of lactic acid, create a

vaginal environment characterized by a low pH, and this, in combination with bacteriocins and possibly other components, inhibits the colonization and growth of potentially pathogenic microorganisms and reduces the phylogenetic diversity of the vaginal microbiome. In a state of dysbiosis termed bacterial vaginosis (BV), the lactobacilli are depleted and replaced with a polymicrobial, anaerobic microflora that includes *Gardnerella vaginalis*, *Atopobium vaginae*, *Sneathia*, *Prevotella spp*, *Megasphaera*, and others. Although BV is frequently asymptomatic, this dysbiotic state is significantly associated with other clinical

These authors contributed equally: G.A. Buck, C.L. Wagner

Supplementary information The online version of this article (<https://doi.org/10.1038/s41372-019-0343-8>) contains supplementary material, which is available to authorized users.

✉ Gregory A. Buck
gregory.buck@vcuhealth.org

✉ Carol L. Wagner
wagnercl@muscd.edu

¹ Department of Microbiology and Immunology, Virginia Commonwealth University, Richmond, VA, USA

² Center for the Study of Biological Complexity, Virginia Commonwealth University, Richmond, VA, USA

³ Department of Statistical Sciences and Operations Research, Virginia Commonwealth University, Richmond, VA, USA

⁴ Institute of Metabolism and Systems Research, University of Birmingham, Edgbaston, Birmingham, UK

⁵ Division of Neonatology, Shawn Jenkins Children's Hospital, Medical University of South Carolina Children's Hospital, Charleston, SC, USA

⁶ Department of Obstetrics and Gynecology, Johns Hopkins University, Baltimore, MD, USA

⁷ Department of Obstetrics and Gynecology, Virginia Commonwealth University, Richmond, VA, USA

complications including pelvic inflammatory disease, infertility, and spontaneous abortion. When BV occurs during pregnancy, it is associated with a more than twofold increased risk for preterm birth. This is important because preterm birth accounts for as much as 70% of neonatal mortality, 75% of neonatal morbidity, and nearly 50% of long-term neurologic sequelae. A significant portion of moderate and late preterm births and the majority of very preterm births (<32 weeks) may be attributable to infection and subsequent inflammation and recent microbiome analyses reveal an association between term birth and abundance of healthy lactobacilli and an association between preterm birth and BV-associated bacterial taxa [1, 2].

Numerous studies support the important role for sufficient serum or plasma concentrations of 25-hydroxyvitamin D (25(OH)D) during pregnancy in preventing negative outcomes [3–10]. Although some studies failed to detect an association, possibly because vitamin D deficiency is low among certain populations, many others and a recent meta-analysis suggest that vitamin D deficiency increases the rate of preterm birth, particularly early preterm birth [3]. It plays roles in maintenance of maternal and fetal calcium homeostasis, which is linked to skeletal integrity, in general fetal growth and development, and in the regulation of immune function [11, 12]. There also appears to be a relationship between vitamin D and BV status during pregnancy. Multiple reports describe higher rates of BV among women with insufficient 25(OH)D concentrations (often defined as <15 ng/mL or <37.5 nmol/L) [13–15]. However, studies in non-pregnant women have yielded conflicting results. One randomized controlled trial that investigated the effectiveness of vitamin D supplementation in conjunction with metronidazole in eliminating BV in non-pregnant women did not detect a positive effect of vitamin D [16] while another study found that vitamin D supplementation was effective in eliminating BV [17].

Race/ethnicity has significant population-level impacts on vitamin D status, BV status, and pregnancy outcomes. Vitamin D deficiency is more prevalent among people with limited sun exposure and among people with dark skin. Children of African ancestry are more likely than those of European ancestry to experience hypocalcemic seizures and to develop rickets [18]. Women of African ancestry are also twice as likely to receive a clinical diagnosis of BV, and analyses of the vaginal microbiota reveal that they are more likely to be colonized by certain BV-associated bacteria [19]. Furthermore, African American women (though not African-born women) are nearly twice as likely to give birth preterm (<37 weeks' gestation) and more than twice as likely to give birth early preterm (<34 weeks) [20, 21] and vitamin D deficiency has been linked to preterm birth and small-for-gestational-age infants born to women of African ancestry [22, 23]. There is an urgent need to understand and address these important health disparities and an important

first step in understanding whether or not the three parameters, vitamin D status, vaginal microbiome, and preterm birth, are linked. As a component of a prospective vitamin D supplementation trial in pregnant women funded by the W. F. F. Kellogg Foundation, referred to as the Kellogg Study, we analyzed a cohort of 230 healthy women and investigated the relationship between plasma 25(OH)D concentrations, the vaginal microbiome, and preterm birth.

Materials and methods

Participant recruitment

This study was approved by the Medical University of South Carolina Institutional Review Board for Human Research (PRO 00020570), registered at clinicaltrials.gov as NCT 01932788, and conducted from 1 January 2013 to 30 April 2018 at the Medical University of South Carolina (Charleston, South Carolina). All subjects gave written, informed consent. Women of diverse racial/ethnic backgrounds presenting to their obstetrician or midwife at the Medical University of SC (MUSC), Charleston, SC obstetrical facilities within the first 14 weeks after last menstrual period (LMP) with confirmation of a singleton pregnancy were eligible for enrollment in the Kellogg Study. Obstetrical dating was confirmed by early ultrasound by an MUSC obstetrician at the first prenatal visit. In those women with greater than a 2-week discrepancy between dating by LMP and ultrasound, the ultrasound dating was used. Mothers with pre-existing calcium, uncontrolled thyroid disease, parathyroid conditions, or who required chronic diuretic or cardiac medication therapy including calcium channel blockers were excluded. Mothers with pre-existing sickle cell disease (not trait only), sarcoidosis, Crohn's disease, or ulcerative colitis were also ineligible to participate in the study. In addition, because of the potentially confounding effect of multiple fetuses, mothers with multiple gestations as determined by obstetrical ultrasound were not eligible for participation in the study. Women who met the entry criteria were approached by the Kellogg Study team for enrollment in the study following their written, informed consent. A subgroup of approximately 100 subjects with known diabetes, hypertension, human immunodeficiency virus (HIV), or morbid obesity (body mass index > 49) did participate but were subsequently excluded from analysis in the present study as the present study's focus was on healthy women at the start of pregnancy.

The women were randomized into one of two groups: (1) group A: 400 IU vitamin D₃/day—standard dose treatment of placebo (0 IU vitamin D₃ gummy) plus prenatal vitamin (400 IU/day); or (2) group B: 4400 IU/day (4000 IU/2

gummies/day + 400 IU/day in prenatal). A sample size of 240 subjects, 120 in each arm, was calculated to be sufficient to detect a clinically important difference of 0.3 between groups in achieving a total circulating 25(OH)D concentration of at least 40 ng/mL assuming a standard deviation of 10 using a two-tailed *t*-test of difference between means with 80% power and a 5% level of significance. Considering a dropout rate of 10% the sample size required is 264 (132 per group). To attain equal group numbers and balanced numbers by racial/ethnic group (African American, Caucasian and Hispanic, which represent the predominant groups delivering at MUSC), a stratified block randomization was used. Adherence to treatment was assessed by monthly pill count and women with an adherence rate of <60% predicted pill intake for two consecutive months exited the study ($n = 60$: 7 white/Caucasian; 18 Hispanic; and 35 black/African American women). Because of the effect of adherence to treatment on vitamin D status [24], the biomarker—25(OH)D was used as an additional measure of maternal vitamin D status.

Sampling and sample processing

Samples were self-obtained using CultureSwab EZ (Becton Dickinson, Franklin Lakes, NJ) from the mid-vaginal wall at visit 1 (baseline; 7.6–15.1 weeks), visit 4 (21.5–28.3 weeks gestation), and visit 7 (31.6–40.4 weeks). The swab handles were broken off and swabs were stored in microtubes at -80°C until they were transported from MUSC to VCU. DNA was extracted from the swabs using the Powersoil[®] kit (MoBio). The swabs were swirled directly in the Powerbead tubes supplied with the kit and processing was according to the manufacturer's instructions. Nugent score was calculated and assigned by a single, blinded obstetrician by assessment of presence of Gram-positive rods, small Gram-variable rods, and curved Gram-variable rods per high power field as described [25].

Measurement of total circulating 25(OH)D

Whole blood samples were obtained, placed in EDTA-treated vacutainers, and centrifuged to achieve plasma separation. The plasma was stored at -80°C until measurement for total circulating 25(OH)D, which was measured using a commercially available radioimmunoassay (Diasorin, Stillwater, MN).

Maternal sociodemographic and health characteristics

Baseline maternal sociodemographic and health data were obtained at the first study visit. During each subsequent study visit, maternal health status was obtained by

utilizing a validated pregnancy health status questionnaire used in two prior studies by this group. Pregnancy complications were obtained by electronic medical record and patient records after obtaining written informed consent, and utilized definitions of preterm labor, preterm birth (delivery <37 weeks), and significant preterm birth (<34 weeks at delivery), preeclampsia as defined by the clinical team using American College of Obstetricians and Gynecologists (ACOG) guidelines, BV, abnormal glucose screening, and gestational diabetes, as previously defined by this group [26]. Sexually transmitted diseases were categorized as trichomoniasis, chlamydia, herpes simplex virus, gonorrhea, and human papilloma virus. BV was diagnosed by a single obstetrician who was blinded to treatment using ACOG guidelines [27].

16S rRNA gene survey

The V1–V3 hypervariable regions of the bacterial 16S rRNA gene was PCR amplified using barcoded primers [28–30]. Briefly, for each reaction, 2 μL of extracted DNA was combined with 12.5 μL 2X Phusion Hot Start II High-Fidelity PCR Master Mix (Thermo Fisher Scientific, Waltham, MA), 3% dimethyl sulfoxide, and 100 nM each of forward and reverse primers. The primers included an Illumina linker adaptor (used for binding and sequencing), a unique index sequence followed immediately by a variable sequence spacer (0–6 bases), and 16S rRNA gene primers (see S1 Appendix). The forward primer was a mixture (4:1) of primers Fwd-P1 (5'-CCATCTCATCCCTGCGTGTCTCC GACTCAG BBBBAGAGTTYGATYMTGGCTYAG) and Fwd-P2 (5'-CCATCTCATCCCTGCGTGTCTCCGACT CAG BBBBAGARTTTTGATCYTGGTTCAG). The reverse primer was Rev1B (5'-CCTATCCCCTGTGTGC CTTGGCAGTCTCAG ATTACCGCGGCTGCTGG). The PCR was carried out in a 25 μL reaction in a Thermal Cycler (Applied BioSystem GeneAmp PCR system 9700) with the following parameters: initial denaturation at 98°C for 30 s, followed by 30 cycles of 98°C for 15 s, 58°C for 15 s, and 72°C for 15 s with a final extension at 72°C for 1 min. The PCR was performed in 96-well format with two PCR controls, a negative water control, and a positive MOMS-PI Mock Community control. All amplicons were quantified using Picogreen (Invitrogen/Thermo Scientific) and a spectrofluorimeter (Biotek). Equal amounts of amplicon were combined followed by removal of unincorporated primers, salts, and enzymes using Agencourt AMPure XP beads. The DNA concentration of this concentrated pool was verified by quantitative PCR using the KAPA Library Quantification Kit for Illumina platform using Thermo Fisher Scientific ViiA[™] 7 Real-Time PCR System. The library pool was diluted and denatured according to the Illumina MiSeq library preparation guide.

The sequencing run was conducted on the Illumina MiSeq sequencer using the 2 × 300 PE reagent kit 3.

Bioinformatics analysis

Sequence reads were de-multiplexed using an in-house python script. The raw overlapping paired-end sequence data were merged and quality-filtered using MeFiT [31] with *meep* (maximum expected error rate) cutoff of 1.0. Non-overlapping high-quality paired-end reads were retained for downstream analysis by linking them with 15 *N*'s. Samples with <1000 high-quality reads were removed. Species-level taxonomic assignments were performed using STIRRUPS [32], employing the USEARCH algorithm [33] combined with a curated vaginal 16S rRNA gene database. All below-threshold (<97% identity) assignments were removed. Relative abundance of each species was calculated as the proportion of total classified reads at >97% identity.

Spurious taxonomic assignments, i.e. relative abundance <0.01%, were removed from each sample's microbial profile and relative abundances were re-normalized. For downstream statistical analysis (diversity measures and linear discriminant analysis effect size (LEfSe) analysis), species present in <5% samples were not taken into consideration.

Statistical analysis

The predominant taxon in a sample refers to the taxon for which the largest number of reads were assigned taxonomic classification with confidence (i.e. the highest percentage of reads in the sample). Vaginal 16S rRNA gene profiles were assigned to community state types/vagitypes based on the taxon with the largest proportion of reads. Samples where the largest proportion was <30% were not assigned a vagitype. This "predominant taxon" rule has been shown to exhibit over 90% agreement with clustering-based methods across a variety of vaginal microbiome datasets [34], yet is not population- or dataset-dependent and is therefore more conducive to use in a clinical setting.

Within-sample alpha diversity was measured using the Shannon and inverse Simpson's index. Pairwise Bray-Curtis dissimilarity distances were calculated for between-sample beta diversity comparisons. Diversity measures were calculated using the *vegan* R package. Principal coordinates analysis on dissimilarity distances was performed using the *phyloseq* R package.

For each race/ethnicity, a linear mixed effect model with random subject effect (to account for the longitudinal nature of the data) was fit to relate 25(OH)D concentration as a function of treatment group, gestational age, bacteria relative abundances from a subject's microbiome profile (*A. vaginalis*, *G. vaginalis*, BVAB1, *Lactobacillus crispatus*,

Lactobacillus gasseri, *Lactobacillus iners*, and *Megasphaera*), as well as the interaction of treatment with gestational age and relative abundances. Interaction terms were included to determine if the effect of gestational age and/or taxa relative abundances on 25(OH)D depend on treatment group. Given the significant association between education level and treatment/control group (Table 1), education level was also included as a covariate. All linear mixed effect models were fit using JMP software (version 14).

LEfSe applies a Kruskal-Wallis rank sum test for each bacterium, then uses linear discriminant analysis to estimate effect size. The effect size is the contribution of a variable to the ability to distinguish two different groups.

Results

Subject characteristics

In all, 402 women were consented and enrolled in the Kellogg Study. Of these, 387 were randomized and received supplement: 191 received 400 IU (control group), of whom, 142 were followed through to delivery; 196 received 4400 IU (treatment group), of whom, 155 were followed through to delivery. There were 19 miscarriages (10 in black/African American women and 9 in Hispanic women). The concentration of 25(OH)D was determined from blood samples collected monthly starting at 10–14 weeks of gestation through delivery. Women ($n = 108$) exited the study due to loss of interest, moving away, and nonadherence to protocol ($n = 60$). Three women were enrolled in the study for two separate pregnancies; the results from the second pregnancy was not included in the analyses, but all 3 women delivered at term gestation. Women who exited the Kellogg Study were more likely to be black/African American ($p < 0.0001$), had less than college education ($p = 0.005$), and were less likely to be employed full time ($p = 0.03$); more likely to have Medicaid/self-pay insurance ($p < 0.0001$), higher gravidity ($p = 0.008$) and parity (0.0015); and lower 25(OH)D concentration at baseline ($p = 0.0003$), but did not differ by body mass index, season, dietary vitamin D intake, or serum calcium or phosphorus concentrations.

While a circulating 25(OH)D concentration was obtained for each participant on a monthly basis, vaginal swab samples were obtained in women who chose to provide a sample, and in some cases, women were late to be seen for their appointments, and chose not to provide a vaginal swab. This resulted in fewer vaginal swabs than blood samples, and hence the discrepancy in numbers. For the purpose of this nested study, we analyzed circulating 25 (OH)D concentrations and vaginal swab samples that were obtained at visit 1 (baseline; 7.6–15.1 weeks' gestation; $n =$

Table 1 Baseline demographics

Baseline maternal characteristic	All subjects	Controls (400 IU)	Treated (4000 IU)	<i>p</i> -Value
Subjects	236	112	124	
Age in years, mean (range)	29 (18–42)	29 (18–42)	29 (20–41)	0.642 ^a
BMI at baseline visit 1, mean	33.19	30.33	35.79	0.494 ^a
Race/ethnicity, <i>N</i> (%)				0.390 ^b
American Indian	2 (1%)	1 (1%)	1 (1%)	
African American	83 (35%)	39 (35%)	44 (35%)	
Hispanic	61 (26%)	34 (30%)	27 (22%)	
White/Caucasian	90 (38%)	38 (34%)	52 (42%)	
Gravidity, median (range)	2 (1–14)	2 (1–14)	2 (1–6)	
Parity, median (range)	1 (0–4)	1 (0–4)	1 (0–3)	
Education, <i>N</i> (%)				
<High school	25 (10%)	17 (15%)	8 (6%)	
High school	56 (24%)	31 (28%)	25 (20%)	0.018 ^b
College	155 (66%)	64 (57%)	91 (74%)	
Insurance status, <i>N</i> (%)				
Private insurance	105 (44%)	44 (39%)	61 (49%)	
Medicaid	84 (36%)	44 (39%)	40 (32%)	0.312 ^b
Self-pay	47 (20%)	24 (22%)	23 (19%)	
Mean gestational age at delivery (weeks, days)	38.90	39.0	38.81	0.410 ^a
Number term (≥37 weeks)	217	107	111	
Number preterm (<37, ≥34 weeks)	16	4	12	0.091 ^b
Number early preterm (<34 weeks)	2	1	1	

BMI body mass index

p-Values calculated using ^aWelch's *t*-test or ^bFisher's exact test, as appropriate

177), visit 4 (21.5–28.3 weeks' gestation; *n* = 220), and visit 7 (31.6–40.4 weeks' gestation; *n* = 200) from 112 women in the control group and 124 women in the treatment group. Baseline demographics for the cohort are summarized in Table 1. Subjects were randomly stratified into control and treatment groups in blocks of 10 and, by chance, there was a significant difference in education between the groups. This was accounted for in analyses as appropriate.

Effect of treatment on total circulating 25(OH)D concentration

The treatment group demonstrated a greater increase in 25(OH)D concentrations over the three time points; however, even the control group exhibited a significant increase over the time course of the study (Fig. 1a). Ethnicity was a significant covariate, so the effect of treatment on 25(OH)D concentration was also examined within each ethnic group (Fig. 1b–d). Women of European ancestry and Hispanic women in the control group exhibited a significant increase in 25(OH)D but women of African ancestry in the control

group did not. All ethnic groups in the treatment group displayed a significant increase in 25(OH)D over the three time points. Gestational age was a significant covariate; however, a linear mixed effect model indicated that the effect of treatment on 25(OH)D concentrations remained significant for each of the three groups (*p* < 0.0001 for African ancestry, *p* = 0.0127 for European ancestry, and *p* < 0.0001 for Hispanic).

Association between vitamin D sufficiency and the vaginal microbiome

DNA from vaginal swab samples was analyzed by 16S rRNA gene survey. 16S amplicon sequences were classified against a custom database of vaginally relevant species, using STIRRUPS [32], and relative abundance was computed as a percentage of reads classified. Taxonomic assignments below 0.01% relative abundance were removed and the proportions were re-normalized. Furthermore, bacterial species not present in at least 5% of samples were removed. This protects against features with small mean and trivially large coefficient of variance.




Fig. 1 Total circulating 25-hydroxyvitamin D (25(OH)D) concentrations increase over pregnancy. Boxplots of 25(OH)D concentration in plasma collected longitudinally over pregnancy. Whiskers extend to the highest/lowest value within 1.5 times the interquartile range and outliers beyond the whiskers are plotted as points. Significant

differences in 25(OH)D between visits are indicated for **a** all participants, **b** women of African ancestry, **c** women of European ancestry, and **d** Hispanic women. Symbols indicating statistical significance—ns: $p > 0.05$; * $p \leq 0.05$; ** $p \leq 0.01$; **** $p \leq 0.0001$. Figure was prepared using *ggpubr* [61] R package

Vaginal 16S rRNA gene profiles from all samples were grouped based on total circulating plasma 25(OH)D concentrations of either <30 or >40 ng/mL. There was no significant association between 25(OH)D concentration and alpha diversity as measured by Shannon index or Inverse Simpson Index (Supplementary figure S1a). Principal coordinates analysis performed on the Bray-Curtis dissimilarity distances between samples did not reveal any patterns between plasma 25(OH)D concentration and bacterial species (supplementary figure S1b). When the samples were grouped by dominant taxon, it was apparent, as expected, that women with vaginal microbiome dominated by lactobacilli had lower Nugent scores, overall (Fig. 2). Furthermore, it appeared that women with >40 ng/mL 25(OH)D at the time of sampling had vaginal microbiomes with greater abundance of lactobacilli relative to women with <30 ng/mL 25(OH)D (Fig. 2a, b). The microbiomes in this group were 1.6 times, 2.3 times, and 1.5 times as likely to be dominated by the healthy *Lactobacillus* species *L. crispatus*, *L. jensenii*, and *L. gasseri* (respectively), and 0.2 times as likely

to be dominated by the unhealthy BV-associated species *G. vaginalis* or BVAB1 (Table 2). When separated by ethnicity, the effect appeared to be more pronounced among women of African ancestry (Fig. 3).

We analyzed the abundance of bacterial taxa by LEfSe to determine whether any were associated with 25(OH)D status. Among samples from women with >40 ng/mL 25(OH)D, *L. crispatus*, *L. gasseri*, and *Bifidobacterium* species were more abundant (Fig. 4). In addition to these lactic acid-producing taxa, additional taxa that are not associated with vaginal health, including *A. vaginae*, were more abundant in this group. Among samples from women with <30 ng/mL 25(OH)D, *G. vaginalis*, BVAB1, TM7_OTU_H1, *Prevotella*, BVAB1, and *Sneathia* were more abundant.

Relationship between *G. vaginalis*, plasma 25(OH)D, ethnicity, and gestational age

Because of its strong association with BV and because it appeared to be negatively associated with vitamin D sufficiency, we examined the abundance of *G. vaginalis* over the




Fig. 2 Vaginal microbial taxa differ between women with plasma 25-hydroxyvitamin D (25(OH)D) <30 or >40 ng/mL. **a, b** Stacked bar plots showing vaginal microbial community profiles from three visits from each of the 236 women in the cohort. The profiles are grouped by

the most abundant species and samples within each community group are clustered on Bray distances using ward method. The microbial profiles are annotated by Nugent score. The distances were calculated using *vegan* R package and the figure was prepared using *ggplot2*

Table 2 Prevalence of communities in 25(OH)D-deficient and -sufficient subjects

	<i>L. crispatus</i>	<i>L. jensenii</i>	<i>L. gasseri</i>	<i>L. iners</i>	<i>G. vaginalis</i>	BVAB1	Other	No type	Total
<30 ng/mL	43 (19%)	8 (3%)	15 (6%)	103 (44%)	25 (11%)	12 (5%)	16 (7%)	10 (4%)	232 (100%)
>40 ng/mL	68 (30%)	17 (7%)	21 (9%)	94 (41%)	4 (2%)	2 (1%)	16 (7%)	8 (3%)	230 (100%)

Number of samples and (percent of total) dominated by one of the named bacterial taxa, a rare taxon “Other”, or not dominated by a single taxon “No type”

25(OH)D 25-hydroxyvitamin D

course of pregnancy more carefully. We examined a subset of the women of African ancestry who had a relative abundance of at least 1% *G. vaginalis* at their first visit. Among the controls within this subset, there was not a significant increase in total circulating 25(OH)D between visit 1 and visit 7 (Wilcoxon, $p = 0.14$) but there was a significant increase in the treated subjects (Wilcoxon $p < 0.0001$) (Fig. 5a). There was a decrease in *G. vaginalis* abundance between visits 1 and 7 in both the control and treatment group ($p = 0.0029$ and 0.0153 , respectively) (Fig. 5b).

Collinearity between plasma 25(OH)D and gestational age

Because of the observed decrease in the BV-associated taxon *G. vaginalis* over pregnancy, even among the control group, we examined baseline (before treatment) 25(OH)D concentrations and Nugent scores and found that women of African ancestry with 25(OH)D <20 ng/mL at baseline had significantly higher Nugent score than women with 25(OH)D >40 ng/mL. We also developed a model to incorporate the mixed effects of treatment group and gestational age,

Fig. 3 The association between 25-hydroxyvitamin D (25(OH)D) and microbiome differs among ethnic groups. Stacked bar plots showing vaginal microbial community profiles from three visits from each of the 236 women in the cohort grouped by ethnicity and 25(OH)D status





Fig. 4 Statistical association analysis using linear discriminant analysis effect size (LEfSe). Bacterial species with significant differential abundance between women with plasma 25-hydroxyvitamin D (25(OH)D) concentrations >40 or <30 ng/mL were identified using LEfSe. Features with linear discriminant analysis (LDA) score > 3.0 are shown in the figure, with bacterial species associated with plasma 25(OH)D concentrations >40 ng/mL shown in green, while those associated with <30 ng/mL are shown in red



All Participants with *Gardnerella vaginalis* relative abundance > 1% at visit1




Fig. 5 *G. vaginalis* abundance decreases over pregnancy irrespective of 25-hydroxyvitamin D (25(OH)D) concentration. Boxplots showing plasma 25(OH)D concentration (left panel) and *G. vaginalis* abundance (right panel) for all subjects with *G. vaginalis* abundance >1% at visit 1. Data points from same subject at visit 1 and visit 7 are

connected by gray line. There is a significant increase in 25(OH)D among the treatment group and significant decrease in *G. vaginalis* abundance in both control and treatment group. The figure was prepared using *ggpubr* R package

and we analyzed the different ethnic groups separately. Because education differed between the control and treatment groups, education was accounted for in the model. When treatment group and gestational age were fixed, both treatment and gestational age significantly correlated with 25(OH)D concentrations, but the associations between many of the taxa and 25(OH)D were no longer significant, suggesting that covariation of abundance levels with gestational age, treatment group, and/or ethnicity exerted an effect. Despite the covariation, there remained a significant correlation ($p = 0.0088$) between *Megasphaera* species and 25(OH)D concentrations among women of African ancestry insofar as women with higher 25(OH)D had less *Megasphaera*. Among

women of European ancestry, a significant interaction between *L. crispatus* and treatment ($p = 0.0123$) was observed and suggests that the association of *L. crispatus* with 25(OH)D depends on treatment/control. In particular, among women of European ancestry in the control group, but not the treatment group, total circulating 25(OH)D concentrations correlated positively with abundance of *L. crispatus*.

Discussion

The hypothesis driving this study was that vitamin D status is associated with the vaginal microbiome. Vitamin D could

affect the vaginal microbiome through a number of mechanisms. First, it could promote the barrier integrity of the vaginal epithelium. Vitamin D was shown to upregulate genes encoding epithelial cell junction proteins and stimulated proliferation of the vaginal epithelium [35, 36]. Furthermore, two studies found that postmenopausal women receiving vitamin D treatment had increased numbers of superficial vaginal epithelial cells compared to women not receiving treatment, and that this increase was accompanied by decreases in vaginal pH [37, 38]. Together these studies suggest that vitamin D may promote vaginal epithelial cell growth, differentiation, and function, which may increase vaginal thickness to prevent atrophy and improve barrier function. Second, adequate vitamin D during pregnancy may impact the vaginal environment so that it is more supportive of beneficial flora. It is well established that insulin stimulates glycogen synthesis [39], and more recently it has been proposed that vitamin D induces insulin synthesis [40, 41]. Vitamin D was also shown to increase the phosphorylation and inactivation of glycogen synthase kinase, an inhibitor of glycogen synthesis, in adipose tissue [42]. Therefore, there is also a possibility that vitamin D sufficiency alters glucose homeostasis in the vagina to promote increased glycogen deposition. Increased vaginal levels of free glycogen positively correlate with *Lactobacillus* relative abundance [43], and so if vitamin D is indeed promoting vaginal glycogen production, the accompanying increase in lactobacilli and decrease in pH may result in the decreased abundance of pathogenic and BV-associated organisms. A third possibility is that vitamin D influences abundance of BV-associated bacteria through its effects on immune function. Previous studies have shown that vitamin D treatment induces the expression of the LL-37 antimicrobial peptide in keratinocytes, neutrophils, monocytes, and bladder and gingival epithelial cells as well as β 2 defensin in keratinocytes, neutrophils, and monocytes [44–46], as well as potentiates the secretion of interleukin-8 and CXCL10 from airway epithelial cells [47]. These effects could promote leukocyte recruitment to the vaginal epithelium and modulate the antimicrobial response of leukocytes upon arrival to the tissue [48].

In this study, we analyzed a sub-cohort from a randomized, placebo-controlled clinical trial of vitamin D supplementation of pregnant women who were enrolled during the first trimester of pregnancy and followed until delivery. We analyzed 16S rRNA gene survey data to determine whether plasma 25(OH)D concentrations were associated with the vaginal microbiome. Women in both the control (400 IU vitamin D daily) and treatment groups (4400 IU vitamin D daily) displayed higher circulating 25(OH)D concentrations with increasing gestational age. While initial analyses suggested a substantive association between 25(OH)D and the vaginal microbiome, we found that

gestational age was a strong covariate. While multiple studies have shown that the abundances of certain BV-associated bacterial taxa decrease during pregnancy [49, 50], a correlation with gestational age has not been demonstrated and a report of a longitudinal study suggested that the vaginal microbiome is remarkably constant over the course of pregnancy [51]. In our study, however, *G. vaginalis* decreased significantly in abundance between visit 1 (7.6–15.1 weeks) and visit 7 (31.6–40.4 weeks) (Fig. 5).

A model that took into account the covariation between the vaginal microbiome and gestational age, found only three taxa that were associated with 25(OH)D concentrations, and these associations were dependent upon ethnicity. Specifically, among women of African ancestry, there was a negative correlation between 25(OH)D and abundance of *Megasphaera* spp. The majority of reads that were categorized with the genus *Megasphaera* were *Megasphaera* type 1, which we have found in another study (Glascock, Fettweis, manuscript in preparation) to be the most common vaginal type. Detection of *Megasphaera* by PCR has been shown to accurately predict of BV [52] and *Megasphaera* is associated with bacteria vaginosis and spontaneous preterm delivery [53–55]. *Megasphaera* has also been linked to genital tract inflammation [56] and HIV acquisition among African women [57]. Among women of European ancestry in the control group, women with higher 25(OH)D concentrations were more likely to have higher abundance of *L. crispatus*. *L. crispatus* is a healthy H₂O₂-producing lactobacillus species that is associated with decreased risk for developing BV and decreased risk for HIV acquisition [58, 59]. Thus, although only two taxa were significantly associated with 25(OH)D, these results suggest that vitamin D could have a positive impact on the vaginal microbiome.

A limitation of this study is that the cohort contained few women who had profound vitamin D deficiency (defined as 25(OH)D <12 ng/mL). In our prior National Institute of Child Health and Human Development vitamin D pregnancy study involving 350 women randomized to three treatment groups (400, 2000, and 4000 IU vitamin D₃/day) [60], even the control group showed an increase in 25(OH)D over the course of pregnancy, especially in white/Caucasian women. Most women are not taking a prenatal vitamin or a multivitamin at the start of pregnancy and since most women have limited sunlight exposure or use sunscreen limiting their endogenous synthesis of vitamin D, and because the average American woman is provided ~200 IU vitamin D/day from her diet [60], the addition of 400 IU/day results in a slight but measurable increase in 25(OH)D over time.

Another potential limitation of the study is the loss of subjects during the study period. There were 108 women who exited the study, 60 due to nonadherence to protocol. The women who exited the study tended to be of lower

socioeconomic status, were black/African American, less educated, more likely to be of higher gravidity and parity, and with a lower baseline 25(OH)D concentration than those women who completed the study. The loss of these subjects in the study could have created potential bias in the results.

It is possible that within the range of total circulating 25(OH)D concentration among women in the cohort, there is little impact on the vaginal microbiome, but that severe vitamin D deficiency has a greater effect. In this study, plasma 25(OH)D was the only marker of vitamin D status that was measured; however, maternal 1,25D and other vitamin D metabolites such as 24,25(OH)₂D and the 3-epi-25(OH)D could also be important factors. Because the subjects in this cohort were pregnant, plasma 1,25D concentration would be expected to be higher and could influence immune function and the vaginal microbiota. Another limitation was that the relatively small cohort size may not have provided the study with adequate power to detect significant associations between 25(OH)D concentrations and bacterial taxa, particularly in light of other covariates (gestational age, ethnicity, and random treatment group assignment). While all participants in the study provided a blood sample to measure total circulating 25(OH)D, only a subset of those women provided a vaginal sample at the three time points, which further decreased the sample size. The vaginal microbiome is a complex parameter and consequently, microbiome studies often require large cohorts to detect biologically relevant results. An additional potential confounder is that women who were diagnosed with BV by Nugent score were prescribed treatment, and while treatment occurred after vaginal swab samples were obtained, it could have influenced the composition of the microbiome at subsequent visits.

Acknowledgements This work was funded in part from a grant from the W. K. Kellogg Foundation and by the South Carolina Clinical & Translational Research (SCTR) Institute, with an academic home at the Medical University of South Carolina, NIH/NCAT Grant number UL1 TR000062. This work was also supported by National Institutes of Health [grant U54 DE023786 “A Multi-omic Analysis of the Vaginal Microbiome during Pregnancy”]. All sequencing and analysis of sequence data were performed in the Genomics Core of the Nucleic Acids Research Facilities at VCU.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Publisher’s note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. Tabatabaei N, Eren AM, Barreiro LB, Yotova V, Dumaine A, Allard C et al. Vaginal microbiome in early pregnancy and subsequent risk of spontaneous preterm birth: a case-control study. *BJOG*. 2018. <https://doi.org/10.1111/1471-0528.15299>.
2. Callahan BJ, DiGiulio DB, Goltsman DSA, Sun CL, Costello EK, Jeganathan P, et al. Replication and refinement of a vaginal microbial signature of preterm birth in two racially distinct cohorts of US women. *Proc Natl Acad Sci USA*. 2017;114:9966–71.
3. Amegah AK, Klever MK, Wagner CL. Maternal vitamin D insufficiency and risk of adverse pregnancy and birth outcomes: a systematic review and meta-analysis of longitudinal studies. *PLoS ONE*. 2017;12:e0173605.
4. Tabatabaei N, Auger N, Herba CM, Wei S, Allard C, Fink GD, et al. Maternal vitamin D insufficiency early in pregnancy is associated with increased risk of preterm birth in ethnic minority women in Canada. *J Nutr*. 2017;147:1145–51.
5. Zhao X, Fang R, Yu R, Chen D, Zhao J, Xiao J. Maternal vitamin D status in the late second trimester and the risk of severe pre-eclampsia in southeastern China. *Nutrients*. 2017;9. <https://doi.org/10.3390/nu9020138>.
6. Wen J, Hong Q, Zhu L, Xu P, Fu Z, Cui X, et al. Association of maternal serum 25-hydroxyvitamin D concentrations in second and third trimester with risk of gestational diabetes and other pregnancy outcomes. *Int J Obes (Lond)*. 2017;41:489–96.
7. Toko EN, Sumba OP, Daud II, Ogolla S, Majiwa M, Krisher JT et al. Maternal vitamin D status and adverse birth outcomes in children from rural Western Kenya. *Nutrients*. 2016;8. <https://doi.org/10.3390/nu8120794>.
8. Kiely ME, Zhang JY, Kinsella M, Khashan AS, Kenny LC. Vitamin D status is associated with uteroplacental dysfunction indicated by pre-eclampsia and small-for-gestational-age birth in a large prospective pregnancy cohort in Ireland with low vitamin D status. *Am J Clin Nutr*. 2016;104:354–61.
9. Miliku K, Vinkhuyzen A, Blanken LM, McGrath JJ, Eyles DW, Burne TH, et al. Maternal vitamin D concentrations during pregnancy, fetal growth patterns, and risks of adverse birth outcomes. *Am J Clin Nutr*. 2016;103:1514–22.
10. Wagner CL, Hollis BW, Kotsa K, Fakhoury H, Karras SN. Vitamin D administration during pregnancy as prevention for pregnancy, neonatal and postnatal complications. *Rev Endocr Metab Disord*. 2017;18:307–22.
11. Weinert LS, Silveiro SP. Maternal-fetal impact of vitamin D deficiency: a critical review. *Matern Child Health J*. 2015;19:94–101.
12. Tamblyn JA, Hewison M, Wagner CL, Bulmer JN, Kilby MD. Immunological role of vitamin D at the maternal-fetal interface. *J Endocrinol*. 2015;224:R107–121.
13. Bodnar LM, Krohn MA, Simhan HN. Maternal vitamin D deficiency is associated with bacterial vaginosis in the first trimester of pregnancy. *J Nutr*. 2009;139:1157–61.
14. Dunlop AL, Taylor RN, Tangpricha V, Fortunato S, Menon R. Maternal vitamin D, folate, and polyunsaturated fatty acid status and bacterial vaginosis during pregnancy. *Infect Dis Obstet Gynecol*. 2011;2011:216217.
15. Hensel KJ, Randis TM, Gelber SE, Ratner AJ. Pregnancy-specific association of vitamin D deficiency and bacterial vaginosis. *Am J Obstet Gynecol*. 2011;204:41.e1–9.
16. Turner AN, Carr Reese P, Fields KS, Anderson J, Ervin M, Davis JA, et al. A blinded, randomized controlled trial of high-dose vitamin D supplementation to reduce recurrence of bacterial vaginosis. *Am J Obstet Gynecol*. 2014;211:479.e1–479.e13.
17. Taheri M, Baheiraei A, Foroushani AR, Nikmanesh B, Modarres M. Treatment of vitamin D deficiency is an effective method in the elimination of asymptomatic bacterial vaginosis: a placebo-controlled randomized clinical trial. *Indian J Med Res*. 2015;141:799–806.

18. O'Callaghan KM, Kiely ME. Ethnic disparities in the dietary requirement for vitamin D during pregnancy: considerations for nutrition policy and research. *Proc Nutr Soc.* 2018;77:164–73.
19. Fettweis JM, Brooks JP, Serrano MG, Sheth NU, Girerd PH, Edwards DJ, et al. Differences in vaginal microbiome in African American women versus women of European ancestry. *Microbiol Read Engl.* 2014;160:2272–82.
20. Oliver EA, Klebanoff M, Yossef-Salameh L, Oza-Frank R, Moosavinasab S, Reagan P, et al. Preterm birth and gestational length in four race-nativity groups, including Somali Americans. *Obstet Gynecol.* 2018;131:281–9.
21. Hamilton BE, Martin JA, Osterman MJK, Curtin SC, Matthews TJ. Births: Final Data for 2014. *Natl Vital Stat Rep.* 2015;64:1–64.
22. Seto TL, Tabangin ME, Langdon G, Mangeot C, Dawodu A, Steinhoff M, et al. Racial disparities in cord blood vitamin D levels and its association with small-for-gestational-age infants. *J Perinatol.* 2016;36:623–8.
23. Wagner CL, Baggerly C, McDonnell S, Baggerly KA, French CB, Baggerly L, et al. Post-hoc analysis of vitamin D status and reduced risk of preterm birth in two vitamin D pregnancy cohorts compared with South Carolina March of Dimes 2009–11 rates. *J Steroid Biochem Mol Biol.* 2016;155:245–51.
24. Abercrombie M, Shary J, Ebeling M, Hollis B, Wagner C. Analysis of the NICHD vitamin D pregnancy cohort on a per-protocol vs. intent-to-treat basis: the effect of adherence on trial results. *J Nutr Food Sci.* 2018;8:696.
25. Nugent RP, Krohn MA, Hillier SL. Reliability of diagnosing bacterial vaginosis is improved by a standardized method of gram stain interpretation. *J Clin Microbiol.* 1991;29:297–301.
26. Hollis BW, Wagner CL. Vitamin D supplementation during pregnancy: improvements in birth outcomes and complications through direct genomic alteration. *Mol Cell Endocrinol.* 2017;453:113–30.
27. ACOG Committee on Practice Bulletins—Gynecology. ACOG Practice Bulletin. Clinical management guidelines for obstetrician-gynecologists, Number 72, May 2006: Vaginitis. *Obstet Gynecol.* 2006;107:1195–206.
28. Fadrosch DW, Ma B, Gajer P, Sengamaly N, Ott S, Brotman RM, et al. An improved dual-indexing approach for multiplexed 16 S rRNA gene sequencing on the Illumina MiSeq platform. *Microbiome.* 2014;2:6.
29. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl Environ Microbiol.* 2013;79:5112–20.
30. Bartram AK, Lynch MDJ, Stearns JC, Moreno-Hagelsieb G, Neufeld JD. Generation of multimillion-sequence 16 S rRNA gene libraries from complex microbial communities by assembling paired-end Illumina reads. *Appl Environ Microbiol.* 2011;77:3846–52.
31. Parikh HI, Koparde VN, Bradley SP, Buck GA, Sheth NU. MeFIT: merging and filtering tool for illumina paired-end reads for 16 S rRNA amplicon sequencing. *BMC Bioinformatics.* 2016;17:491.
32. Fettweis JM, Serrano MG, Sheth NU, Mayer CM, Glascock AL, Brooks JP, et al. Species-level classification of the vaginal microbiome. *BMC Genomics.* 2012;13(Suppl 8):S17.
33. Edgar RC. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics.* 2010;26:2460–1.
34. Brooks JP, Buck GA, Chen G, Diao L, Edwards DJ, Fettweis JM, et al. Changes in vaginal community state types reflect major shifts in the microbiome. *Microb Ecol Health Dis.* 2017;28:1303265.
35. Gniadecki R, Gajkowska B, Hansen M. 1,25-dihydroxyvitamin D3 stimulates the assembly of adherens junctions in keratinocytes: involvement of protein kinase C. *Endocrinology.* 1997;138:2241–8.
36. Lee A, Lee MR, Lee H-H, Kim Y-S, Kim J-M, Enkhbold T, et al. Vitamin D proliferates vaginal epithelium through RhoA expression in postmenopausal atrophic vagina tissue. *Mol Cells.* 2017;40:677–84.
37. Rad P, Tadayon M, Abbaspour M, Latifi SM, Rashidi I, Delaviz H. The effect of vitamin D on vaginal atrophy in postmenopausal women. *Iran J Nurs Midwifery Res.* 2015;20:211–5.
38. Yildirim B, Kaleli B, Düzcan E, Topuz O. The effects of postmenopausal Vitamin D treatment on vaginal atrophy. *Maturitas.* 2004;49:334–7.
39. Forde JE, Dale TC. Glycogen synthase kinase 3: a key regulator of cellular fate. *Cell Mol Life Sci.* 2007;64:1930–44.
40. Maestro B, Campión J, Dávila N, Calle C. Stimulation by 1,25-dihydroxyvitamin D3 of insulin receptor expression and insulin responsiveness for glucose transport in U-937 human promonocytic cells. *Endocr J.* 2000;47:383–91.
41. Maestro B, Molero S, Bajo S, Dávila N, Calle C. Transcriptional activation of the human insulin receptor gene by 1,25-dihydroxyvitamin D(3). *Cell Biochem Funct.* 2002;20:227–32.
42. Parker L, Levinger I, Mousa A, Howlett K, de Courten B. Plasma 25-hydroxyvitamin D is related to protein signaling involved in glucose homeostasis in a tissue-specific manner. *Nutrients.* 2016;8. <https://doi.org/10.3390/nu8100631>.
43. Mirmonsef P, Hotton AL, Gilbert D, Burgad D, Landay A, Weber KM, et al. Free glycogen in vaginal fluids is associated with Lactobacillus colonization and low vaginal pH. *PLoS ONE.* 2014;9:e102467.
44. Wang T-T, Nestel FP, Bourdeau V, Nagai Y, Wang Q, Liao J, et al. Cutting edge: 1,25-dihydroxyvitamin D3 is a direct inducer of antimicrobial peptide gene expression. *J Immunol.* 2004;173:2909–12.
45. Hertting O, Holm Å, Lühthje P, Brauner H, Dyrdak R, Jonasson AF, et al. Vitamin D induction of the human antimicrobial Peptide cathelicidin in the urinary bladder. *PLoS ONE.* 2010;5:e15580.
46. McMahon L, Schwartz K, Yilmaz O, Brown E, Ryan LK, Diamond G. Vitamin D-mediated induction of innate immunity in gingival epithelial cells. *Infect Immun.* 2011;79:2250–6.
47. Brockman-Schneider RA, Pickles RJ, Gern JE. Effects of vitamin D on airway epithelial cell morphology and rhinovirus replication. *PLoS ONE.* 2014;9:e86755.
48. Zheng Y, Niyonsaba F, Ushio H, Nagaoka I, Ikeda S, Okumura K, et al. Cathelicidin LL-37 induces the generation of reactive oxygen species and release of human alpha-defensins from neutrophils. *Br J Dermatol.* 2007;157:1124–31.
49. Romero R, Hassan SS, Gajer P, Tarca AL, Fadrosch DW, Nikita L, et al. The composition and stability of the vaginal microbiota of normal pregnant women is different from that of non-pregnant women. *Microbiome.* 2014;2:4.
50. Stout MJ, Zhou Y, Wylie KM, Tarr PI, Macones GA, Tuuli MG. Early pregnancy vaginal microbiome trends and preterm birth. *Am J Obstet Gynecol.* 2017;217:356.e1–18.
51. DiGiulio DB, Callahan BJ, McMurdie PJ, Costello EK, Lyell DJ, Robaczewska A, et al. Temporal and spatial variation of the human microbiota during pregnancy. *Proc Natl Acad Sci USA.* 2015;112:11060–5.
52. Fredricks DN, Fiedler TL, Thomas KK, Oakley BB, Marrazzo JM. Targeted PCR for detection of vaginal bacteria associated with bacterial vaginosis. *J Clin Microbiol.* 2007;45:3270–6.
53. Kusters JG, Reuland EA, Bouter S, Koenig P, Dorigo-Zetsma JW. A multiplex real-time PCR assay for routine diagnosis of bacterial vaginosis. *Eur J Clin Microbiol Infect Dis.* 2015;34:1779–85.

54. Nelson DB, Hanlon A, Nachamkin I, Haggerty C, Mastrogiannis DS, Liu C, et al. Early pregnancy changes in bacterial vaginosis-associated bacteria and preterm delivery. *Paediatr Perinat Epidemiol.* 2014;28:88–96.
55. Zozaya-Hinchliffe M, Martin DH, Ferris MJ. Prevalence and abundance of uncultivated *Megasphaera*-like bacteria in the human vaginal environment. *Appl Environ Microbiol.* 2008;74:1656–9.
56. Lennard K, Dabee S, Barnabas SL, Havyarimana E, Blakney A, Jaumdally SZ et al. Microbial composition predicts genital tract inflammation and persistent bacterial vaginosis in South African adolescent females. *Infect Immun.* 2018;86. <https://doi.org/10.1128/IAI.00410-17>.
57. McClelland RS, Lingappa JR, Srinivasan S, Kinuthia J, John-Stewart GC, Jaoko W, et al. Evaluation of the association between the concentrations of key vaginal bacteria and the increased risk of HIV acquisition in African women from five cohorts: a nested case-control study. *Lancet Infect Dis.* 2018;18:554–64.
58. Gajer P, Brotman RM, Bai G, Sakamoto J, Schütte UME, Zhong X, et al. Temporal dynamics of the human vaginal microbiota. *Sci Transl Med.* 2012;4:132ra52.
59. Verstraelen H, Verhelst R, Claeys G, De Backer E, Temmerman M, Vaneechoutte M. Longitudinal analysis of the vaginal microflora in pregnancy suggests that *L. crispatus* promotes the stability of the normal vaginal microflora and that *L. gasseri* and/or *L. iners* are more conducive to the occurrence of abnormal vaginal microflora. *BMC Microbiol.* 2009;9:116.
60. Hollis BW, Johnson D, Hulsey TC, Ebeling M, Wagner CL. Vitamin D supplementation during pregnancy: double-blind, randomized clinical trial of safety and effectiveness. *J Bone Miner Res.* 2011;26:2341–57.
61. Wickham H. *ggplot2 - elegant graphics for data analysis.* Springer. <http://www.springer.com/us/book/9780387981413> (accessed 13 Jul 2016).