



Vitamin D and the epigenome

Irfete S. Fetahu, Julia Höbaus and Enikő Kállay*

Department of Pathophysiology and Allergy Research, Center of Pathophysiology, Infectiology and Immunology, Comprehensive Cancer Center, Medical University of Vienna, Vienna, Austria

Edited by:

Carsten Carlberg, University of Eastern Finland, Finland

Reviewed by:

Patsie Polly, The University of New South Wales, Australia

Moray J. Campbell, Roswell Park Cancer Institute, USA

*Correspondence:

Enikő Kállay, Department of Pathophysiology and Allergy Research, Medical University of Vienna, Währinger Gürtel 18-20, A-1090 Vienna, Austria
e-mail: enikoe.kallay@medunivwien.ac.at

Epigenetic mechanisms play a crucial role in regulating gene expression. The main mechanisms involve methylation of DNA and covalent modifications of histones by methylation, acetylation, phosphorylation, or ubiquitination. The complex interplay of different epigenetic mechanisms is mediated by enzymes acting in the nucleus. Modifications in DNA methylation are performed mainly by DNA methyltransferases (DNMTs) and ten-eleven translocation (TET) proteins, while a plethora of enzymes, such as histone acetyltransferases (HATs), histone deacetylases (HDACs), histone methyltransferases (HMTs), and histone demethylases (HDMs) regulate covalent histone modifications. In many diseases, such as cancer, the epigenetic regulatory system is often disturbed. Vitamin D interacts with the epigenome on multiple levels. Firstly, critical genes in the vitamin D signaling system, such as those coding for vitamin D receptor (VDR) and the enzymes 25-hydroxylase (CYP2R1), 1 α -hydroxylase (CYP27B1), and 24-hydroxylase (CYP24A1) have large CpG islands in their promoter regions and therefore can be silenced by DNA methylation. Secondly, VDR protein physically interacts with coactivator and corepressor proteins, which in turn are in contact with chromatin modifiers, such as HATs, HDACs, HMTs, and with chromatin remodelers. Thirdly, a number of genes encoding for chromatin modifiers and remodelers, such as HDMs of the Jumonji C (JmjC)-domain containing proteins and lysine-specific demethylase (LSD) families are primary targets of VDR and its ligands. Finally, there is evidence that certain VDR ligands have DNA demethylating effects. In this review we will discuss regulation of the vitamin D system by epigenetic modifications and how vitamin D contributes to the maintenance of the epigenome, and evaluate its impact in health and disease.

Keywords: VDR, VDRE, 1,25-dihydroxyvitamin D₃, CYP27B1, CYP24A1, DNA methylation, histone modifications, CpG island

INTRODUCTION

The role of vitamin D in regulating gene expression has become increasingly evident since the discovery of the transcription factor vitamin D receptor (VDR), a member of the steroid nuclear receptor superfamily. The effect of liganded VDR depends on the epigenetic landscape of the target gene. Genome wide analysis in the human leukemia cell line THP-1 showed that VDR binds mainly at loci of open chromatin. Upon treatment with the VDR ligand 1,25-dihydroxyvitamin D₃ (1,25-D₃), chromatin accessibility further increases in more than 30% of these loci (Seuter et al., 2013). The mechanism of action of the liganded VDR is dependent on binding and action of histone acetyltransferases (HATs) and histone methyltransferases (HMTs). It has been shown that co-treatment of cells with 1,25-D₃, and histone deacetylase or DNA methyltransferase inhibitors often have synergistic effects (Pan et al., 2010).

Many common diseases have both genetic and epigenetic components, which communicate in an intricate and multilayered manner. Currently, it is not clear to what extent epigenetic alterations contribute to onset and progress of common diseases, such as cancer. Epigenetics refers to processes that alter gene activity without changing the DNA sequence. They play an important role in regulating key processes during development, including embryonic developmental events, gene imprinting, and

inactivation of chromosome X in females (Bird, 2002; Meissner et al., 2008; Tsai and Baylin, 2011). Maintenance of normal functioning of these biological processes is dependent on the intricate interaction between several epigenetic mechanisms, such as DNA methylation, histone modifications, and non-coding RNAs (Jones and Baylin, 2007). Therefore, at a given promoter the marks arising from DNA methylation and histone modifications determine whether the chromatin is in an open (active) or a closed (repressed) state. Deregulation of the epigenetic mechanisms can lead to aberrant DNA methylation patterns and chromatin architecture, which is a common feature in cancer (Baylin and Jones, 2011; Tsai and Baylin, 2011; Helin and Dhanak, 2013).

EPIGENETIC CHANGES MEDIATED BY THE VITAMIN D RECEPTOR AND ITS LIGANDS

The effect of nutrition on the methylation equilibrium of the genome is already accepted as one of the mechanisms preventing either promoter hyper- or global hypomethylation. Several nutrients are renowned for their impact on DNA methylation, such as folic acid, vitamin B, green tea, and alcohol (Arasaradnam et al., 2008). The effect of vitamin D is currently under debate.

Primary epigenetic effects of vitamin D are linked to histone modifications, mainly acetylation. The VDR/RXR dimer interacts with HATs to induce transcriptional activation (Karlic and Varga,

2011). Several studies have suggested that vitamin D may affect also DNA methylation. A recent study associated severe vitamin D deficiency with methylation changes in leukocyte DNA, although the observed differences were relatively small (Zhu et al., 2013). This study suggested that subjects with vitamin D deficiency were more likely to show reduced synthesis and increased catabolism of active vitamin D. Whether this was the cause of the vitamin D deficiency or the consequence thereof is not clear and needs further studies.

EFFECT OF VITAMIN D ON DNA METHYLATION

DNA methylation is the most extensively studied epigenetic mark (Esteller, 2008). In humans, DNA methylation occurs on cytosines followed by guanine (CpG) (Bird, 1980; Gruenbaum et al., 1981). Regions of DNA enriched in CpG clusters form CpG islands (CGI) (Wang and Leung, 2004). DNA methylation is necessary for regulating and orchestrating key biological processes, including cell cycle, differentiation, as well as genomic imprinting (Feinberg et al., 2002; Reik and Lewis, 2005; Jones and Baylin, 2007). DNA hypermethylation is mainly found in intergenic regions and repetitive genomic sequences to maintain these in a transcriptionally inactive chromatin state (Herman and Baylin, 2003).

DNA methyltransferases (DNMTs) mediate DNA methylation (Robertson, 2005). DNMT1 encodes for a maintenance methyltransferase, whereas DNMT3A/3B encode for de novo methyltransferases, which are pivotal to maintain and establish genomic methylation (Okano et al., 1998, 1999; Jin and Robertson, 2013). However, *in vivo* studies suggest that all three DNMTs might exert both de novo and maintenance functions (Rhee et al., 2000, 2002; Kim et al., 2002; Esteller, 2007a). Recently, a new group of enzymes that induce active demethylation of the DNA was discovered, the ten-eleven translocation (TET) enzyme family, which plays an important role both in development and tumorigenesis (Kriaucionis and Heintz, 2009; Ficz et al., 2011; Williams et al., 2011; Yamaguchi et al., 2012; Hackett et al., 2013).

Alterations in the cancer epigenome are generally associated with loss of global DNA methylation and gain of methylation in specific gene promoters (Ting et al., 2006). Loss of global methylation may lead to chromosomal instability (Eden et al., 2003), loss of imprinting (Cui et al., 2003; Bjornsson et al., 2007), and activation of transposable elements, thereby leading to disturbances in the genome (Bestor, 2005; Esteller, 2008). Conversely, hypermethylation of promoter regions of tumor suppressor genes (Greger et al., 1989; Sakai et al., 1991; Esteller, 2008) leads to loss of expression of key genes affecting pathways involved in maintenance of cellular functions, including cell cycle, apoptosis, and DNA repair (Esteller, 2007b). Several bona fide tumor suppressor genes are silenced by promoter hypermethylation in tumors. For instance, hypermethylation of the promoter of the DNA repair gene *hMLH1* is associated with early stages of endometrial and colon cancer, and microsatellite instability phenotype (Esteller et al., 1999). Epigenetically mediated silencing of cyclin-dependent kinase inhibitor 2A, which is crucial for control of cell cycle has been reported in several cancers (Brock et al., 2008; Liau et al., 2014). Additionally, pathways regulated by microRNAs

have been associated with DNA hypermethylation-dependent silencing (Saito et al., 2006).

Besides methylating cytosines, DNMTs may coordinate other chromatin-mediated aspects of gene expression at sites of gene promoters (Herman and Baylin, 2003). For example, hypermethylation of promoters of tumor suppressor genes is associated with recruitment of proteins belonging to the methyl CpG-binding domain (MBD) family, MeCP2, MBD1, MBD2, MBD3, and MBD4 (Ballestar and Esteller, 2005). It has been shown that MeCP2 represses transcription of methylated DNA by recruiting histone deacetylases (HDACs), providing the first evidence for interactions between DNA methylation and histone modifications (Jones et al., 1998; Nan et al., 1998).

There is evidence that 1,25-D₃ is able to induce DNA demethylation, however, the mechanisms behind the effect of 1,25-D₃ on DNA methylation are not clear. In most cases it is probably passive demethylation that happens over several cycles of DNA replication. However, in some cases demethylation occurs within 1–4 h, indicative of an active process (Doig et al., 2013). The fact that vitamin D can alter methylation of DNA in the promoter of certain genes is novel. Tapp and colleagues suggested that in healthy subjects global, age-related CGI methylation of human rectal mucosa was influenced not only by gender, folate availability, and selenium, but also by vitamin D status (Tapp et al., 2013). The authors show negative association between serum 25-D₃ level and CGI methylation of the adenomatous polyposis coli (*APC*) promoter region, a tumor suppressor often inactive in colorectal cancer. Interestingly, they observed a weak positive correlation of vitamin D level with methylation of *LINE-1* (genomic long interspersed nuclear element-1), a mammalian autonomous retrotransposon, increasing stability of this region (Tapp et al., 2013). A recent study in colorectal cancer patients investigating two Canadian populations (from Newfoundland and Ontario) found that high dietary vitamin D intake was associated with lower methylation of the two WNT antagonists *dickkopf 1* (*DKK1*) and *WNT5A* (Rawson et al., 2012). This relationship became even more significant in females in the Newfoundland population, while in the Ontario population the association between vitamin D intake and lower methylation was observed only in early stage tumors, but not in late stage tumors (Rawson et al., 2012). These data confer further insights in the mechanisms regulating the transcriptional activating effect of vitamin D on *DKK1* expression described *in vitro* (Aguilera et al., 2007; Pendas-Franco et al., 2008).

Moreover, treatment of the triple negative breast cancer cell line MDA-MB-231 with 1,25-D₃ reduced DNA methylation of the *e-cadherin* promoter (Lopes et al., 2012), while another study showed that 1,25-D₃ induced demethylation of the *PDZ-LIM* domain-containing protein 2 promoter, leading to increased expression (Vanoirbeek et al., 2014). In non-malignant and malignant prostate epithelial cells, treatment with 1,25-D₃ caused clear changes in site-specific methylation of the *p21* promoter, in a cell line-specific manner (Doig et al., 2013).

An interesting interaction between vitamin D and DNA methylation is induction of the expression of GADD45 (growth arrest and DNA damage) protein by 1,25-D₃ in several tumor

cells (Jiang et al., 2003; Zhang et al., 2006; Bremner et al., 2012). GADD45A is one of the enzymes that promote epigenetic gene activation by repair mediated DNA demethylation in *Xenopus laevis* (Barreto et al., 2007).

In summary, alterations in DNA methylation lead to aberrant gene expression and disruptions of genomic integrity, which contribute to development and progression of diseases. Vitamin D can regulate these processes; the mechanisms behind need further investigations.

INTERACTIONS OF VITAMIN D WITH CHROMATIN MODULATORS AND REMODELERS


Nuclear receptors, such as the VDR contain DNA-binding domains that mediate binding to the DNA, presuming the DNA is available and is not wound tightly around nucleosomes. The chromatin context determines nuclear receptor binding and determines which epigenetic modifications will occur thereafter. Upon binding to their genomic response elements, nuclear hormone receptors will then recruit different regulatory cofactor complexes (Lee et al., 2001). The unliganded VDR is able to bind also genomic DNA, where it usually forms complexes with corepressor proteins that either exert HDAC activity, e.g., ALIEN (Polly et al., 2000), or are associated with HDACs, such as NCOR1 and SMRT. The corepressors dissociate upon binding of 1,25-D₃, and are replaced by coactivator complexes.

The chromatin environment dictates gene activity throughout the genome. Post-translational modifications of the N-terminal tails of histone proteins allow nucleosomes to shift, the chromatin to relax, and genes to become activated. Histone modifications change in response to environmental stimuli (Meyer et al., 2013). Histones are major protein components of chromatin that undergo post-translational modifications, including acetylation of lysines, methylation of lysines and arginines, and phosphorylation of serine and threonine residues (Esteller, 2008). In epigenetically silenced genes, hypermethylation of CGIs is often associated with loss of acetylation on histone 3 and 4 (H3 and H4), loss of methylation of lysine (K) 4 on H3 (H3K4), and gain of methylation of K9 and K27 on H3 (H3K9 and H3K27) (Esteller, 2008).

Histone acetylation generally correlates with transcriptional activation (Hebbes et al., 1988; Kouzarides, 2007) and is dependent on a dynamic interaction between histone acetyltransferases (HATs) and histone deacetylases (HDACs) (Marks et al., 2001). The balance between the actions of these enzymes is crucial in controlling gene expression, and governs several developmental processes and disease states (Haberland et al., 2009). Generally, HATs are defined as activators of transcription, whereas HDACs as transcription repressors (Parbin et al., 2014). In various cancer types, including prostate, gastric, and breast cancers, overexpression of HDAC1 is often associated with poor clinical outcome (Choi et al., 2001; Halkidou et al., 2004; Zhang et al., 2005). In colorectal cancer patients HDAC1, 2, and 3 are overexpressed, and high HDAC1 and 2 expression is linked with reduced patient survival (Zhu et al., 2004; Wilson et al., 2006). Overexpression of HDAC1 plays a crucial role in regulating proliferation by repressing the expression of the cyclin-dependent kinase inhibitor p21 (Lagger et al., 2003). Additionally, silencing of HDAC4 leads to re-expression of p21, which in turn induces cell growth arrest and tumor growth inhibition, both *in vitro* and *in vivo* in a human glioblastoma model (Mottet et al., 2009). In addition to classical HDACs, another group of enzymes, the sirtuins (silent information regulator 2 proteins) are involved in histone deacetylation (Schwer and Verdin, 2008). Sirtuins have been linked to metabolic disorders, cancer, aging, and also regulation of the circadian rhythm (Guarente, 2006; Longo and Kennedy, 2006; Jung-Hynes et al., 2010).

Many of the coactivators recruited by the VDR, including p160 steroid receptor coactivator proteins (SRC1, 2, and 3), p300, or CBP have lysine acetyltransferase activity (Figure 1). Indeed, treatment of THP-1 cells with 1,25-D₃ increased H3K27ac at the promoter of several early VDR target genes (Seuter et al., 2013). In genetic hypercalciuric stone forming rats inhibition of bone morphogenetic protein 2 (BMP2) by 1,25-D₃, seems to involve H3 deacetylation and H3K9 di-methylation (Fu et al., 2013).

In MDA-MB453 breast cancer cells 1,25-D₃ treatment regulates expression of p21 through a mechanism involving both histone acetylation and methylation, probably by dynamic



chromatin looping from distal 1,25-D₃ responsive elements to the TSS of *p21* (Saramäki et al., 2009).

Histone methylation can lead either to gene activation or repression, depending on the histone site that is methylated, the degree of methylation (e.g., mono-methylation, di-methylation, or tri-methylation), amino acid residues affected, and their position in the histone tail (Esteller, 2008). Methylation of histones depends on a dynamic process arising from the actions of methyltransferases (HMTs) and demethylases (HDMs) (Shi and Whetstine, 2007; Mosammamaparast and Shi, 2010; Greer and Shi, 2012). So far, two protein families capable of demethylating lysines are known, the amine oxidases (Shi et al., 2004) and jumonji C (JmjC)-domain-containing proteins (Cloos et al., 2006; Tsukada et al., 2006). The first histone demethylase discovered was the lysine-specific demethylase 1 (LSD1/KDM1A), an amine oxidase, which demethylates H3K4me2/me1 (Table 1) (Shi et al., 2004). High expression of KDM1A in various cancers, including colorectal cancer, prostate cancer, and neuroblastomas is associated with increased cancer recurrence and poor clinical outcome (Kahl et al., 2006; Schulte et al., 2009; Ding et al., 2013). Additionally, it has been shown that LSD1 is essential for androgen and estrogen receptor-dependent gene activation via H3K9me2/me1 demethylation (Metzger et al., 2005; Garcia-Bassets et al., 2007; Perillo et al., 2008). There is a reciprocal regulatory effect between the activity of VDR and histone demethylases. In the colon cancer cell line SW480-ADH 1,25-D₃

increased the expression of the lysine-specific demethylase 1 and 2 (Pereira et al., 2012).

1,25-D₃ treatment affected also the expression of a series of different JmjC histone demethylases. The first identified member of the JmjC family was KDM2A/JHDM1A (Tsukada et al., 2006). Expression profiling data showed altered expression of KDM2A and KDM2B in several tumors, however, it seems that their pro- or antioncogenic functions are tissue-dependent (Frescas et al., 2007, 2008; Pfau et al., 2008). 1,25-D₃ inhibited the expression of several histone demethylases (e.g., KDM4A/4C/4D/5A/2B, JMJD5/6, PLA2G4B), and induced the expression of others, JARID2 and KDM5B (Pereira et al., 2012). Members of the KDM4 family catalyze tri-demethylation of H3K9 and/or H3K36 (Cloos et al., 2006; Fodor et al., 2006; Klose et al., 2006; Whetstine et al., 2006; Lin et al., 2008). H3K9me3 is a mark for heterochromatin and demethylation of H3K9 is suggested to be linked with chromosomal instability (Cloos et al., 2006). Inhibition of expression of KDM4 family members by 1,25-D₃ could thus contribute to genome stability. Members of KDM5 cluster catalyze demethylation of H3K4me3/me2, which is a mark for open chromatin (Christensen et al., 2007; Iwase et al., 2007; Klose et al., 2007; Tahiliani et al., 2007) and their upregulation upon 1,25-D₃ treatment might lead to gene repression (Pereira et al., 2011). Overexpression of KDM5B has been reported in breast and prostate cancers (Lu et al., 1999; Xiang et al., 2007). Deletion of *kdm5b* inhibits tumor growth in a syngeneic mouse mammary

Table 1 | A simplified list of the members of the two classes of histone demethylases (mentioned in the manuscript).

Class of histone demethylases	Histone demethylase family	Histone demethylase	Histone substrate	Gene expression
Amine oxidases	KDM1	KDM1A	H3K4me2/me1 H3K9me2/me1	Repression Activation
		KDM1B	H3K4me2/me1	Repression
Jumonji C-domain-containing proteins	KDM2	KDM2A	H3K36me2/me1	Repression
		KDM2B	H3K4me3 H3K36me2/me1	
	KDM3	KDM3A	H3K9me2/me1	Activation
		KDM3B	H3K9me3/me2/me1	
	KDM4	KDM4A	H3K9me3/me2	Activation
		KDM4B	H3K36me3/me2	Repression
		KDM4C		
	KDM5	KDM4D	H3K9me3/me2	Activation
		KDM5A	H3K4me3/me2	Repression
		KDM5B KDM5C KDM5D		
KDM6	KDM6A	H3K27me3/me2	Activation	
	KDM6B			
PHF	JHDM1D	H3K9me2/me1 H3K27me2/me1	Activation	
	PHF8	H3K9me2/me1		

Reviewed in Pedersen and Helin (2010), Greer and Shi (2012). The main histone demethylase families and submembers are indicated. Degree of methylation and site of lysine residue are given. References for individual enzymes can be found throughout the text. H, Histone; K, lysine; me1, mono-methylation; me2, di-methylation; me3, tri-methylation; KDM1A, lysine-specific demethylase 1A; JHDM1D, JmjC-domain-containing histone demethylation protein 1D; PHF, plant homeodomain finger protein.

tumor (Yamane et al., 2007), suggestive of its potential role in tumor development. 1,25-D₃ induced the expression of the histone demethylase KDM6B as well, which is the only other known enzyme, besides KDM6A that is able to demethylate H3K27me₃, a histone mark that correlates with gene repression. Furthermore, the authors showed positive correlation between KDM6B and VDR in 96 colon tumor patients, and inverse correlation of KDM6B with SNAIL1, which is involved in epithelial to mesenchymal transition, indicating that probably the antiproliferative role of 1,25-D₃ via KDM6B upregulation might take place *in vivo* (Pereira et al., 2011). Interestingly, treatment of SW480-ADH cells with 1,25-D₃ had no effect on global H3K27me₃ levels, in spite of KDM6B upregulation (Pereira et al., 2011, 2012). The effect of 1,25-D₃ on the expression of histone demethylases may well be indirect and could be mediated by microRNAs (Padi et al., 2013). KDM2A is one of the direct targets of microRNA-627. 1,25-D₃-dependent upregulation of the microRNA-627 expression both *in vitro*, in the HT-29 colorectal cancer cells and *in vivo*, in tumor xenografts, led to lower KDM2A levels (Padi et al., 2013).

In different pathologies, the expression pattern of the nuclear receptor cofactors is altered, compromising the effect of 1,25-D₃ (Doig et al., 2013; Singh et al., 2013). The initial interactions between VDR and coactivators are the seed for the assembly of intricate multiprotein complexes that remodel the chromatin structure, recruit the core transcriptional machinery, and induce expression of 1,25-D₃ target genes (Figure 1). Often, differences in responsiveness to 1,25-D₃ depend on the expression pattern of the coregulators of VDR. In prostate cancer cells, the temporal distribution of the nuclear corepressor NCOR1 at VDR target genes is different in 1,25-D₃ responsive cells compared with unresponsive cells (Doig et al., 2013; Singh et al., 2013).

The liganded VDR is able both to transactivate and transrepress target genes. The mechanisms of action are probably different between transactivation and transrepression, and also highly dependent on the motifs of the vitamin D response elements. A highly complex mechanism regulates the ligand-dependent repression of CYP27B1 (Kim et al., 2007a). CYP27B1 repression requires two epigenetic modifications: deacetylation of histones and methylation of the CYP27B1 gene promoter and exon regions. This is dependent on the presence of the VDR interacting repressor (VDIR) and the chromatin remodeler Williams Syndrome transcription factor. In the absence of 1,25-D₃, VDIR is bound directly to the E-box motifs in the negative VDRE and recruits histone acetyltransferases to induce CYP27B1 gene transcription. In the presence of 1,25-D₃, VDIR acts as a scaffold for the 1,25-D₃-VDR complex to repress transcription of CYP27B1 through recruitment of HDAC2, DNMT1, and DNMT3B (Kim et al., 2007a). It seems that VDIR and HDAC2 are involved also in the 1,25-D₃-dependent transrepression of the human parathyroid hormone gene (Kim et al., 2007b). It is not clear, whether this mechanism of transrepression by liganded VDR also applies to other genes. In mesenchymal stem cells 1,25-D₃ represses gene expression by binding to promoters with enhanced H3K9Ac and H3K9me₂ (Tan et al., 2009). Whether H3K9 acetylation/methylation enabled VDR binding or VDR binding caused H3K9 acetylation, is not clear.

Proper orchestration of histone modifications in crosstalk with other chromatin regulators is crucial in maintaining the epigenetic landscape and governing gene expression. Any disturbances in these constellations may lead to aberrant gene expression. Whether 1,25-D₃ affects regulation of other chromatin modulators as well, is not yet known.

REGULATION OF THE VITAMIN D SYSTEM


The vitamin D system has pleiotropic functions and regulates approximately 3% of the human genome (Bouillon et al., 2008). To maintain balance, a strict regulation of the vitamin D system genes is of utmost importance. The main role of liganded VDR in tissues not involved in calcium homeostasis is to control expression of genes that regulate cell proliferation, differentiation, and apoptosis. One major limitation in the therapeutic exploitation of these effects is the resistance of cancer cells to 1,25-D₃. Epigenetic corruption of VDR signaling is suggested to be one of the mechanisms that leads to reduced responsiveness to 1,25-D₃ actions. This can be caused by promoter methylation of key vitamin D system genes or by skewed accumulation of VDR-associated corepressors, preferentially at promoters of anti-proliferative target genes (Abedin et al., 2006).

Expression of the vitamin D degrading and metabolizing enzymes is regulated through binding of 1,25-D₃-liganded VDR to vitamin D responsive elements (VDREs). However, the major regulators of 1,25-D₃ levels and signaling CYP2R1, CYP24A1, CYP27B1, and VDR, “the vitamin D tool” genes, are prone to epigenetic regulation. CpG islands span the promoters of CYP2R1, CYP24A1, and VDR, while a CpG island is located within the CYP27B1 gene locus (Figure 2). Therefore, DNA methylation and histone modifications in these regions can change the chromatin state from an open to closed conformation and lead to transcriptional repression of these genes. Expression of vitamin D tool genes becomes deregulated in various types of cancer, and these changes may be partially attributed to epigenetic alterations (reviewed in Hobaus et al., 2013). As early as 1984, Yoneda et al. (1984) have shown that the histone acetyltransferase inhibitor butyrate augments 1,25-D₃ actions. Several studies confirmed these findings (e.g., Rashid et al., 2001) and have suggested that the action of butyrate could be through upregulation of VDR or CYP27B1 expression (Gaschott and Stein, 2003). Whether this effect is mediated by direct acetylation of the VDR or CYP27B1 promoters, has not been determined.

In this chapter we discuss evidence for epigenetic regulation through DNA methylation of these genes in health and disease.

EPIGENETIC REGULATION OF THE VITAMIN D RECEPTOR

The VDR is a nuclear receptor mediating 1,25-D₃ signaling. It is expressed by at least 38 cell types in the human body (Norman and Bouillon, 2010). In the absence of its ligand 1,25-D₃, VDR is mainly found in the cytoplasm (Nagpal et al., 2005). Upon ligand binding, VDR heterodimerizes with the retinoid X receptor (RXR) and translocates to the nucleus, where it binds to vitamin D responsive elements (VDREs) to regulate transcription of 1,25-D₃ target genes. This is achieved through recruitment of coactivators or corepressors to the VDR-RXR complex bound to DNA (Nagpal et al., 2005; Pike et al., 2012; Haussler et al., 2013;



Pike and Meyer, 2013). As VDR is rarely mutated during carcinogenesis (Miller et al., 1997), the disturbance of the vitamin D signaling and apparent 1,25-D₃ insensitivity in cancer (Marik et al., 2010) must be attributed to other alterations, which may include epigenetic changes, such as promoter methylation.

The VDR gene is located on the long arm of chromosome 12 (12q13.11) and contains 4 potential promoter regions. Exons 1a, 1c, and 1d of the VDR are well conserved, while 1b, 1e, and 1f show low homology (Halsall et al., 2007). Exon 1a appears to contain a strong promoter, including several transcription factor binding sites (AP-2 and SP1). Transcription was reported to originate also in exons 1d, 1e, and 1f, while translation starts in exon 2 (Halsall et al., 2007). Marik and colleagues performed an *in silico* analysis of the VDR gene sequence and reported three CpG islands located in exon 1a spanning from -790 bp to 380 bp relative to the TSS in exon 1a (Marik et al., 2010). According to the UCSC Genome browser, however, only one large CpG island spanning 892 bp in length is found in the VDR promoter region (Gardiner-Garden and Frommer, 1987; Karolchik et al., 2014). This discrepancy is likely due to the different search parameters used for CpG island identification.

Epigenetic silencing of VDR was suggested to cause the slow normalization of VDR levels in the parathyroid glands of uremic rats after kidney transplantation (Lewin et al., 2002; Hofman-Bang et al., 2012). However, sequencing of the VDR promoter [-250 to 300 bp relative to exon 1 (43 CpGs)] in normal and

uremic rats showed no difference between methylation patterns (Hofman-Bang et al., 2012). Further, the authors reported that methylation levels coincided with the negative control, thus showing that promoter methylation does not play a role in regulating VDR expression in the parathyroid glands.

In contrast, promoter methylation was reported to cause repression of VDR gene expression in HIV infected T cells. In normal T cells, activation or priming causes an upregulation of VDR expression (Von Essen et al., 2010). In comparison, infection of previously activated T cells with human immunodeficiency virus (HIV) led to upregulation of DNMT3B, increased promoter methylation of VDR (45–70%), and decreased VDR gene expression (Chandel et al., 2013). This downregulation of VDR could be reversed upon treatment with 5-azacytidine (AZA) suggesting that the decreased expression of VDR by HIV is, at least partially, caused by DNA methylation (Chandel et al., 2013). There is evidence for an inverse correlation between the vitamin D status and infections, however, many trials failed to show a protective effect of vitamin D (reviewed in Yamshchikov et al., 2009). Thus, reduced sensitivity to vitamin D metabolites due to, e.g., downregulation of VDR may account for inconclusive trials. This is supported by a study investigating methylation of the 3' end of VDR in two South African groups revealing differences with respect to ethnicity and tuberculosis status of the patients (Andraos et al., 2011).

In breast tumors, methylation of exon 1a of the VDR gene was significantly higher (65% of CpGs methylated) compared with normal breast tissue (15% of CpGs methylated) (Marik et al., 2010). *In vitro*, in breast cancer cell lines, three hypermethylated regions in exon 1a became demethylated after treatment with the DNMT1 inhibitor 5-aza-2'-deoxycytidine (DAC) and VDR mRNA expression increased. These regions were in proximity to the SP1 binding sites (approximately 790 bp from TSS), NFκB binding sites (approximately -480 from TSS), and the exon 1a TSS. Treatment with 1,25-D₃ had no effect on methylation of these regions (Marik et al., 2010). In other types of cancer, e.g., the choriocarcinoma-derived trophoblast cell lines JEG-3 and JAR, the VDR promoter was densely methylated (Novakovic et al., 2009). In contrast, no methylation of the VDR promoter region was observed in colon cancer cell lines, and treatment with DAC did not increase gene expression (Habano et al., 2011; Höbaus et al., 2013a). In parathyroid tumors the expression of VDR is decreased (Gogusev et al., 1997; Carling et al., 2000), however, no differences in DNA methylation of VDR were observed between parathyroid tumors and healthy controls (Sulaiman et al., 2013). Similar results were seen in parathyroid adenoma samples, which showed decreased expression of VDR, but showed no promoter methylation (Varshney et al., 2013).

Additionally, it has been suggested that expression of 5' truncated variants of VDR is linked to methylation of the VDR promoter. These variants are predominantly found in breast cancer compared with the full length variants expressed in normal breast tissue. Treatment with DAC restored expression of the active transcript variant of VDR in breast cancer cell lines, indicating promoter methylation as cause of truncated protein expression (Marik et al., 2010). The significance of these potentially untranslated truncated variants remains to be investigated, however, as

they are not found in normal breast tissue aberrant expression of truncated isoforms may further disrupt vitamin D signaling in tumor tissue.

A recent study suggested that in colorectal cancer metastases, VDR becomes the target of the polycomb group protein enhancer of zeste homolog 2 (EZH2) that mediates VDR downregulation by H3K27 trimethylation in the VDR promoter (Lin et al., 2013). The histone deacetylase HDAC3, one of the most frequently upregulated genes in cancer, seems to inhibit VDR expression. In two colorectal cancer cell lines, HCT116 and SW480 knock down of HDAC3 increased VDR expression and restored sensitivity of these cells to 1,25-D₃ (Godman et al., 2008).

Taken together, there is evidence that in various diseases the decreased tissue sensitivity to 1,25-D₃ could have been caused by the epigenetic silencing of the VDR.

EPIGENETIC REGULATION OF THE CYP2R1

CYP2R1 is a microsomal P450 enzyme, which hydroxylates both vitamin D₂ and D₃ at position C-25 to form the circulating storage form of vitamin D 25-D₃. The promoter region of *CYP2R1* is located within a CpG island, which can be subjected to epigenetic regulation. So far, only two studies investigated the promoter methylation status of this gene. Genome wide association studies found increased *CYP2R1* promoter methylation in leukocyte DNA from individuals with severe vitamin D deficiency compared with control group (Zhu et al., 2013). Further, methylation levels of *CYP2R1* promoter decreased within 12 months of vitamin D supplementation in DNA extracted from serum of non-Hispanic white American post-menopausal women aged ≥ 55 years (Zhu et al., 2013), indicating an effect of vitamin D supplementation on *CYP2R1* promoter methylation. These data indicate that under low vitamin D serum levels, the promoter of the major 25-hydroxylase *CYP2R1* may become methylated, and that event appears to be reversible upon exposure to increased vitamin D.

EPIGENETIC REGULATION OF THE CYP27B1

CYP27B1 is an inner mitochondrial membrane P450 enzyme that converts 25-D₃ to its active form 1,25-D₃. It is mainly expressed in the proximal tubule of the kidneys, but it is also expressed in many vitamin D target tissues, albeit at lower levels (Hendrix et al., 2004). The *CYP27B1* gene harbors a CpG island. However, recent sequence updates (Ensembl 74, November 2013) shift the CpG island from the *CYP27B1* promoter region into the gene coding sequence (Flicek et al., 2014). This explains the differences between the location of the CpG island depicted in **Figure 2** and the location of the CpG island described in literature. For simplicity, statements on nVDRES and CpG island location below refer to reports in the published articles and not to **Figure 2**.

The promoter region of *CYP27B1* contains a negative VDRE (nVDRE) located at around 500 bp, consisting of two E-box like motifs (Murayama et al., 2004). This region is responsible for 1,25-D₃-dependent transrepression, which seems to be achieved through recruitment of both HDACs and DNMTs by VDR/RXR to the promoter region of *CYP27B1* (Takeyama and Kato, 2011). For further details see subsection Interactions of Vitamin D with Chromatin Modulators and Remodelers.

In cancer, expression of CYP27B1 is often downregulated. This may be explained by increased methylation of the CpG island located within *CYP27B1*. In the breast cancer cells MDA-MB-231, *CYP27B1* hypermethylation led to gene silencing, which could be reversed by treatment with deoxyC (Shi et al., 2002). In prostate cancer cell lines, combination of the DNMT1 inhibitor DAC and the HDAC inhibitor TSA resulted in increased activity of CYP27B1 (Khorchide et al., 2005). In the choriocarcinoma cell lines BeWo and JAR the promoter of *CYP27B1* was densely methylated (Novakovic et al., 2009). The *CYP27B1* promoter was hypermethylated (61%) in Non-Hodgkin's lymphoma, but not in benign follicular hyperplasia. Two out of four non-Hodgkin's lymphoma cell lines showed strong methylation of the *CYP27B1* promoter. Interestingly, all four responded to DAC-TSA treatments with upregulation of gene expression independent of the methylation status of their *CYP27B1* promoter, which may be explained by other regions prone to methylation not investigated in this study or by differences in silencing mechanisms (Shi et al., 2007). Further, the methylation level of *CYP27B1* was increased in primary lymphoma and leukemia cells also compared with normal peripheral blood lymphocytes (Lagger et al., 2003; Wjst et al., 2010).

Methylation of *CYP27B1* in diseases might cause reduced local activation of 25-D₃ to 1,25-D₃, thus reducing local 1,25-D₃ levels and restricting its functions.

EPIGENETIC REGULATION OF CYP24A1

The 1,25-dihydroxyvitamin D₃ 24-hydroxylase is an inner mitochondrial membrane P450 enzyme, which catalyzes both 25-D₃ and 1,25-D₃ (Kawashima et al., 1981; Pedersen et al., 1983; Sakaki et al., 2000). Its primary site of expression are the kidneys, where it plays a crucial role in regulating systemic vitamin D metabolite levels, however, expression is found in many other vitamin D target tissues.

The promoter of *CYP24A1* is spanned by a CpG island making it prone to regulation by DNA methylation. Several responsive elements are located within this area, including two VDREs, a vitamin stimulating element (VSE), and SP1 binding sites (see **Figure 2**).

In healthy kidney, skeletal muscle, whole blood, brain, skin fibroblasts, and sperm the *CYP24A1* promoter is not methylated (Novakovic et al., 2009), although the expression levels are highly variable. In peripheral blood lymphocytes methylation of *CYP24A1* was low (5%) (Wjst et al., 2010). Interestingly, in full term human placenta 56.5% of the *CYP24A1* promoter is methylated. *CYP24A1* methylation was also observed in the placenta of the marmoset and mouse, however, at a lower level.

In the choriocarcinoma cell lines JEG-3, BeWo, and JAR the promoter of *CYP24A1* was densely methylated and the methylation level correlated inversely with the low gene expression (Novakovic et al., 2009). Treatment of osteoblastic ROS cells with 1,25-D₃ did not induce CYP24A1 expression. Considering the strong methylation of the *CYP24A1* promoter region, epigenetic silencing of CYP24A1 may account for the unresponsiveness of this gene to 1,25-D₃ (Ohyama et al., 2002). In the human prostate cancer cell line PC3, methylation of the *CYP24A1*

promoter reduced reporter gene expression in a methylation-dependent manner (Luo et al., 2010). In prostate cancer cells, the methylation status of *CYP24A1* promoter inversely correlated with gene expression. Demethylating agents restored *CYP24A1* expression only in cell lines where the *CYP24A1* promoter was methylated prior to treatment (Luo et al., 2010). Only DNA demethylation by DAC treatment permitted recruitment of VDR to the *CYP24A1* promoter (Luo et al., 2010). In patients, development from benign toward malignant prostate lesions was paralleled by increasing methylation levels of the *CYP24A1* promoter (Luo et al., 2010). Prostate tumor derived endothelial cells (TDEC) expressed less *CYP24A1* compared with endothelial cells derived from normal cells or matrigel plugs, which may be attributed to increased *CYP24A1* promoter methylation in TDECs (Johnson et al., 2010). We have shown recently that in colon cancer cell lines DAC induced *CYP24A1* expression in a cell line-specific manner, independent of the methylation level of the promoter. In these cells induction of *CYP24A1* expression by DAC seems to be independent of *CYP24A1* promoter methylation (Höbaus et al., 2013a). Moreover, the methylation level of the *CYP24A1* promoter was comparably low both in colon adenocarcinomas and the adjacent mucosa, although the expression of *CYP24A1* was significantly higher in the tumors (Höbaus et al., 2013b).

Taken together, the regulation of *CYP24A1* by DNA methylation appears to be tissue-dependent, both in health and disease.

CONCLUSIONS

There is a strong reciprocity between the vitamin D system and epigenetic mechanisms. The vitamin D system is, on the one hand regulated by epigenetic mechanisms and, on the other hand, is involved in regulating epigenetic events. Critical vitamin D tool genes can be silenced by DNA methylation. The VDR protein interacts, directly or indirectly, with chromatin modifiers and remodelers. Liganded VDR regulates expression of several of these chromatin modifiers and remodelers, and it might even regulate DNA methylation.

Epigenetic regulation of gene expression is a fine-tuned mechanism and its deregulation can lead to pathological conditions. The impact of vitamin D in the maintenance of the normal epigenetic landscape underlines the central role of this hormone in physiology.

PERSPECTIVES

One of the most fundamental questions in the control of gene expression is the way how epigenetic marks are established, erased, and recognized. Regulating epigenetic events could be a further mechanism by which 1,25-D₃ may prevent or delay tumorigenesis and onset of chronic diseases. Therefore, we need to understand better the impact of vitamin D on the epigenome, and plan thorough and comprehensive studies to examine this interplay.

ACKNOWLEDGMENTS

We thank the Austrian Research Fund (FWF) Grant P22200-B11 and the European Union, FP7-PEOPLE-2010 Grant 264663 for support.

REFERENCES

- Abedin, S. A., Banwell, C. M., Colston, K. W., Carlberg, C., and Campbell, M. J. (2006). Epigenetic corruption of VDR signalling in malignancy. *Anticancer Res.* 26, 2557–2566.
- Aguilera, O., Pena, C., Garcia, J. M., Larrriba, M. J., Ordóñez-Moran, P., Navarro, D., et al. (2007). The Wnt antagonist DICKKOPF-1 gene is induced by 1 α ,25-dihydroxyvitamin D₃ associated to the differentiation of human colon cancer cells. *Carcinogenesis* 28, 1877–1884. doi: 10.1093/carcin/bgm094
- Andraos, C., Koorsen, G., Knight, J. C., and Bornman, L. (2011). Vitamin D receptor gene methylation is associated with ethnicity, tuberculosis, and TaqI polymorphism. *Hum. Immunol.* 72, 262–268. doi: 10.1016/j.humimm.2010.12.010
- Arasaradnam, R. P., Commane, D. M., Bradburn, D., and Mathers, J. C. (2008). A review of dietary factors and its influence on DNA methylation in colorectal carcinogenesis. *Epigenetics* 3, 193–198. doi: 10.4161/epi.3.4.6508
- Ballestar, E., and Esteller, M. (2005). Methyl-CpG-binding proteins in cancer: blaming the DNA methylation messenger. *Biochem. Cell Biol.* 83, 374–384. doi: 10.1139/o05-035
- Barreto, G., Schafer, A., Marhold, J., Stach, D., Swaminathan, S. K., Handa, V., et al. (2007). Gadd45a promotes epigenetic gene activation by repair-mediated DNA demethylation. *Nature* 445, 671–675. doi: 10.1038/nature05515
- Baylin, S. B., and Jones, P. A. (2011). A decade of exploring the cancer epigenome - biological and translational implications. *Nat. Rev. Cancer* 11, 726–734. doi: 10.1038/nrc3130
- Bestor, T. H. (2005). Transposons reanimated in mice. *Cell* 122, 322–325. doi: 10.1016/j.cell.2005.07.024
- Bird, A. (2002). DNA methylation patterns and epigenetic memory. *Genes Dev.* 16, 6–21. doi: 10.1101/gad.947102
- Bird, A. P. (1980). DNA methylation and the frequency of CpG in animal DNA. *Nucleic Acids Res.* 8, 1499–1504. doi: 10.1093/nar/8.7.1499
- Bjornsson, H. T., Brown, L. J., Fallin, M. D., Rongione, M. A., Bibikova, M., Wickham, E., et al. (2007). Epigenetic specificity of loss of imprinting of the IGF2 gene in Wilms tumors. *J. Natl. Cancer Inst.* 99, 1270–1273. doi: 10.1093/jnci/djm069
- Bouillon, R., Carmeliet, G., Verlinden, L., Van Etten, E., Verstuyf, A., Luderer, H. E., et al. (2008). Vitamin D and human health: lessons from vitamin D receptor null mice. *Endocr. Rev.* 29, 726–776. doi: 10.1210/er.2008-0004
- Bremmer, F., Thelen, P., Pottek, T., Behnes, C. L., Radzun, H. J., and Schwyer, S. (2012). Expression and function of the vitamin D receptor in malignant germ cell tumour of the testis. *Anticancer Res.* 32, 341–349.
- Brock, M. V., Hooker, C. M., Ota-Machida, E., Han, Y., Guo, M., Ames, S., et al. (2008). DNA methylation markers and early recurrence in stage I lung cancer. *N. Engl. J. Med.* 358, 1118–1128. doi: 10.1056/NEJMoa0706550
- Carling, T., Rastad, J., Szabó, E., Westin, G., and Akerström, G. (2000). Reduced parathyroid vitamin D receptor messenger ribonucleic acid levels in primary and secondary hyperparathyroidism. *J. Clin. Endocrinol. Metab.* 85, 2000–2003. doi: 10.1210/jc.85.5.2000
- Chandel, N., Husain, M., Goel, H., Salhan, D., Lan, X., Malhotra, A., et al. (2013). VDR hypermethylation and HIV-induced T cell loss. *J. Leukoc. Biol.* 93, 623–631. doi: 10.1189/jlb.0812383
- Choi, J. H., Kwon, H. J., Yoon, B. I., Kim, J. H., Han, S. U., Joo, H. J., et al. (2001). Expression profile of histone deacetylase 1 in gastric cancer tissues. *Jpn. J. Cancer Res.* 92, 1300–1304. doi: 10.1111/j.1349-7006.2001.tb02153.x
- Christensen, J., Agger, K., Cloos, P. A., Pasini, D., Rose, S., Sennels, L., et al. (2007). RBP2 belongs to a family of demethylases, specific for tri- and dimethylated lysine 4 on histone 3. *Cell* 128, 1063–1076. doi: 10.1016/j.cell.2007.02.003
- Cloos, P. A., Christensen, J., Agger, K., Maiolica, A., Rappsilber, J., Antal, T., et al. (2006). The putative oncogene GASC1 demethylates tri- and dimethylated lysine 9 on histone H3. *Nature* 442, 307–311. doi: 10.1038/nature04837
- Cui, H., Cruz-Correa, M., Giardiello, F. M., Hutcheon, D. F., Kafonek, D. R., Brandenburg, S., et al. (2003). Loss of IGF2 imprinting: a potential marker of colorectal cancer risk. *Science* 299, 1753–1755. doi: 10.1126/science.1080902
- Ding, J., Zhang, Z. M., Xia, Y., Liao, G. Q., Pan, Y., Liu, S., et al. (2013). LSD1-mediated epigenetic modification contributes to proliferation and metastasis of colon cancer. *Br. J. Cancer* 109, 994–1003. doi: 10.1038/bjc.2013.364
- Doig, C. L., Singh, P. K., Dhiman, V. K., Thorne, J. L., Battaglia, S., Sobolewski, M., et al. (2013). Recruitment of NCOR1 to VDR target genes is enhanced in prostate cancer cells and associates with altered DNA methylation patterns. *Carcinogenesis* 34, 248–256. doi: 10.1093/carcin/bgs331

- Eden, A., Gaudet, F., Waghmare, A., and Jaenisch, R. (2003). Chromosomal instability and tumors promoted by DNA hypomethylation. *Science* 300, 455. doi: 10.1126/science.1083557
- Esteller, M. (2007a). Epigenetic gene silencing in cancer: the DNA hypermethylation. *Hum. Mol. Genet.* 16, Spec No 1, R50–R59. doi: 10.1093/hmg/ddm018
- Esteller, M. (2007b). Epigenetics provides a new generation of oncogenes and tumour-suppressor genes. *Br. J. Cancer* 96(Suppl.), R26–R30. doi: 10.1038/sj.bjc.6602918
- Esteller, M. (2008). Epigenetics in cancer. *N. Engl. J. Med.* 358, 1148–1159. doi: 10.1056/NEJMra072067
- Esteller, M., Catusas, L., Matias-Guiu, X., Mutter, G. L., Prat, J., Baylin, S. B., et al. (1999). hMLH1 promoter hypermethylation is an early event in human endometrial tumorigenesis. *Am. J. Pathol.* 155, 1767–1772. doi: 10.1016/S0002-9440(10)65492-2
- Feinberg, A. P., Cui, H., and Ohlsson, R. (2002). DNA methylation and genomic imprinting: insights from cancer into epigenetic mechanisms. *Semin. Cancer Biol.* 12, 389–398. doi: 10.1016/S1044-579X(02)00059-7
- Ficz, G., Branco, M. R., Seisenberger, S., Santos, F., Krueger, F., Hore, T. A., et al. (2011). Dynamic regulation of 5-hydroxymethylcytosine in mouse ES cells and during differentiation. *Nature* 473, 398–402. doi: 10.1038/nature10008
- Flicek, P., Amode, M. R., Barrell, D., Beal, K., Billis, K., Brent, S., et al. (2014). Ensembl 2014. *Nucleic Acids Res.* 42, D749–D755. doi: 10.1093/nar/gkt1196
- Fodor, B. D., Kubicek, S., Yonezawa, M., O'Sullivan, R. J., Sengupta, R., Perez-Burgos, L., et al. (2006). Jmjd2b antagonizes H3K9 trimethylation at pericentric heterochromatin in mammalian cells. *Genes Dev.* 20, 1557–1562. doi: 10.1101/gad.388206
- Frescas, D., Guardavaccaro, D., Bassermann, F., Koyama-Nasu, R., and Pagano, M. (2007). JHDM1B/FBXL10 is a nucleolar protein that represses transcription of ribosomal RNA genes. *Nature* 450, 309–313. doi: 10.1038/nature06255
- Frescas, D., Guardavaccaro, D., Kuchay, S. M., Kato, H., Poleshko, A., Basrur, V., et al. (2008). KDM2A represses transcription of centromeric satellite repeats and maintains the heterochromatic state. *Cell Cycle* 7, 3539–3547. doi: 10.4161/cc.7.22.7062
- Fu, B., Wang, H., Wang, J., Barouhas, I., Liu, W., Shuboy, A., et al. (2013). Epigenetic regulation of BMP2 by 1,25-dihydroxyvitamin D3 through DNA methylation and histone modification. *PLoS ONE* 8:e61423. doi: 10.1371/journal.pone.0061423
- Garcia-Bassets, I., Kwon, Y. S., Telese, F., Prefontaine, G. G., Hutt, K. R., Cheng, C. S., et al. (2007). Histone methylation-dependent mechanisms impose ligand dependency for gene activation by nuclear receptors. *Cell* 128, 505–518. doi: 10.1016/j.cell.2006.12.038
- Gardiner-Garden, M., and Frommer, M. (1987). CpG islands in vertebrate genomes. *J. Mol. Biol.* 196, 261–282. doi: 10.1016/0022-2836(87)90689-9
- Gaschott, T., and Stein, J. (2003). Short-chain fatty acids and colon cancer cells: the vitamin D receptor–butyrate connection. *Recent Results Cancer Res.* 164, 247–257. doi: 10.1007/978-3-642-55580-0_18
- Godman, C. A., Joshi, R., Tierney, B. R., Greenspan, E., Rasmussen, T. P., Wang, H. W., et al. (2008). HDAC3 impacts multiple oncogenic pathways in colon cancer cells with effects on Wnt and vitamin D signaling. *Cancer Biol. Ther.* 7, 1570–1580. doi: 10.4161/cbt.7.10.6561
- Gogusev, J., Duchambon, P., Hory, B., Giovannini, M., Goureau, Y., Sarfati, E., et al. (1997). Depressed expression of calcium receptor in parathyroid gland tissue of patients with hyperparathyroidism. *Kidney Int.* 51, 328–336. doi: 10.1038/ki.1997.41
- Greer, E. L., and Shi, Y. (2012). Histone methylation: a dynamic mark in health, disease and inheritance. *Nat. Rev. Genet.* 13, 343–357. doi: 10.1038/nrg3173
- Greger, V., Passarge, E., Höpping, W., Messmer, E., and Horsthemke, B. (1989). Epigenetic changes may contribute to the formation and spontaneous regression of retinoblastoma. *Hum. Genet.* 83, 155–158. doi: 10.1007/BF00286709
- Gruenbaum, Y., Stein, R., Cedar, H., and Razin, A. (1981). Methylation of CpG sequences in eukaryotic DNA. *FEBS Lett.* 124, 67–71. doi: 10.1016/0014-5793(81)80055-5
- Guarente, L. (2006). Sirtuins as potential targets for metabolic syndrome. *Nature* 444, 868–874. doi: 10.1038/nature05486
- Habano, W., Gamo, T., Terashima, J., Sugai, T., Otsuka, K., Wakabayashi, G., et al. (2011). Involvement of promoter methylation in the regulation of Pregnane X receptor in colon cancer cells. *BMC Cancer* 11:81. doi: 10.1186/1471-2407-11-81
- Haberland, M., Montgomery, R. L., and Olson, E. N. (2009). The many roles of histone deacetylases in development and physiology: implications for disease and therapy. *Nat. Rev. Genet.* 10, 32–42. doi: 10.1038/nrg2485
- Hackett, J. A., Sengupta, R., Zyllicz, J. J., Murakami, K., Lee, C., Down, T. A., et al. (2013). Germline DNA demethylation dynamics and imprint erasure through 5-hydroxymethylcytosine. *Science* 339, 448–452. doi: 10.1126/science.1229277
- Halkidou, K., Gaughan, L., Cook, S., Leung, H. Y., Neal, D. E., and Robson, C. N. (2004). Upregulation and nuclear recruitment of HDAC1 in hormone refractory prostate cancer. *Prostate* 59, 177–189. doi: 10.1002/pros.20022
- Hallsall, J. A., Osborne, J. E., Hutchinson, P. E., and Pringle, J. H. (2007). *In silico* analysis of the 5' region of the Vitamin D receptor gene: functional implications of evolutionary conservation. *J. Steroid. Biochem. Mol. Biol.* 103, 352–356. doi: 10.1016/j.jsbmb.2006.12.046
- Haussler, M. R., Whitfield, G. K., Kaneko, I., Haussler, C. A., Hsieh, D., Hsieh, J. C., et al. (2013). Molecular mechanisms of vitamin D action. *Calcif. Tissue Int.* 92, 77–98. doi: 10.1007/s00223-012-9619-0
- Hebbes, T. R., Thorne, A. W., and Crane-Robinson, C. (1988). A direct link between core histone acetylation and transcriptionally active chromatin. *EMBO J.* 7, 1395–1402.
- Helin, K., and Dhanak, D. (2013). Chromatin proteins and modifications as drug targets. *Nature* 502, 480–488. doi: 10.1038/nature12751
- Hendrix, I., Anderson, P., May, B., and Morris, H. (2004). Regulation of gene expression by the CYP27B1 promoter–study of a transgenic mouse model. *J. Steroid Biochem. Mol. Biol.* 89–90, 139–142. doi: 10.1016/j.jsbmb.2004.03.093
- Herman, J. G., and Baylin, S. B. (2003). Gene silencing in cancer in association with promoter hypermethylation. *N. Engl. J. Med.* 349, 2042–2054. doi: 10.1056/NEJMra023075
- Höbaus, J., Fetahu, I. S. H., Khorchide, M., Manhardt, T., and Kallay, E. (2013a). Epigenetic regulation of the 1,25-dihydroxyvitamin D3 24-hydroxylase (CYP24A1) in colon cancer cells. *J. Steroid Biochem. Mol. Biol.* 136, 296–299. doi: 10.1016/j.jsbmb.2012.08.003
- Höbaus, J., Hummel, D. M., Thiem, U., Fetahu, I. S., Aggarwal, A., Müllauer, L., et al. (2013b). Increased copy-number and not DNA hypomethylation causes overexpression of the candidate proto-oncogene CYP24A1 in colorectal cancer. *Int. J. Cancer* 133, 1380–1388. doi: 10.1002/ijc.28143
- Höbaus, J., Thiem, U., Hummel, D. M., and Kallay, E. (2013). Role of calcium, vitamin D, and the extrarenal vitamin D hydroxylases in carcinogenesis. *Anticancer Agents Med. Chem.* 13, 20–35. doi: 10.2174/187152013804487434
- Hofman-Bang, J., Gravesen, E., Olgaard, K., and Lewin, E. (2012). Epigenetic methylation of parathyroid CaR and VDR promoters in experimental secondary hyperparathyroidism. *Int. J. Nephrol.* 2012:123576. doi: 10.1155/2012/123576
- Iwase, S., Lan, F., Bayliss, P., De La Torre-Ubieta, L., Huarte, M., Qi, H. H., et al. (2007). The X-linked mental retardation gene SMCX/JARID1C defines a family of histone H3 lysine 4 demethylases. *Cell* 128, 1077–1088. doi: 10.1016/j.cell.2007.02.017
- Jiang, F., Li, P., Fornace, A. J., Nicosia, S. V., and Bai, W. (2003). G2/M arrest by 1,25-dihydroxyvitamin D3 in ovarian cancer cells mediated through the induction of GADD45 via an exonic enhancer. *J. Biol. Chem.* 278, 48030–48040. doi: 10.1074/jbc.M308430200
- Jin, B., and Robertson, K. D. (2013). DNA methyltransferases, DNA damage repair, and cancer. *Adv. Exp. Med. Biol.* 754, 3–29. doi: 10.1007/978-1-4419-9967-2_1
- Johnson, C. S., Chung, I., and Trump, D. L. (2010). Epigenetic silencing of CYP24 in the tumor microenvironment. *J. Steroid Biochem. Mol. Biol.* 121, 338–342. doi: 10.1016/j.jsbmb.2010.03.046
- Jones, P. A., and Baylin, S. B. (2007). The epigenomics of cancer. *Cell* 128, 683–692. doi: 10.1016/j.cell.2007.01.029
- Jones, P. L., Veenstra, G. J., Wade, P. A., Vermaak, D., Kass, S. U., Landsberger, N., et al. (1998). Methylated DNA and MeCP2 recruit histone deacetylase to repress transcription. *Nat. Genet.* 19, 187–191. doi: 10.1038/561
- Jung-Hynes, B., Reiter, R. J., and Ahmad, N. (2010). Sirtuins, melatonin and circadian rhythms: building a bridge between aging and cancer. *J. Pineal Res.* 48, 9–19. doi: 10.1111/j.1600-079X.2009.00729.x
- Kahl, P., Gullotti, L., Heukamp, L. C., Wolf, S., Friedrichs, N., Vorreuther, R., et al. (2006). Androgen receptor coactivators lysine-specific histone demethylase 1 and four and a half LIM domain protein 2 predict risk of prostate cancer recurrence. *Cancer Res.* 66, 11341–11347. doi: 10.1158/0008-5472.CAN-06-1570
- Karlic, H., and Varga, F. (2011). Impact of vitamin D metabolism on clinical epigenetics. *Clin. Epigenetics* 2, 55–61. doi: 10.1007/s13148-011-0021-y

- Karolchik, D., Barber, G. P., Casper, J., Clawson, H., Cline, M. S., Diekhans, M., et al. (2014). The UCSC Genome Browser database: 2014 update. *Nucleic Acids Res.* 42, D764–770. doi: 10.1093/nar/gkt1168
- Kawashima, H., Torikai, S., and Kurokawa, K. (1981). Localization of 25-hydroxyvitamin D₃ 1 alpha-hydroxylase and 24-hydroxylase along the rat nephron. *Proc. Natl. Acad. Sci. U.S.A.* 78, 1199–1203. doi: 10.1073/pnas.78.2.1199
- Khorchide, M., Lechner, D., and Cross, H. S. (2005). Epigenetic regulation of vitamin D hydroxylase expression and activity in normal and malignant human prostate cells. *J. Steroid Biochem. Mol. Biol.* 93, 167–172. doi: 10.1016/j.jsbmb.2004.12.022
- Kim, G. D., Ni, J., Kelesoglu, N., Roberts, R. J., and Pradhan, S. (2002). Cooperation and communication between the human maintenance and de novo DNA (cytosine-5) methyltransferases. *EMBO J.* 21, 4183–4195. doi: 10.1093/emboj/cdf401
- Kim, M. S., Fujiki, R., Kitagawa, H., and Kato, S. (2007a). 1alpha,25(OH)2D3-induced DNA methylation suppresses the human CYP27B1 gene. *Mol. Cell Endocrinol.* 265–266, 168–173. doi: 10.1016/j.mce.2006.12.014
- Kim, M. S., Fujiki, R., Murayama, A., Kitagawa, H., Yamaoka, K., Yamamoto, Y., et al. (2007b). 1Alpha,25(OH)2D3-induced transrepression by vitamin D receptor through E-box-type elements in the human parathyroid hormone gene promoter. *Mol. Endocrinol.* 21, 334–342. doi: 10.1210/me.2006-0231
- Kim, S., Li, M., Paik, H., Nephew, K., Shi, H., Kramer, R., et al. (2008). Predicting DNA methylation susceptibility using CpG flanking sequences. *Pac. Symp. Biocomput.* 13, 315–326.
- Klose, R. J., Yamane, K., Bae, Y., Zhang, D., Erdjument-Bromage, H., Tempst, P., et al. (2006). The transcriptional repressor JHDM3A demethylates trimethyl histone H3 lysine 9 and lysine 36. *Nature* 442, 312–316. doi: 10.1038/nature04853
- Klose, R. J., Yan, Q., Tothova, Z., Yamane, K., Erdjument-Bromage, H., Tempst, P., et al. (2007). The retinoblastoma binding protein RBP2 is an H3K4 demethylase. *Cell* 128, 889–900. doi: 10.1016/j.cell.2007.02.013
- Kouzarides, T. (2007). Chromatin modifications and their function. *Cell* 128, 693–705. doi: 10.1016/j.cell.2007.02.005
- Kriaucionis, S., and Heintz, N. (2009). The nuclear DNA base 5-hydroxymethylcytosine is present in Purkinje neurons and the brain. *Science* 324, 929–930. doi: 10.1126/science.1169786
- Lagger, G., Doetzlhofer, A., Schuettengruber, B., Haidweger, E., Simboeck, E., Tischler, J., et al. (2003). The tumor suppressor p53 and histone deacetylase 1 are antagonistic regulators of the cyclin-dependent kinase inhibitor p21/WAF1/CIP1 gene. *Mol. Cell Biol.* 23, 2669–2679. doi: 10.1128/MCB.23.8.2669-2679.2003
- Lee, J. W., Lee, Y. C., Na, S. Y., Jung, D. J., and Lee, S. K. (2001). Transcriptional coregulators of the nuclear receptor superfamily: coactivators and corepressors. *Cell Mol. Life Sci.* 58, 289–297. doi: 10.1007/PL00000856
- Lewin, E., Garfia, B., Recio, F. L., Rodriguez, M., and Olgaard, K. (2002). Persistent downregulation of calcium-sensing receptor mRNA in rat parathyroids when severe secondary hyperparathyroidism is reversed by an isogenic kidney transplantation. *J. Am. Soc. Nephrol.* 13, 2110–2116. doi: 10.1097/01.ASN.0000024439.38838.03
- Liau, J. Y., Liao, S. L., Hsiao, C. H., Lin, M. C., Chang, H. C., and Kuo, K. T. (2014). Hypermethylation of the CDKN2A gene promoter is a frequent epigenetic change in periocular sebaceous carcinoma and is associated with younger patient age. *Hum Pathol.* 45, 533–539. doi: 10.1016/j.humpath.2013.10.019
- Lin, C. H., Li, B., Swanson, S., Zhang, Y., Florens, L., Washburn, M. P., et al. (2008). Heterochromatin protein 1a stimulates histone H3 lysine 36 demethylation by the Drosophila KDM4A demethylase. *Mol. Cell* 32, 696–706. doi: 10.1016/j.molcel.2008.11.008
- Lin, Y. W., Ren, L. L., Xiong, H., Du, W., Yu, Y. N., Sun, T. T., et al. (2013). Role of STAT3 and vitamin D receptor in EZH2-mediated invasion of human colorectal cancer. *J. Pathol.* 230, 277–290. doi: 10.1002/path.4179
- Longo, V. D., and Kennedy, B. K. (2006). Sirtuins in aging and age-related disease. *Cell* 126, 257–268. doi: 10.1016/j.cell.2006.07.002
- Lopes, N., Carvalho, J., Duraes, C., Sousa, B., Gomes, M., Costa, J. L., et al. (2012). 1Alpha,25-dihydroxyvitamin D3 induces de novo E-cadherin expression in triple-negative breast cancer cells by CDH1-promoter demethylation. *Anticancer Res.* 32, 249–257.
- Lu, P. J., Sundquist, K., Baekstrom, D., Poulsom, R., Hanby, A., Meier-Ewert, S., et al. (1999). A novel gene (PLU-1) containing highly conserved putative DNA/chromatin binding motifs is specifically up-regulated in breast cancer. *J. Biol. Chem.* 274, 15633–15645. doi: 10.1074/jbc.274.22.15633
- Luo, W., Karpf, A. R., Deeb, K. K., Muindi, J. R., Morrison, C. D., Johnson, C. S., et al. (2010). Epigenetic regulation of vitamin D 24-hydroxylase/CYP24A1 in human prostate cancer. *Cancer Res.* 70, 5953–5962. doi: 10.1158/0008-5472.CAN-10-0617
- Marik, R., Fackler, M., Gabrielson, E., Zeiger, M. A., Sukumar, S., Stearns, V., et al. (2010). DNA methylation-related vitamin D receptor insensitivity in breast cancer. *Cancer Biol. Ther.* 10, 44–53. doi: 10.4161/cbt.10.1.11994
- Marks, P., Rifkind, R. A., Richon, V. M., Breslow, R., Miller, T., and Kelly, W. K. (2001). Histone deacetylases and cancer: causes and therapies. *Nat. Rev. Cancer* 1, 194–202. doi: 10.1038/35106079
- Meissner, A., Mikkelsen, T. S., Gu, H., Wernig, M., Hanna, J., Sivachenko, A., et al. (2008). Genome-scale DNA methylation maps of pluripotent and differentiated cells. *Nature* 454, 766–770. doi: 10.1038/nature07107
- Metzger, E., Wissmann, M., Yin, N., Müller, J. M., Schneider, R., Peters, A. H., et al. (2005). LSD1 demethylates repressive histone marks to promote androgen-receptor-dependent transcription. *Nature* 437, 436–439. doi: 10.1038/nature04020
- Meyer, M. B., Benkusky, N. A., and Pike, J. W. (2013). 1,25-Dihydroxyvitamin D3 induced histone profiles guide discovery of VDR action sites. *J. Steroid Biochem. Mol. Biol.* doi: 10.1016/j.jsbmb.2013.09.005. [Epub ahead of print].
- Miller, C. W., Morosetti, R., Campbell, M. J., Mendoza, S., and Koeffler, H. P. (1997). Integrity of the 1,25-dihydroxyvitamin D3 receptor in bone, lung, and other cancers. *Mol. Carcinog.* 19, 254–257. doi: 10.1002/(SICI)1098-2744(199708)19:4<254::AID-MC6>3.0.CO;2-C
- Mosammamaparast, N., and Shi, Y. (2010). Reversal of histone methylation: biochemical and molecular mechanisms of histone demethylases. *Annu. Rev. Biochem.* 79, 155–179. doi: 10.1146/annurev.biochem.78.070907.103946
- Mottet, D., Pirotte, S., Lamour, V., Hagedorn, M., Javerzat, S., Bikfalvi, A., et al. (2009). HDAC4 represses p21(WAF1/Cip1) expression in human cancer cells through a Sp1-dependent, p53-independent mechanism. *Oncogene* 28, 243–256. doi: 10.1038/onc.2008.371
- Murayama, A., Kim, M. S., Yanagisawa, J., Takeyama, K., and Kato, S. (2004). Transrepression by a liganded nuclear receptor via a bHLH activator through co-regulator switching. *EMBO J.* 23, 1598–1608. doi: 10.1038/sj.emboj.7600157
- Nagpal, S., Na, S., and Rathnachalam, R. (2005). Noncalcemic actions of vitamin D receptor ligands. *Endocr. Rev.* 26, 662–687. doi: 10.1210/er.2004-0002
- Nan, X., Ng, H. H., Johnson, C. A., Laherty, C. D., Turner, B. M., Eisenman, R. N., et al. (1998). Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a histone deacetylase complex. *Nature* 393, 386–389. doi: 10.1038/30764
- Norman, A. W., and Bouillon, R. (2010). Vitamin D nutritional policy needs a vision for the future. *Exp. Biol. Med. (Maywood)* 235, 1034–1045. doi: 10.1258/ebm.2010.010014
- Novakovic, B., Sibson, M., Ng, H. K., Manuelpillai, U., Rakyar, V., Down, T., et al. (2009). Placenta-specific methylation of the vitamin D 24-hydroxylase gene: implications for feedback autoregulation of active vitamin D levels at the fetomaternal interface. *J. Biol. Chem.* 284, 14838–14848. doi: 10.1074/jbc.M809542200
- Ohyama, Y., Kusada, T., Yamasaki, T., and Ide, H. (2002). Extensive methylation of CpG island of CYP24 gene in osteoblastic ROS17/2.8 cells. *Nucleic Acids Res. Suppl.* 249–250. doi: 10.1093/nass/2.1.249
- Okano, M., Bell, D. W., Haber, D. A., and Li, E. (1999). DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell* 99, 247–257. doi: 10.1016/S0092-8674(00)81656-6
- Okano, M., Xie, S., and Li, E. (1998). Cloning and characterization of a family of novel mammalian DNA (cytosine-5) methyltransferases. *Nat. Genet.* 19, 219–220. doi: 10.1038/890
- Padi, S. K., Zhang, Q., Rustum, Y. M., Morrison, C., and Guo, B. (2013). MicroRNA-627 mediates the epigenetic mechanisms of vitamin D to suppress proliferation of human colorectal cancer cells and growth of xenograft tumors in mice. *Gastroenterology* 145, 437–446. doi: 10.1053/j.gastro.2013.04.012
- Pan, L., Matloob, A. F., Du, J., Pan, H., Dong, Z., Zhao, J., et al. (2010). Vitamin D stimulates apoptosis in gastric cancer cells in synergy with trichostatin A /sodium butyrate-induced and 5-aza-2'-deoxycytidine-induced PTEN upregulation. *FEBS J.* 277, 989–999. doi: 10.1111/j.1742-4658.2009.07542.x

- Parbin, S., Kar, S., Shilpi, A., Sengupta, D., Deb, M., Rath, S. K., et al. (2014). Histone deacetylases: a saga of perturbed acetylation homeostasis in cancer. *J. Histochem. Cytochem.* 62, 11–33. doi: 10.1369/0022155413506582
- Pedersen, J. I., Shobaki, H. H., Holmberg, I., Bergseth, S., and Björkhem, I. (1983). 25-Hydroxyvitamin D₃-24-hydroxylase in rat kidney mitochondria. *J. Biol. Chem.* 258, 742–746.
- Pedersen, M. T., and Helin, K. (2010). Histone demethylases in development and disease. *Trends Cell Biol.* 20, 662–671. doi: 10.1016/j.tcb.2010.08.011
- Pendas-Franco, N., Aguilera, O., Pereira, F., Gonzalez-Sancho, J. M., and Munoz, A. (2008). Vitamin D and Wnt/beta-catenin pathway in colon cancer: role and regulation of DICKKOPF genes. *Anticancer Res.* 28, 2613–2623.
- Pereira, F., Barbachano, A., Silva, J., Bonilla, F., Campbell, M. J., Munoz, A., et al. (2011). KDM6B/JMJD3 histone demethylase is induced by vitamin D and modulates its effects in colon cancer cells. *Hum. Mol. Genet.* 20, 4655–4665. doi: 10.1093/hmg/ddr399
- Pereira, F., Barbachano, A., Singh, P. K., Campbell, M. J., Munoz, A., and Larriba, M. J. (2012). Vitamin D has wide regulatory effects on histone demethylase genes. *Cell Cycle* 11, 1081–1089. doi: 10.4161/cc.11.6.19508
- Perillo, B., Ombra, M. N., Bertoni, A., Cuzzo, C., Sacchetti, S., Sasso, A., et al. (2008). DNA oxidation as triggered by H3K9me2 demethylation drives estrogen-induced gene expression. *Science* 319, 202–206. doi: 10.1126/science.1147674
- Pfau, R., Tzatsos, A., Kampranis, S. C., Serebrennikova, O. B., Bear, S. E., and Tschlis, P. N. (2008). Members of a family of JmjC domain-containing oncoproteins immortalize embryonic fibroblasts via a JmjC domain-dependent process. *Proc. Natl. Acad. Sci. U.S.A.* 105, 1907–1912. doi: 10.1073/pnas.0711865105
- Pike, J. W., and Meyer, M. B. (2013). Fundamentals of vitamin D hormone-regulated gene expression. *J. Steroid Biochem. Mol. Biol.* doi: 10.1016/j.jsbmb.2013.11.004. [Epub ahead of print].
- Pike, J. W., Meyer, M. B., and Bishop, K. A. (2012). Regulation of target gene expression by the vitamin D receptor - an update on mechanisms. *Rev. Endocr. Metab. Disord.* 13, 45–55. doi: 10.1007/s11154-011-9198-9
- Polly, P., Herdick, M., Moehren, U., Banihmad, A., Heinzl, T., and Carlberg, C. (2000). VDR-Alien: a novel, DNA-selective vitamin D(3) receptor-corepressor partnership. *FASEB J.* 14, 1455–1463. doi: 10.1096/fj.14.10.1455
- Rashid, S. F., Moore, J. S., Walker, E., Driver, P. M., Engel, J., Edwards, C. E., et al. (2001). Synergistic growth inhibition of prostate cancer cells by 1 alpha,25 Dihydroxyvitamin D(3) and its 19-nor-hexafluoride analogs in combination with either sodium butyrate or trichostatin A. *Oncogene* 20, 1860–1872. doi: 10.1038/sj.onc.1204269
- Rawson, J. B., Sun, Z., Dicks, E., Daftary, D., Parfrey, P. S., Green, R. C., et al. (2012). Vitamin D intake is negatively associated with promoter methylation of the Wnt antagonist gene DKK1 in a large group of colorectal cancer patients. *Nutr. Cancer* 64, 919–928. doi: 10.1080/01635581.2012.711418
- Reik, W., and Lewis, A. (2005). Co-evolution of X-chromosome inactivation and imprinting in mammals. *Nat. Rev. Genet.* 6, 403–410. doi: 10.1038/nrg1602
- Rhee, I., Bachman, K. E., Park, B. H., Jair, K. W., Yen, R. W., Schuebel, K. E., et al. (2002). DNMT1 and DNMT3b cooperate to silence genes in human cancer cells. *Nature* 416, 552–556. doi: 10.1038/416552a
- Rhee, I., Jair, K. W., Yen, R. W., Lengauer, C., Herman, J. G., Kinzler, K. W., et al. (2000). CpG methylation is maintained in human cancer cells lacking DNMT1. *Nature* 404, 1003–1007. doi: 10.1038/35010000
- Robertson, K. D. (2005). DNA methylation and human disease. *Nat. Rev. Genet.* 6, 597–610. doi: 10.1038/nrg1655
- Saito, Y., Liang, G., Egger, G., Friedman, J. M., Chuang, J. C., Coetzee, G. A., et al. (2006). Specific activation of microRNA-127 with downregulation of the proto-oncogene BCL6 by chromatin-modifying drugs in human cancer cells. *Cancer Cell* 9, 435–443. doi: 10.1016/j.ccr.2006.04.020
- Sakai, T., Toguchida, J., Ohtani, N., Yandell, D. W., Rapaport, J. M., and Dryja, T. P. (1991). Allele-specific hypermethylation of the retinoblastoma tumor-suppressor gene. *Am. J. Hum. Genet.* 48, 880–888.
- Sakaki, T., Sawada, N., Komai, K., Shiozawa, S., Yamada, S., Yamamoto, K., et al. (2000). Dual metabolic pathway of 25-hydroxyvitamin D₃ catalyzed by human CYP24. *Eur. J. Biochem.* 267, 6158–6165. doi: 10.1046/j.1432-1327.2000.01680.x
- Saramäki, A., Diermeier, S., Kellner, R., Laitinen, H., Väisänen, S., and Carlberg, C. (2009). Cyclical chromatin looping and transcription factor association on the regulatory regions of the p21 (CDKN1A) gene in response to 1alpha,25-dihydroxyvitamin D₃. *J. Biol. Chem.* 284, 8073–8082. doi: 10.1074/jbc.M808090200
- Schulte, J. H., Lim, S., Schramm, A., Friedrichs, N., Koster, J., Versteeg, R., et al. (2009). Lysine-specific demethylase 1 is strongly expressed in poorly differentiated neuroblastoma: implications for therapy. *Cancer Res.* 69, 2065–2071. doi: 10.1158/0008-5472.CAN-08-1735
- Schwer, B., and Verdin, E. (2008). Conserved metabolic regulatory functions of sirtuins. *Cell Metab.* 7, 104–112. doi: 10.1016/j.cmet.2007.11.006
- Seuter, S., Pehkonen, P., Heikkinen, S., and Carlberg, C. (2013). Dynamics of 1alpha,25-dihydroxyvitamin D₃-dependent chromatin accessibility of early vitamin D receptor target genes. *Biochim. Biophys. Acta* 1829, 1266–1275. doi: 10.1016/j.bbagr.2013.10.003
- Shi, H., Guo, J., Duff, D. J., Rahmatpanah, F., Chitima-Matsiga, R., Al-Kuhlani, M., et al. (2007). Discovery of novel epigenetic markers in non-Hodgkin's lymphoma. *Carcinogenesis* 28, 60–70. doi: 10.1093/carcin/bgl092
- Shi, H., Yan, P. S., Chen, C. M., Rahmatpanah, F., Lofton-Day, C., Caldwell, C. W., et al. (2002). Expressed CpG island sequence tag microarray for dual screening of DNA hypermethylation and gene silencing in cancer cells. *Cancer Res.* 62, 3214–3220.
- Shi, Y., Lan, F., Matson, C., Mulligan, P., Whetstone, J. R., Cole, P. A., et al. (2004). Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. *Cell* 119, 941–953. doi: 10.1016/j.cell.2004.12.012
- Shi, Y., and Whetstone, J. R. (2007). Dynamic regulation of histone lysine methylation by demethylases. *Mol. Cell* 25, 1–14. doi: 10.1016/j.molcel.2006.12.010
- Singh, P. K., Doig, C. L., Dhiman, V. K., Turner, B. M., Smiraglia, D. J., and Campbell, M. J. (2013). Epigenetic distortion to VDR transcriptional regulation in prostate cancer cells. *J. Steroid Biochem. Mol. Biol.* 136, 258–263. doi: 10.1016/j.jsbmb.2012.10.002
- Sulaiman, L., Juhlin, C. C., Nilsson, I. L., Fotouhi, O., Larsson, C., and Hashemi, J. (2013). Global and gene-specific promoter methylation analysis in primary hyperparathyroidism. *Epigenetics* 8, 646–655. doi: 10.4161/epi.24823
- Tahiliani, M., Mei, P., Fang, R., Leonor, T., Rutenberg, M., Shimizu, F., et al. (2007). The histone H3K4 demethylase SMCX links REST target genes to X-linked mental retardation. *Nature* 447, 601–605. doi: 10.1038/nature05823
- Takeyama, K., and Kato, S. (2011). The vitamin D₃ 1alpha-hydroxylase gene and its regulation by active vitamin D₃. *Biosci. Biotechnol. Biochem.* 75, 208–213. doi: 10.1271/bbb.100684
- Tan, J., Lu, J., Huang, W., Dong, Z., Kong, C., Li, L., et al. (2009). Genome-wide analysis of histone H3 lysine9 modifications in human mesenchymal stem cell osteogenic differentiation. *PLoS ONE* 4:e6792. doi: 10.1371/journal.pone.0006792
- Tapp, H. S., Commane, D. M., Bradburn, D. M., Arasaradnam, R., Mathers, J. C., Johnson, I. T., et al. (2013). Nutritional factors and gender influence age-related DNA methylation in the human rectal mucosa. *Aging Cell* 12, 148–155. doi: 10.1111/acer.12030
- Ting, A. H., McGarvey, K. M., and Baylin, S. B. (2006). The cancer epigenome—components and functional correlates. *Genes Dev.* 20, 3215–3231. doi: 10.1101/gad.1464906
- Tsai, H. C., and Baylin, S. B. (2011). Cancer epigenetics: linking basic biology to clinical medicine. *Cell Res.* 21, 502–517. doi: 10.1038/cr.2011.24
- Tsukada, Y., Fang, J., Erdjument-Bromage, H., Warren, M. E., Borchers, C. H., Tempst, P., et al. (2006). Histone demethylation by a family of JmjC domain-containing proteins. *Nature* 439, 811–816. doi: 10.1038/nature04433
- Vanoirbeek, E., Eelen, G., Verlinden, L., Carmeliet, G., Mathieu, C., Bouillon, R., et al. (2014). PDLIM2 expression is driven by vitamin D and is involved in the pro-adhesion, and anti-migration and -invasion activity of vitamin D. *Oncogene* 33, 1904–1911. doi: 10.1038/onc.2013.123
- Varshney, S., Bhadada, S. K., Sachdeva, N., Arya, A. K., Saikia, U. N., Behera, A., et al. (2013). Methylation status of the CpG islands in vitamin D and calcium-sensing receptor gene promoters does not explain the reduced gene expressions in parathyroid adenomas. *J. Clin. Endocrinol. Metab.* 98, E1631–E1635. doi: 10.1210/jc.2013-1699
- Von Essen, M. R., Kongsbak, M., Schjerling, P., Olgaard, K., Odum, N., and Geisler, C. (2010). Vitamin D controls T cell antigen receptor signaling and activation of human T cells. *Nat. Immunol.* 11, 344–349. doi: 10.1038/ni.1851
- Wang, Y., and Leung, F. C. (2004). An evaluation of new criteria for CpG islands in the human genome as gene markers. *Bioinformatics* 20, 1170–1177. doi: 10.1093/bioinformatics/bth059
- Whetstone, J. R., Nottke, A., Lan, F., Huarte, M., Smolnikov, S., Chen, Z., et al. (2006). Reversal of histone lysine trimethylation by the JMJD2 family of histone demethylases. *Cell* 125, 467–481. doi: 10.1016/j.cell.2006.03.028

- Williams, K., Christensen, J., Pedersen, M. T., Johansen, J. V., Cloos, P. A., Rappsilber, J., et al. (2011). TET1 and hydroxymethylcytosine in transcription and DNA methylation fidelity. *Nature* 473, 343–348. doi: 10.1038/nature10066
- Wilson, A. J., Byun, D. S., Popova, N., Murray, L. B., L'Italien, K., Sowa, Y., et al. (2006). Histone deacetylase 3 (HDAC3) and other class I HDACs regulate colon cell maturation and p21 expression and are deregulated in human colon cancer. *J. Biol. Chem.* 281, 13548–13558. doi: 10.1074/jbc.M510023200
- Wjst, M., Heimbeck, I., Kutschke, D., and Pukelsheim, K. (2010). Epigenetic regulation of vitamin D converting enzymes. *J. Steroid Biochem. Mol. Biol.* 121, 80–83. doi: 10.1016/j.jsbmb.2010.03.056
- Xiang, Y., Zhu, Z., Han, G., Ye, X., Xu, B., Peng, Z., et al. (2007). JARID1B is a histone H3 lysine 4 demethylase up-regulated in prostate cancer. *Proc. Natl. Acad. Sci. U.S.A.* 104, 19226–19231. doi: 10.1073/pnas.0700735104
- Yamaguchi, S., Hong, K., Liu, R., Shen, L., Inoue, A., Diep, D., et al. (2012). Tet1 controls meiosis by regulating meiotic gene expression. *Nature* 492, 443–447. doi: 10.1038/nature11709
- Yamane, K., Tateishi, K., Klose, R. J., Fang, J., Fabrizio, L. A., Erdjument-Bromage, H., et al. (2007). PLU-1 is an H3K4 demethylase involved in transcriptional repression and breast cancer cell proliferation. *Mol. Cell* 25, 801–812. doi: 10.1016/j.molcel.2007.03.001
- Yamshchikov, A. V., Desai, N. S., Blumberg, H. M., Ziegler, T. R., and Tangpricha, V. (2009). Vitamin D for treatment and prevention of infectious diseases: a systematic review of randomized controlled trials. *Endocr. Pract.* 15, 438–449. doi: 10.4158/EP09101.ORR
- Yoneda, T., Aya, S., and Sakuda, M. (1984). Sodium butyrate (SB) augments the effects of 1,25 dihydroxyvitamin D3 (1,25(OH)2D3) on neoplastic and osteoblastic phenotype in clonal rat osteosarcoma cells. *Biochem. Biophys. Res. Commun.* 121, 796–801. doi: 10.1016/0006-291X(84)90748-4
- Zhang, X., Nicosia, S. V., and Bai, W. (2006). Vitamin D receptor is a novel drug target for ovarian cancer treatment. *Curr. Cancer Drug. Targets* 6, 229–244. doi: 10.2174/156800906776842939
- Zhang, Z., Yamashita, H., Toyama, T., Sugiura, H., Ando, Y., Mita, K., et al. (2005). Quantitation of HDAC1 mRNA expression in invasive carcinoma of the breast*. *Breast Cancer Res. Treat.* 94, 11–16. doi: 10.1007/s10549-005-6001-1
- Zhu, H., Wang, X., Shi, H., Su, S., Harshfield, G. A., Gutin, B., et al. (2013). A genome-wide methylation study of severe vitamin D deficiency in African American adolescents. *J. Pediatr.* 162, 1004–1009.e1001. doi: 10.1016/j.jpeds.2012.10.059
- Zhu, P., Martin, E., Mengwasser, J., Schlag, P., Janssen, K. P., and Göttlicher, M. (2004). Induction of HDAC2 expression upon loss of APC in colorectal tumorigenesis. *Cancer Cell* 5, 455–463. doi: 10.1016/S1535-6108(04)00114-X

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 09 February 2014; paper pending published: 07 March 2014; accepted: 09 April 2014; published online: 29 April 2014.

Citation: Fetahu IS, Höbaus J and Kállay E (2014) Vitamin D and the epigenome. *Front. Physiol.* 5:164. doi: 10.3389/fphys.2014.00164

This article was submitted to *Integrative Physiology*, a section of the journal *Frontiers in Physiology*.

Copyright © 2014 Fetahu, Höbaus and Kállay. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.