

Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Essential sufficiency of zinc, ω -3 polyunsaturated fatty acids, vitamin D and magnesium for prevention and treatment of COVID-19, diabetes, cardiovascular diseases, lung diseases and cancer

Michael J. Story

PII: S0300-9084(21)00140-1

DOI: https://doi.org/10.1016/j.biochi.2021.05.013

Reference: BIOCHI 6136

To appear in: *Biochimie*

Received Date: 22 February 2021

Revised Date: 19 May 2021

Accepted Date: 25 May 2021

Please cite this article as: M.J. Story, Essential sufficiency of zinc, ω-3 polyunsaturated fatty acids, vitamin D and magnesium for prevention and treatment of COVID-19, diabetes, cardiovascular diseases, lung diseases and cancer, *Biochimie* (2021), doi: https://doi.org/10.1016/j.biochi.2021.05.013.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier B.V.



Essential sufficiency of zinc, ω -3 polyunsaturated fatty acids, vitamin D and magnesium for prevention and treatment of COVID-19, diabetes, cardiovascular diseases, lung diseases and cancer

Abstract

Despite the development of a number of vaccines for COVID-19, there remains a need for prevention and treatment of the virus SARS-CoV-2 and the ensuing disease COVID-19. This report discusses the key elements of SARS-CoV-2 and COVID-19 that can be readily treated: viral entry, the immune system and inflammation, and the cytokine storm. It is shown that the essential nutrients zinc, ω -3 polyunsaturated fatty acids (PUFAs), vitamin D and magnesium provide the ideal combination for prevention and treatment of COVID-19: prevention of SARS-CoV-2 entry to host cells, prevention of proliferation of SARS-CoV-2, inhibition of excessive inflammation, improved control of the regulation of the immune system, inhibition of the cytokine storm, and reduction in the effects of acute respiratory distress syndrome (ARDS) and associated noncommunicable diseases. It is emphasized that the non-communicable diseases associated with COVID-19 are inherently more prevalent in the elderly than the young, and that the maintenance of sufficiency of zinc, ω -3 PUFAs, vitamin D and magnesium is essential for the elderly to prevent the occurrence of noncommunicable diseases such as diabetes, cardiovascular diseases, lung diseases and cancer. Annual checking of levels of these essential nutrients is recommended for those over 65 years of age, together with appropriate adjustments in their intake, with these services and supplies being at government cost. The cost:benefit ratio would be huge as the cost of the nutrients and the testing of their levels would be very small compared with the cost savings of specialists and hospitalization.

Essential sufficiency of zinc, ω -3 polyunsaturated fatty acids, vitamin D and magnesium for prevention and treatment of COVID-19, diabetes, cardiovascular diseases, lung diseases and cancer

Michael J Story PhD

Story Pharmaceutics Pty Ltd, PO Box 6086, Linden Park, South Australia, 5065, Australia

storypharm@bigpond.com

Abstract

Despite the development of a number of vaccines for COVID-19, there remains a need for prevention and treatment of the virus SARS-CoV-2 and the ensuing disease COVID-19. This report discusses the key elements of SARS-CoV-2 and COVID-19 that can be readily treated: viral entry, the immune system and inflammation, and the cytokine storm. It is shown that the essential nutrients zinc, ω -3 polyunsaturated fatty acids (PUFAs), vitamin D and magnesium provide the ideal combination for prevention and treatment of COVID-19: prevention of SARS-CoV-2 entry to host cells, prevention of proliferation of SARS-CoV-2, inhibition of excessive inflammation, improved control of the regulation of the immune system, inhibition of the cytokine storm, and reduction in the effects of acute respiratory distress syndrome (ARDS) and associated noncommunicable diseases. It is emphasized that the non-communicable diseases associated with COVID-19 are inherently more prevalent in the elderly than the young, and that the maintenance of sufficiency of zinc, ω -3 PUFAs, vitamin D and magnesium is essential for the elderly to prevent the occurrence of noncommunicable diseases such as diabetes, cardiovascular diseases, lung diseases and cancer. Annual checking of levels of these essential nutrients is recommended for those over 65 years of age, together with appropriate adjustments in their intake, with these services and supplies being at government cost. The cost:benefit ratio would be huge as the cost of the nutrients and the testing of their levels would be very small compared with the cost savings of specialists and hospitalization.

Keywords: zinc, ω -3 PUFAs, vitamin D, magnesium, essentiality, SARS-CoV-2, COVID-19, diabetes, cardiovascular diseases, lung diseases, cancer, supplementation

Abbreviations

1a,25(OH) ₂ D ₃ 25(OH)D ACE ACE2 ADAM17	1a,25-dihydroxyvitamin D ₃ , calcitriol 25-hydroxyvitamin D ₃ angiotensin converting enzyme angiotensin converting enzyme 2 disintegrin and metalloproteinase	IL-x IL-1RN IP-10 M1,2 MMP-2,9	interleukin x IL-1 receptor antagonist protein interferon- γ -inducible protein macrophage type 1, (or 2) matrix metallopeptidase 2, (or 9)
	domain 17	NF-vB	nuclear factor k-light-chain-enhancer
			of activated B cells
Ang	angiotensin	РЗ8 МАРК	p38 mitogen-activated protein kinases
ARDS	acute respiratory distress syndrome	PDGF	platelet-derived growth factor
ATR1	AT1 receptor	PGE2	prostaglandin E ₂
CCL2 (MCP-1)	monocyte chemotactic protein-1	PUFA	polyunsaturated fatty acid
CCL3	macrophage inflammatory protein	RANTES	regulated upon activation, normal T
(MIP-1a)	1a		cell expressed and presumably secreted
c-Kit	stem cell factor receptor	RAS	renin-angiotensin system
CRP	C-reactive protein	ROS	reactive oxygen species
CYP27B1	1a-hydroxylase	SAA	serum amyloid A
DHA	docosahexaenoic acid	SCF	stem cell factor
EPA	eicosapentaenoic acid	TGF-B	transforming growth factor B
ERK1/2	extracellular signal-regulated kinase 1/2	Th	T helper
FGF	fibroblast growth factor	TLR	toll-like receptor
G-CSF	granulocyte colony-stimulating factor	TMPRSS2	transmembrane serine protease 2
GM-CSF	granulocyte-macrophage colony- stimulating factor	TNF-a	tumour necrosis factor a
HCQ	hydroxychloroquine	Treg	regulatory T cell
HIF-1a	hypoxia-inducible factor 1-a	VDR	vitamin D receptor
IFN-γ	interferon-γ	VEGF	vascular endothelial growth factor
			_

1. Introduction

SARS-CoV is the original SARS that was virulent in the early 2000s; it is the virus that leads to the COVID-19 disease state. There is an immediate need for a prophylaxis and treatment for COVID-19. Even if the vaccines that are being distributed and further developed are successful in their efficacy and prevention of transference, there will always be a need for a prophylaxis and treatment for like viruses and comorbidities, particularly in poorer countries. It is therefore timely to present the advantages of the essential combination of zinc, ω -3 PUFAs, vitamin D and magnesium as these treatments are low in cost, extensive in their actions, and safe as they are naturally present in the human body.

The essential sufficiency of zinc, ω -3 PUFAs and vitamin D for prevention and treatment of cancers has been previously presented [1]. The prevention and treatment of COVID-19 and associated comorbidities is currently of the highest concern. Furthermore, the four principal non-communicable diseases that are

most prevalent world-wide are diabetes, cardiovascular diseases, lung diseases and cancer [2]. The comorbidities discussed here in the context of COVID-19 include these non-communicable diseases as well as ageing and obesity. Zinc, ω -3 PUFAs, vitamin D and magnesium, although very different in their modes of action, have many similar end-effects. The commonality of zinc, ω -3 PUFAs, vitamin D and magnesium in inhibiting inflammatory situations, as well as their ability to correct immune dysfunction, together with the low cost and safety of these nutrients, makes them ideal as front-line prevention measures and adjuvants in treating COVID-19 and associated comorbidities, as well as the principal non-communicable diseases in non-COVID situations.

Extensive literature searches were made in preparation of this manuscript. PubMed was principally used with a wide range of search strings. Separate searches were made for key concepts together with each of the individual components: ie, zinc; (PUFAs OR DHA OR EPA); vitamin D; magnesium. A PubMed search was conducted using the following search string: (COVID-19 OR SARS-CoV-2 OR coronavirus) AND zinc AND (PUFA OR DHA OR EPA) AND "vitamin D" AND magnesium. This search produced zero results suggesting that there have been virtually no studies examining the beneficial effects of supplementation of all four essential components together: zinc, ω -3 PUFAs, vitamin D and magnesium. Nevertheless, the four components are described or mentioned in a number of reviews of nutritional status and immune function and/or inflammation [eg, 3-6].

2. Zinc

Zinc is a vital cofactor for more than 2000 transcription factors and 300 enzymes in regulating cell differentiation and proliferation, as well as basic metabolic functions of cells [7]. Zinc deficiency is a world-wide problem with approximately 2 billion people subjected to zinc-deficient diets [7,8]. Zinc deficiency is not limited to the developing countries as it also exists in the industrialized world, mainly in the elderly [9,10]. In normal healthy adults, the plasma concentration of zinc is typically 14-23 μ mol/L (0.9-1.5 μ g/mL) [11].

Risk factors associated with Zn deficiency have been well reported in the literature [12-14]. Zinc deficiency can cause an imbalance in both the innate and adaptive immune systems, with severe deficiency leading to infections, skin disorders, gastrointestinal disorders, weight loss, growth retardation and male hypogonadism, amongst other symptoms [14-17]. Low zinc levels have been found to affect the function of various types of immune cells, including macrophages, neutrophils, mast cells and dendritic cells [11,18]. Zinc is also essential for the development, differentiation and activation of T cells [19]. Zinc deficiency can therefore result in impaired production, activation and maturation of natural killer cells (cell-mediated innate immunity), T cells (cell-mediated adaptive immunity) and B cells (humoral adaptive immunity) [5].

Zinc deficiency, which is commonly reported in the elderly, lowers immune function, decreases resistance to invading pathogens and increases the risk of contracting pneumonia [20,21]. Zinc deficiency also occurs frequently in patients with cardiovascular disease, chronic pulmonary disease, diabetes or obesity [22,23].

Zinc exerts its anti-inflammatory activity by inhibiting activation and signalling of NF- κ B, and through controlling regulatory T cell (Treg) functions [12]. Zinc supplementation causes pro-inflammatory Th17 cells to skew towards anti-inflammatory Treg cells [24]. Zinc supplementation inhibits NF- κ B through expression of the A20 protein, resulting in suppression of TNF-a, IL-1 β , IL-6, IL-8, IL-12, IL-18, IFN- γ , iNOS, COX-2, GM-CSF [12,25].

Macrophages, neutrophils, and T cells are activated through elevation of cytokines including IL-1, IL-6, and TNF-a, often leading to acute respiratory distress syndrome (ARDS) [26]. IL-6, IL-8 and TNF-a levels are elevated in elderly people who are zinc deficient, as well as in the obese [17,27], and zinc supplementation has been found to reduce these levels [10].

When levels of reactive oxygen species (ROS) are elevated, as is common in zinc deficiency, oxidative damage results. Zinc supplementation decreases ROS production and this has beneficial results, especially in the aged and in diabetes mellitus [13].

A number of comprehensive reviews of zinc and its involvement in ageing, COVID-type viruses and comorbidities have been presented [eg, 27,28]. Zinc sufficiency is vital for reducing risk factors associated with COVID-19, such as obesity, diabetes, cardiovascular diseases, lung diseases and ageing [12,29]. Physiological supplementation of Zn in ageing and in age-related degenerative diseases corrects immune defects and reduces infection relapse [30].

One of the problems of zinc supplementation has been the variability of zinc bioavailability in cells. It has been found that increasing the intracellular levels of zinc using ionophores such as pyrithione can effectively decrease the replication of a variety of viruses, including the replication of SARS-CoV [31]. Unfortunately, pyrithione is not recommended for use internally, whereas it is efficacious and safe when used topically. Other proven zinc ionophores include chloroquine and hydroxychloroquine (HCQ) [12,32,33], disulfiram [33], quercetin and epigallocatechin-gallate [34], and zincophorin [35]. In addition, Rizzo [36] presented a sound rationale for ivermectin being an ionophore for zinc. A number of clinical studies are planned or underway that are based on HCQ and zinc, and ivermectin and zinc, together in some cases with an antibiotic such as azithromycin or doxycycline [37]. Interestingly, the HCQ studies are fundamentally based on testing whether zinc complements HCQ, and not whether HCQ complements zinc which would be expected if HCQ was recognized as an ionophore for zinc. A similar comment applies to ivermectin. The latest summary of ongoing studies at the time of preparing this manuscript was that of Pal et al [38].

Zinc is also known for its capacity to modulate antiviral and antibacterial immunity [12]. The antibacterial properties of zinc are well demonstrated against *S. pneumoniae* [12]. Moreover, zinc has the capacity to reduce the risk of bacterial co-infection by improving lung function through mucociliary clearance and protecting lung barrier function.

Zinc inhibits SARS-CoV RNA polymerase, and thus its replication capacity [17]. Zinc has also been postulated as a stabilizer of cell membranes which could assist in blocking virus entry to cells [39]. In this context, zinc decreases activity

of angiotensin converting enzyme 2 (ACE2) which is required for binding with SARS-CoV-2 for cell entry [12]. Wessels and co-workers [40] concluded that zinc has multiple functions in inhibiting viral entry to host cells and activity: prevention of fusion with the host membrane, inhibiting viral polymerase and ensuing replication, impairing protein translation and processing, blocking viral particle release, and destabilizing the viral envelope. It has been shown that zinc deficiency increases the leakage of the epithelium of the respiratory tract using an *ex vivo* model [41], in contrast to zinc supplementation which has been shown to improve lung integrity in a mouse model through decreased recruitment of neutrophils to the lungs [42].

According to the NIH, US National Library of Medicine, there are no formal studies evaluating zinc for COVID-19 management that have been completed and reported to date, although several trials are currently registered to test zinc as part of different regimens to treat COVID-19 [37] Nevertheless, zinc supplementation has been found to have a beneficial effect on COVID-19 patients [43,44]. A study of 47 COVID-19 patients showed that 57% of COVID-19 patients were zinc deficient. These zinc deficient patients developed more complications and had increased mortality than those with normal zinc levels [45].

Regarding the elderly, supplementation with 45 mg elemental zinc per day markedly reduced the incidence of infection in elderly subjects, ranging from 55 years to 87 years [10]. Consuming around 25-40 mg zinc per day is affordable, and less likely to induce human toxicity, as more than 200 to 400 mg per day of zinc consumption can induce adverse events [46].

3. ω-3 Polyunsaturated fatty acids (PUFAs)

 ω -3 PUFAs have not received the attention they deserve in the prevention and treatment of COVID-19 and associated comorbidities. ω -3 PUFAs have properties that are significantly different from those of zinc, vitamin D and magnesium, properties that are nevertheless ideally suited to prevention and treatment of COVID-19, obesity and diabetes, cardiovascular diseases, chronic pulmonary diseases and cancer, and improving immune function and anti-inflammatory effects in general ageing [1,47-49].

SARS-CoV and SARS-CoV-2 are very similar and they are both enveloped viruses that can lead to development of ARDS. ω -3 PUFAs, in particular eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), have been shown to inactivate enveloped viruses as well as to inhibit the proliferation of a range of microbial organisms [50].

The efficacy of a-linolenic acid (ALA) in treating inflammation and immune deficiency has been shown in a number of cases to be similar to that of DAH and EPA, although typically the potency is in the order DHA>EPA>ALA [eg, 49,51]. It has been reported that oil rich in ALA caused an immune modulation in cancer similar to that with fish oil, which was accompanied by a decrease in macrophage production of pro-inflammatory cytokines (eg, TNF-a and IL-6) [52].

 ω -3 PUFAs have an important role in regulating macrophages as they modulate the production of cytokines and chemokines by macrophages; they change macrophage phagocytosis capacity and they convert macrophages from proinflammatory (M1 type macrophages) to anti-inflammatory (M2) type [53]. ω -3 PUFAs and their metabolites have a modulating effect on neutrophils as they affect neutrophil migration, phagocytic capacity and the production of ROS [53].

Weill and co-workers [51] described the action of PUFAs as having two phases in inhibiting inflammation: a promotion phase in which ω -6 PUFAs such as AA lead to the synthesis of pro-inflammatory leukotrienes and prostaglandins through the action of cyclooxygenases, lipoxygenases and cytochrome P450; and a resolution phase where ω -3 PUFAs are precursors to potent active mediators such as resolvins, maresins and protectins which inhibit the synthesis of pro-inflammatory cytokines through downregulation of the NF- κ B pathway. Resolvins are sourced from EPA and DHA and protectins are sourced from DHA; they have anti-inflammatory effects by limiting leucocyte infiltration in infected tissues [51,54]. Maresins are sourced from DHA and they also resolve inflammation [51,55].

ω-3 PUFAs are notable for their influence on the properties of lipid rafts, which in turn perform an important role in the functioning of the outer leaflet of cellular membranes. ω-3 PUFAs regulate membrane properties such as membrane fluidity and signal transduction [1]. SARS-CoV has been shown to rely on lipid raft integrity for virus entry to host cells [56,57]. The entry receptor for the coronavirus, ACE2, is located in lipid rafts. The ACE2 receptor-mediated endocytosis is followed by activation of the spike protein in the viral envelope by the transmembrane serine protease 2 (TMPRSS2) which is located adjacent to the ACE2 receptor [51]. It has been shown that ω-3 PUFAs inhibit cellular entry through ACE2 and the enzymatic activity of TMPRSS2 [51,58]. The disruptive effect of ω-3 PUFAs on the integrity of lipid rafts has been described before [1], where ω-3 PUFAs were described as causing disruption to lipid rafts due to the very poor affinity of ω-3 PUFAs for cholesterol. It is therefore clear that ω-3 PUFAs have multiple inhibitory effects on viral entry into host cells.

There have been a number of clinical studies confirming the anti-inflammatory and immune response effects of ω -3 PUFA supplementation [4,5,59-61]. According to the NIH, US National Library of Medicine, there are no formal studies evaluating ω -3 PUFAs for COVID-19 management that have been completed and reported to date, although four trials are currently registered to test ω -3 PUFAs as part of different regimens [37].

4. Vitamin D

Vitamin D obtained from sunlight or dietary sources is catalysed by vitamin D-25-hydroxylase in the liver to 25-hydroxyvitamin D₃ (25(OH)D), the major circulating form of vitamin D. 25(OH)D is biologically inert until it is hydroxylated by the enzyme 1a-hydroxylase (CYP27B1) in the kidney to the active form 1a,25-dihydroxyvitamin D₃ (calcitriol, 1a,25(OH)₂D₃) [62].

Calcitriol has important immunoregulatory and anti-inflammatory effects that it exerts through interaction with the vitamin D receptor (VDR). The calcitriol/VDR complex can interact with different gene transcription factors that control

inflammatory responses [63]. VDR and CYP27B1 are expressed in many types of immune cells, including lymphocytes, monocytes/macrophages, dendritic cells, T and B cells [64,65], and on pulmonary epithelial cells. These immune cells can convert 25(OH)D into biologically active calcitriol [63,66]. The calcitriol/VDR complex causes transcription of the antimicrobial peptides cathelicidins and defensins. Cathelicidins disrupt bacterial cell membranes as well as enveloped viruses such as SARS-CoV-2, while defensins promote chemotaxis of inflammatory cells through increased capillary permeability [65,67].

Synthesis of vitamin D in the skin is controlled by the season, the time of the exposure during the day and the latitude [68,69]. Vitamin D is poorly synthesized above (to the north) and below (to the south) of 35° latitude during winter months [70]. Lockdowns, implemented to minimize the spread of COVID-19, are therefore detrimental to vitamin D synthesis as people are prevented from going out from their homes and absorbing the sunshine, which has a cumulative effect in the winter months when COVID-19 is more prevalent. The Black and Asian populations produce less vitamin D as a result of a higher skin melanin content than those with white skin [71]. Excess exposure to sunlight is the major cause of skin cancer [72]. However, there is an increased incidence of skin cancer and other cancers in countries with low levels of sunlight compared with those countries with higher levels of sunlight throughout the year [73,74], supporting the proposition that sunlight is beneficial for synthesis of vitamin D and subsequent prevention of cancers. There have been a number of reports where low sun exposure has been shown to have a negative impact on a range of health issues [75-77]. The advantage of sun exposure in providing vitamin D needs to be sensibly balanced against the potential risk of skin cancer from excessive sun exposure [78].

In the early stages of acute inflammation, vitamin D inhibits the proliferation of Th1 and Th17 cells and their abnormal release of IFN- γ , TNF- α , IL-1, IL-2, IL12, IL-23 and IL-17, IL-21 [65]. During the resolution phase of inflammation, vitamin D-mediates differentiation of Th2 cells and release of their antiinflammatory cytokines (IL-4 and IL-10), evading the organ damage that could be caused by an excessive immune response [65]. Vitamin D has powerful antiinflammatory properties that play an important role in controlling immune function in pulmonary infection; eg, it inhibits the effects of TNF- α , it inhibits NF- κ B activity in immune cells, it inhibits the activation of inflammasomes and hence release of IL-1 β , and it decreases expression of IL-6, a major contributor to the so-called 'cytokine storm' [65,79].

The immune response acts in concert with the inflammatory response. The innate immune system acts as the first line of defence against invading pathogens such as viruses. Calcitriol enhances that defence by recruiting neutrophils, monocytes/macrophages and dendritic cells which kill and clear the viral pathogens, ultimately initiating the adaptive immune response. This response can be overactive resulting in the cytokine storm. Calcitriol inhibits this chronic immune response by downregulating the toll-like receptors (TLRs) that identify the viral pathogens initially, and inhibits the TNF- α /NF- κ B and IFN- γ signalling pathways. Calcitriol shifts the T cell profile from the pro-inflammatory Th1 and Th17 forms to the anti-inflammatory Th2 and Treg forms, respectively [80]. Tregs provide a major defence against inflammation, releasing anti-inflammatory cytokines IL-10 and TGF- β . Treg levels are markedly decreased in

severe COVID-19 disease, in contrast to high levels of Treg correlating with reduced levels of viral disease [81].

Natural killer cells are innate immune cells and they are known to possess strong antiviral activity as well as anticancer activity [82]. The count and activity of natural killer cells have been shown to be reduced below normal in COVID-19 patients, and vitamin D has been found to increase the activity of natural killer cells [82].

Although there is inconsistency in the data, it is apparent that vitamin D deficiency is influential in increasing risk of acute respiratory tract infections [83], particularly when considering the decrease in natural synthesis of vitamin D in winter-time when acute respiratory infections are most prevalent. Ali [84] conducted a study of COVID-19 cases and mortality in 20 European countries, finding that vitamin D status correlated negatively with COVID-19 cases but not with mortality. The effectiveness of vitamin D sufficiency in reducing risk of acute viral respiratory tract infections and pneumonia was also shown. Similar results were reported by Kara and co-workers [85], who also discussed the link between latitude, temperature and humidity and season on viral respiratory tract infections.

Allegra and co-workers [86] reported on the deficiency and supplementation of a vitamins including vitamin D, in of particular in correlating range hypovitaminosis with risk of contracting COVID-19 and associated mortality. They reported that there were positive and indeterminate results in their analysis of multiple studies. Vitamin D levels were particularly reduced in the ageing populations of Italy, Spain, and Switzerland, which were the most susceptible populations in relation to SARS-CoV-2 infection [87]. Additionally, Annweiler and co-workers [88] analysed a range of reports with the conclusion that inverse correlations were found between 25(OH)D levels in patients and COVID-19 incidence and mortality. Other reports have also covered the influence of vitamin D on outcomes of COVID-19 patients with the typical finding that vitamin D supplementation leads to an improved outcome for these patients and that vitamin D deficiency increases the risk and susceptibility for severe COVID-19 disease and mortality [69,84,87,89-99].

In another review, Lau and co-workers [100] found that vitamin D deficiency was highly prevalent in patients with severe COVID-19, which correlated in turn with obesity, male sex, advanced age, population concentration in northern climates, coagulopathy and immune dysfunction. A further meta-analysis found that vitamin D deficiency increased risk of severe infections and mortality of the critically ill [101]. Deficiency of vitamin D has been further claimed to increase the risk of contracting osteoporosis, cancer, diabetes, multiple sclerosis, hypertension, and inflammatory and immunological diseases [102]. Although vitamin D and the benefits of supplementation in preventing cancer have been discussed previously [1], it is of note that a number of researchers have demonstrated that the risk of cancer incidence and fatality is reduced with vitamin D supplementation [eg, 103-105]. The mechanism of action of vitamin D in reducing cancer risk has also been addressed in a number of reviews [eg, 106-108].

It has been specified that a reasonable level of 25(OH)D in serum is at least 30 ng/mL (75 nmol/L) [93,109], with a preference for 40-60 ng/mL (100-150 nmol/L) to ensure good health, particularly in the elderly [69,110].

In summary, vitamin D impedes the entry and replication of SARS-CoV-2, reduces the levels of pro-inflammatory cytokines, increases the levels of antiinflammatory cytokines and increases the production of natural antimicrobial peptides [111].

5. Magnesium

Magnesium is an essential element in the optimal biological functioning of the human body. Magnesium is the second most abundant intracellular cation in the human body and it is fundamental for oxidative phosphorylation, glycolysis, DNA transcription, and protein synthesis [112]. Magnesium levels are not routinely analysed in clinical practice [113] which means there has been limited reporting of magnesium correlations for COVID-19 patients [113,114]. Despite this, there have been some reports of magnesium status being lower in severe COVID-19 cases than in less severe cases [115-117]. On the other hand, there have been a number of excellent reviews of magnesium and its essentiality in maintenance of proper immune function and control of oxidative stress and low-grade inflammation, particularly in the elderly [eg, 112,118-122].

Magnesium is essential for the activation of vitamin D [122,123]. Magnesium and vitamin D are therefore both important for immune function and cellular stability and sufficiency of both is required to counteract the detrimental effects of COVID-19 development [122].

Deficiency of magnesium is common and it has been estimated that in a given population up to 30% may have a magnesium deficiency [122]. Magnesium is principally stored in bone (>50%) with only ~1% in serum [124]. Magnesium homeostasis is maintained by absorption from the gastrointestinal tract, renal excretion and exchange from bone. Estimates of magnesium sufficiency are therefore dubious if reliant on serum analysis alone [118]. However, 0.75 mmol/L has been suggested as the serum level below which magnesium deficiency exists [125], and 0.85 mmol/L as the required level for magnesium sufficiency [118].

Magnesium has anti-inflammatory and anti-oxidative effects, as well as providing vasodilation and neuroprotection [120]. Magnesium suppresses NF- κ B, expression of IL-6 and TNF-a, and levels of C-reactive protein (CRP) [6,121,126]. Magnesium therefore regulates the cardiovascular, digestive, neurological and respiratory systems, contributing significantly to maintenance of normal human health [120]. In this context, magnesium dietary intakes correlate negatively with cardiovascular disease, kidney disease and diabetes [118,127].

6. COVID-19

SARS-CoV-2 is an insidious virus that causes the disease COVID-19. There are many similarities between SARS-CoV-2 and the earlier coronavirus SARS-CoV. In

order to present the overall essentiality of ensuring sufficiency of zinc, ω -3 PUFAs, vitamin D and magnesium, the key contributing stages of SARS-CoV-2 incursion and COVID-19 development will be discussed. These are viral entry, the involvement of the immune system and inflammation, and the subsequent cytokine storm that causes the eventual morbidity and mortality that is associated with COVID-19.

6.1 Viral entry

SARS-CoV-2 enters the host cells via the angiotensin-converting enzyme 2 (ACE2) receptor, in the same manner as SARS-CoV [128]. The spike protein of SARS-CoV-2 binds to ACE2, enabling endocytosis, which is followed by activation of the S protein in the viral envelope utilizing the transmembrane serine protease 2 (TMPRSS2), a membrane-bound enzyme located adjacent to the ACE2 receptor [51]. At the same time, the ADAM17 (disintegrin and metalloproteinase domain 17) 'sheddase' is activated by the SARS-CoV-2-ACE2 complex which in turn leads to ACE2 ectodomain shedding. Activation of the ADAM17 sheddase may also cause cleavage of TNF-a and IL-6 as well as other pro-inflammatory mediators [128]. It should be noted that the ACE2-binding affinity of the of S protein of SARS-CoV-2 is 10- to 20-fold higher than that of SARS-CoV [129], suggesting that SARS-CoV-2 is significantly more infectious than its predecessor SARS-CoV.

The balance of the renin-angiotensin system (RAS) is vital in controlling host-cell entry of viruses as well as associated comorbidities, as RAS regulates blood pressure. RAS is essentially a balance between ACE and ACE2, as illustrated in Figure 1. The ACE pathway requires conversion of angiotensin (Ang) I to Ang II and subsequent binding to the AT1 receptor (AT1R), which has dire consequences such as vasoconstriction, proliferation, inflammation, and apoptosis [91]. The alternative route involves conversion of Ang I and Ang II to angiotensin 1-9 and angiotensin 1-7, respectively, through the enzymatic action of ACE2. Angiotensin 1-9 is also converted to angiotensin 1-7 by ACE.

ACE2 is required for viral entry to host cells, but it is also desirable for conversion of Ang I and Ang II to angiotensins 1-9 and 1-7, respectively, leading to activation of the Mas receptor. Figure 1 shows that this in turn will cause positive pathology in terms of vasodilation, and anti-inflammatory, anti-oxidative and anti-fibrosis effects [130].



Figure 1. RAS pathways showing the balance between ACE and ACE2.

There has been fairly extensive discussion about the contradictory actions of ACE2 in viral entry to host cells and its regulation of the RAS where modulation of RAS has a positive pathological effect. The ACE2 receptor is expressed in lung tissues, and a range of other tissues such as nose, heart, endothelium, kidney and intestine [131,132]. It has now been resolved that once the virus binds to ACE2, it effectively removes it from further action, promoting ACE activity which in turn leads to production of more Ang II. The removal of ACE2 from action therefore causes the virus to have a free run allowing it to proliferate, leading to enhanced morbidity.

The expression of ACE2 has been found to be lower in males than in females and lower in older adults than in young people, which could explain the higher incidence of deaths in elderly males with COVID-19 [130,133]. This category of patients has a worse prognosis when they are also implicated with comorbidities such as cardiovascular diseases, diabetes, hypertension, and obesity, all of which are stimulated by RAS [130]. It is therefore important to enhance the expression of ACE2 and its activity, and at the same time ensure that the entry of the virus to host cells is inhibited. This can be achieved by ensuring that sufficient levels of zinc, ω -3 PUFAs, vitamin D and magnesium are maintained at all times during prevention and treatment of COVID-19.

It should be noted that although there are no reported studies of the effect of zinc on ACE2 for host-cell entry, zinc does protect the human body from virus entry by improved mucociliary clearance of viruses as well as preserving tissue barriers [42]. It has been recognized that enhanced expression of ACE2 by calcitriol alleviates acute lung injury induced by SARS-CoV-2 [134-136]. Calcitriol also supresses renin activity and therefore reduces the generation of angiotensin II which causes pulmonary vasoconstriction [134]. As stated above, binding of ACE2 and cellular entry are inhibited by ω -3 PUFAs [51,58].

6.2 The immune system

The immune system provides two lines of defence: innate and adaptive immunity. Innate immunity is the first line of defence, reliant on mucosal barriers, monocytes, macrophages, neutrophils, eosinophils, and dendritic cells. Adaptive immunity is the process by which immunological memory to a specific

antigen is created but more slowly than innate immunity. Dendritic cells also function as antigen-presenting cells, activating B and T lymphocytes of the adaptive immune response [53].

Mast cells are present in the submucosa of the nasal cavity and respiratory tract where they provide a protective barrier against microorganisms and they can be virus-activated [29]. When activated, mast cells initially release preformed inflammatory molecules such as histamine and proteases, while late activation activates the synthesis and release of pro-inflammatory IL-1 family members, including IL-1, IL-6, and IL-33 [137]. Mast cells therefore normally release a wide range of pro-inflammatory mediators. Vitamin D diverts the release characteristics of mast cells to produce and excrete IL-10 without inducing degranulation of pro-inflammatory mediators [138]. IL-10 is an important antiinflammatory cytokine that inhibits the production of pro-inflammatory cytokines, such as IFN- γ , IL-2, IL-3, TNF-a, and GM-CSF [139].

Macrophages are fundamental to the innate immune system as they scavenge invading pathogens: they recognize invading pathogens through use of pathogen-associated molecular patterns which are in turn recognized by TLRs present on their surface. Macrophages then phagocytose the invading pathogen and at the same time secrete ROS and a large range of cytokines and chemokines to recruit and activate other types of immune cells from both the innate and the adaptive immune systems [53]. The most damaging cytokines released by macrophages when over-activated are IL-1 β , IL-6 and TNF-a [140].

Eosinophils release pro-inflammatory mediators, including degranulated cationic proteins, synthesized eicosanoids, and cytokines [141]. Neutrophils are recruited to the initial site of inflammation where they also have a role in removing pathogens. Neutrophils can also interact with the adaptive immune system by promoting naïve T cells to transition into T helper 1 (Th1) cells [53].

T cells are thymus-derived lymphocytes. T cells can be classified into helper (Th) cells that regulate the function of other immune cells, and cytotoxic T cells that destroy virus-infected cells. Th cells differentiate into Th1, Th2, Th17 and Th22 cells. Th1 cells secrete IFN- γ ; Th2 cells secrete IL-4; Th17 cells secrete IL-17A, IL17-F, IL-21, and IL-22; and Th22 cells secrete IL-22. Th1 and Th17 cells are pro-inflammatory, whereas Th2 cells are essentially anti-inflammatory [53]. Regulatory T cells (Tregs) suppress the activation of other immune cells such as Th1 cells, Th17 cells, B cells, macrophages or dendritic cells, through secretion of IL-10 and TGF- β [53].

The impact provided by the various contributing immune system cells on the SARS-CoV and SARS-CoV-2 viruses is given in Table 1. It can be seen that the release of cytokines and chemokines is potentially huge, leading to the potential for production of the cytokine storm.

Cell Type	Mediators released/activated	References
Mast cells	Histamine, tryptase, NF-κB, IL-1α/β, IL-2, IL-3, IL-4, IL-5, IL-6, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-15, IL-16,	29,142-146
	IL-17, IL-18, IL-25, IL-33, TNF-a, IFN-γ, TGF-β, CCL2, CCL3, GM-CSF, VEGF,	

Table 1. Mediators released/activated in cells in SARS and COVID-19

	PDGF, SCF, PGE2, ROS, TLR2, c-Kit	
Monocytes/macrophages	NF-κB, TNFa, IL-1a/β, IL-1RA, IL-6,	6,110,147-150
	IL-8, IL-10, IL-12, IFN-γ, TGF-β, ROS,	
	TLR2, TLR4	
Eosinophils	IL-1a, IL-2, IL-3, IL-4, IL-5, IL-6, IL-8,	145,151-154
	IL-10, IL-11, IL-12, IL-13, IL-16, IL-18,	
	IL-25, TNF-a, IFN-y, TGF-a/B, VEGF,	
	GM-CSF	
Neutrophils	IL-1a/β, IL-1RA, IL-3, IL-4, IL-6, IL-7,	155-158
	IL-8, IL-9, IL-10, IL-12, IL-16, IL-17,	
	IL-18, IL-23, Th1/Th2, TGF-a/β, IFN-a,	
	IFN-v, TNF-a, G-CSG, GM-CSF, SCF,	
	FGF, VEGF, CCL2, ROS, TLR2, TLR4	
Dendritic cells	IL-6, IL-10, IL-12, TNF-a, CCL3,	110,142,159
	RANTES, IP-10, CCL2	
Th1	IFN-γ, IL-1β, IL-2, IL-12, TNF-a	110,148
Th2	TGF-β, IL-4, IL-5, IL-9, IL-10, IL-13	110,148
Th17	IL-17A, IL-17F, IL-21, IL-22	110,160
Treg	IL-10, TGF-β	53,110

IL-10, TGF-β

Table 2 shows the key immune cells and regulators that contribute to immune function and inflammation together with the correcting effects of zinc, ω -3 PUFAs, vitamin D and magnesium on each of these immune cells and regulators when immune function and inflammation are over-activated.

Cell/regulator	Zinc	ω-3 PUFAs	Vitamin D	Magnesium
Mast cells	↓	\checkmark	\checkmark	↓
	[161,162]	[144,163,164]	[138,165-167]	[168,169]
Monocytes	\checkmark	$ $ \downarrow	\downarrow	\downarrow
	[170,171]	[172,173]	[147]	[174,175]
Macrophages	\downarrow	$ $ \downarrow	\downarrow	$ $ \downarrow
	[176]	[177,178]	[179,180]	[175,181,182]
Neutrophils	\checkmark	\checkmark	↓	\checkmark
•	[171,183,184]	[185,186]	[187,188]	[181,189]
Dendritic Cells	\checkmark	\checkmark	\checkmark	\checkmark
	[190,191]	[192,193]	[63,194,195]	[196]
Eosinophils	\checkmark	\checkmark	\bullet	\checkmark
•	[171,197,198]	[172,185,199]	[200,201]	[202]
Th1/Th2 ratio	\downarrow	↓ ↓	\checkmark	\downarrow
	[203,204]	[205,206]	[207]	[208]
Th17	\checkmark	↓ ↓	\checkmark	_
	[24,209-211]	[212-214]	[215,216]	
Treg	↑	\uparrow	↑	—
-	[24,211,217,218]	[213,219-221]	[222-224]	
Inflammasome/	\checkmark	\downarrow	\downarrow	\downarrow
caspase-1	[225,226]	[227-229]	[79,230-232]	[233,234]
NF-κB	↓	\checkmark	\checkmark	\checkmark
	[12,17,25,235]	[132,236,237]	[79,238-240]	[121,174,241]

	Table 2. Inhibition(ψ)/activation(个)	of immune	cells	cells/regulators
--	----------------------	--------	----------------	---	-----------	-------	------------------

- indicates no literature reference found

6.3 Cytokine storm

The uncontrolled release of immune cells and excessive release of proinflammatory cytokines has been termed the 'cytokine storm'. The cytokine storm normally presents as systemic inflammation, excessive oxidative stress and multiple organ failure [29,51], predominantly resulting in ARDS. The key to counteracting the cytokine storm lies in counteracting excessive inflammation. This can be largely addressed through maintenance of sufficiency of the essential nutrients zinc, ω -3 PUFAs, vitamin D and magnesium. Table 3 gives the key proinflammatory cytokines and other mediators involved in a cytokine storm, together with the inhibitory effects of zinc, ω -3 PUFAs, vitamin D and magnesium. It can be seen that these four nutrients are widely effective in inhibiting the key pro-inflammatory mediators of the cytokine storm.

Mechanisms	Effect of Zinc on mediator [Refs]	Effect of ω-3 PUFAs on mediator [Refs]	Effect of Vitamin D on mediator [Refs]	Effect of Magnesium on mediator [Refs]
TNF-a	↓		↓ •	V
	[9,235]	[1//,242-244]	[245-248]	[126,174,249]
IFN-γ	₩ [217,218,250]	₩ [193,212,244]	₩ [247,248,251-253]	₩ [208,249]
IL-1B	\mathbf{V}	\mathbf{V}	↓ · · ·	Ý I
F	[9,210,254]	[177,227,244,255, 256]	[79,231,232,248, 257]	[182,233,249,258, 259]
IL-6	\checkmark	\mathbf{h}	↓ ↓	↓ ↓
	[17,260]	[177,212,243,244, 256]	[65,248,261,262]	[121,126,174,182, 263]
IL-12	\checkmark	\mathbf{A}	$\mathbf{+}$	_
	[264]	[172,265]	[111,195,266,267]	
IL-17	\mathbf{V}	\checkmark	4	—
	[209,210,268]	[185,193,212,244]	[216,248,269,270]	
IL-18	\checkmark	\checkmark	\checkmark	—
	[271]	[228]	[272]	
IL-33	\checkmark	$\mathbf{+}$	\checkmark	—
	[273]	[274]	[275]	•
CCL2	V	$\mathbf{\Psi}$	•	\mathbf{V}
(MCP-1)	[9,171]	[276-279]	[280-282]	[263,283]
CCL3	\downarrow	\mathbf{V}	\downarrow	—
(MIP-1a)	[284]	[276,285]	[216,286]	
C-reactive	\checkmark	$\mathbf{+}$	\checkmark	\checkmark
protein (CRP)	[287-289]	[290-292]	[93,293,294]	[121,126,295]
GM-CSF	\mathbf{V}	\checkmark	\checkmark	—
	[296]	[297,298]	[299-302]	
NF-κB	\checkmark	\mathbf{V}	\mathbf{V}	\mathbf{V}
	[12,25,226,235]	[132,236,237]	[79,238-240]	[121,174,241]

Table 3. Key pro-inflammatory mediators in a cytokine storm. Ψ inhibits the mediator.

- indicates no literature reference found

7. Ageing, obesity and non-communicable diseases

Deficiencies in zinc, ω -3 PUFAs, vitamin D and magnesium have been shown above to provide significant risk factors for severe COVID-19 disease as well as for pre-existing conditions such as ageing, obesity/diabetes, cardiovascular diseases, chronic respiratory diseases and cancer. All of these comorbidities are accompanied by systemic inflammation which likely impacts on the COVID-19 outcome [29].

Table 4 lists the immune cells and mediators that are released in COVID-19, the cytokine storm, ageing, obesity/diabetes and the principal non-communicable diseases. It can be seen that many of the mediators, particularly those that are key pro-inflammatory cytokines such as IL-1 β , IL-6, TNF-a and IFN- γ , are common to COVID-19, the cytokine storm (which in turn is part of COVID-19) and the listed comorbidities. The entry in Table 4 for chronic respiratory diseases has been taken to be the same as the cytokine storm, which is the principal force behind creation of ARDS.

Table 4 Key cells and mediators associated with COVID-19, cytokine storm, ageing and comorbidities

Cells/Mediators/Transcription factors	References
Mast cells, neutrophils, eosinophils,	29,133,142,148,160,303
monocytes, macrophages, dendritic cells,	
NF-κB, IL-1β, IL-1RA, IL-2, IL-6, IL-7,	
IL-8, IL-9, IL-10, IL-12, IL-13, IL-17,	
IL-18, IL-21, IL-22, IL-33, TNF-a, IFN-γ,	
GM-CSF, G-CSF, CCL2, CCL3, IP-10,	
Th1/Th2, PDGF, VEGF, FGF, CRP	
IL-1β, IL-6, IL-7, IL-8, IL-9, IL-12, IL-17,	29,142,155,160,304
IL-18, IL-33, TNF-α, IFN-γ, CCL2, CCL3,	
FGF, G-CSF, GM-CSF, IP-10, PDGF, VEGF,	
CRP	
IL-1, IL-1RN, IL-2, IL-6, IL-8, IL-12,	305,306
IL-13, IL-18, CRP, IFN- γ , IGF- β , INF- a ,	X
	205 201
$M2 \rightarrow M1$, $In2 \rightarrow In1$, $Ireg \rightarrow In1/$, B cells,	305,301
IL-I β , IL-6, IL-7, IL-22, IFN- γ , INF- β ,	
	202.200
NF-KB, IL-IB, IL-2, IL-4, IL-6, IL-17,	302,309
GM-CSF, MMP-2, MMP-9, CCL2, ERK1/2,	
P38 MAPK, INF-U, IFN-Y, HIF-IU, ILKZ, $T \Box A$	
	20 142 155 160 204
IL-ID, IL-0, IL-7, IL-0, IL-9, IL-IZ, IL-I7,	29,142,155,160,304
ECE GLOSE GMLOSE ID-10 PDGE VEGE	
$NE_{\mu}R_{\mu}PE2 COV 2/PCE2 TNEq TI 10$	1
II_{-6} II_{-8} P_{-7}	T
HER2 VEGE Cyclin D1 c-Myc PTEN	
MDM2 HIPK2 A20 $n21$ TGF-B PARP	
caspases-3.7.8.9. Bcl-2. Bcl-xL. Bax	
cytochrome c. ROS, iNOS, MMP9 HIF-1a	
TLR4.	
	Cells/Mediators/Transcription factors Mast cells, neutrophils, eosinophils, monocytes, macrophages, dendritic cells, NF-κB, IL-1β, IL-1RA, IL-2, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12, IL-13, IL-17, IL-18, IL-21, IL-22, IL-33, TNF-a, IFN-γ, GM-CSF, G-CSF, CCL2, CCL3, IP-10, Th1/Th2, PDGF, VEGF, FGF, CRP IL-1β, IL-6, IL-7, IL-8, IL-9, IL-12, IL-17, IL-18, IL-33, TNF-a, IFN-γ, CCL2, CCL3, FGF, G-CSF, GM-CSF, IP-10, PDGF, VEGF, CRP IL-1, IL-1RN, IL-2, IL-6, IL-8, IL-12, IL-13, IL-18, CRP, IFN-γ, TGF-β, TNF-a, SAA M2→M1, Th2→Th1, Treg→Th17, B cells, IL-1β, IL-6, IL-7, IL-22, IFN-γ, TNF-β, CCL2, TNF-a NF-κB, IL-1β, IL-2, IL-4, IL-6, IL-17, GM-CSF, MMP-2, MMP-9, CCL2, ERK1/2, P38 MAPK, TNF-a, IFN-γ, HIF-1a, TLR2, TLR4 IL-1β, IL-6, IL-7, IL-8, IL-9, IL-12, IL-17, IL-18, IL-33, TNF-a, IFN-γ, CCL2, CCL3, FGF, G-CSF, GM-CSF, IP-10, PDGF, VEGF NF-κB, p53, COX-2/PGE2, TNFa, IL-1β, IL-6, IL-8, p27, PPARa,γ, GSK-3, EGFR, HER2, VEGF, Cyclin D1, c-Myc, PTEN, MDM2, HIPK2, A20, p21, TGF-β, PARP, caspases-3,7,8,9, Bcl-2, Bcl-xL, Bax, cytochrome c, ROS, iNOS, MMP9, HIF-1a, TLR4,

7.1 Ageing

As the human body ages, there is a gradual decline in functioning of the innate and adaptive immune systems, designated immunosenescence, as well as an increase in the levels of pro-inflammatory cytokines IL-1 β , IL-2, IL-6, IL-8, TNF-a and IFN- γ , as well as CRP [148,305,310]. There is also a decline in ACE2 expression, similarly to COVID-19 [148]. Ageing also produces excess ROS production which can initiate pro-inflammatory generation through activation of transcription factors such as NF- κ B [148]. T-cell function has been found to become increasingly defective in the elderly, decreasing immune function [20].

Maintenance of healthy functioning of cells is of increasing importance as ageing progresses. Working against this, deficiencies in one or more of zinc, ω -3 PUFAs, vitamin D and magnesium will lead inevitably to a diminution in immune function and an increase in levels of inflammatory mediators [21,27,311]. Deficiencies of zinc, ω -3 PUFAs, vitamin D and magnesium increase with ageing, frequently contributing to age-related diseases such as diabetes, cardiovascular diseases and chronic pulmonary diseases [27].

Diet is very important for adequate intake of zinc, ω -3 PUFAs and magnesium as the importance and interest in quality of food diminishes with old-age, as well as

the degree of absorption [312] In addition, the exposure of the aged to sunlight becomes severely limited, leading to diminished vitamin D levels.

7.2 Obesity/diabetes

Obesity is related to the accumulation of pro-inflammatory cells in visceral adipose tissue, which can lead to insulin resistance and to diabetes mellitus [151]. Obesity is associated with low-grade inflammation, which in turn is associated with diminution of the innate and adaptive immune responses. Low-grade inflammation is linked to adipocyte hypoxia and dysfunction [307]. There is a significant release of pro-inflammatory cytokines (eg, IL-1 β , IL-6, TNF-a) that activate in turn macrophages, T cells and B cells, creating an auto-regenerating loop [307,313]. Obesity is also associated with increased oxidative stress [314].

Zinc deficiency has been shown in a number of studies to be associated with obesity and diabetes [315-317]. Zinc is essential for the normal physiological processing of insulin and is therefore directly associated with diabetes [318].

The ω -6/ ω -3 PUFA ratio has increase dramatically over the past 50 years and it has contributed to the increased proportion of the population who are obese [319]. It has been shown that this trend can be reversed by increasing the EPA and DHA intakes [319]. It has been recommended that fish oil emulsion supplement be administered to those who are obese and at risk of contracting COVID-19, due to the immune modulatory properties of EPA and DHA [320].

Obesity increases the risk of vitamin D deficiency, mainly due to the higher adiposity of the obese individual. Vitamin D is fat-soluble and is predominantly stored in the adipose tissues, leading to low levels of vitamin D in the circulation [321]. Low vitamin D levels have been reported consistently across age groups, ethnicity and geography [322,323]. Meta-analyses found that vitamin D with increased [321,324]. deficiency correlated obesity Vitamin D supplementation has been shown to reduce insulin resistance [325] and diabetes mellitus has been found to correlate with vitamin D deficiency in older adults [326].

There is a positive relationship between magnesium deficiency and obesity and chronic inflammation [327]. In turn, obesity is a major risk factor for chronic diseases which depend on chronic inflammation, such as diabetes, cardiovascular diseases and cancer [327].

7.3 Cardiovascular diseases

A large proportion of COVID-19 patients have risk factors associated with cardiovascular diseases [328]. The high levels of inflammation associated with COVID-19 can induce cardiovascular diseases [80,328]. Studies of COVID-19 individuals with underlying cardiovascular disease were at increased risk of severe disease and mortality [329].

Choi and co-workers [330] reviewed the literature covering zinc status and cardiovascular diseases. They found that zinc deficiency was associated with atherosclerosis, hypertension, myocardial infarction, atrial fibrillation and

congestive heart failure. Similarly, Jurowski and co-workers [331] had reviewed the literature, reporting that zinc deficiency correlated with hypertension, atherosclerosis and heart failure. Further reports support the fact that zinc deficiency is associated with cardiovascular diseases [22,23].

The cardioprotective effects of n-3 PUFAs and their metabolites are ascribed mainly to their immunomodulatory properties. Emerging evidence demonstrates the ability of ω -3 PUFAs to reduce circulating levels of inflammatory chemokines, cytokines, and the pro-inflammatory metabolites derived from ω -6 PUFAs [332,333]. A number of studies have found that higher consumption of ω -3 PUFAs lowers the number of deaths related to cardiovascular disease [334-337]. Darwesh and co-workers [338] presented a detailed report on the positive effects of ω -3 PUFAs in cardiovascular diseases, which included stabilization of atherosclerotic plaques, reducing the incidence of thrombus formation, enriching cellular membranes and altering the structure of lipid rafts and their function to benefit the treatment of cardiovascular diseases.

There is a strong correlation between obesity and vitamin D deficiency, as well as between obesity and cardiovascular disease. It would therefore be anticipated that there would be a benefit in supplemental vitamin D for obese patients at risk of cardiovascular disease [339]. A study of 137 elderly Brazilian patients found that 65% were vitamin D deficient and there was a strong association between vitamin D deficiency and the risk of heart failure [340].] A number of literature reviews have examined the association between vitamin D deficiency and the incidence of cardiovascular diseases, with the conclusion that vitamin D decreases inflammation and pro-inflammatory cytokines causing a strong association with cardiovascular diseases [308,341,342].

The anti-inflammatory and anti-oxidative effects of magnesium provide cardiovascular protection [119,120]. Qu and co-workers [127] provided a metaanalysis that showed an inverse correlation between magnesium serum concentrations and the risk of total cardiovascular events.

7.4 Lung diseases

Lung diseases include pneumonia, bronchitis and asthma. The most common lung disease associated with COVID-19 is acute respiratory distress syndrome (ARDS), promoted most often by the cytokine storm, and which is often lethal [51]. ARDS occurs in approximately 10% of COVID-19 patients [51].

Meydani and co-workers [20] found that nursing home elderly who were zinc deficient were more likely to contract pneumonia with its subsequent consequences. Further reports support the fact that zinc deficiency is associated with chronic pulmonary diseases [21,23]. Skalny and co-workers [12] deduced that zinc has the propensity to alleviate COVID-19 through its properties of reducing inflammation, improving mucociliary clearance and promoting antiviral and antibacterial immunity.

Weill and co-workers [51] discussed the properties of ω -3 PUFAs, which include interference of viral entry and replication and inhibition of inflammation, leading to improvement of the outcome of critically ill patients with ARDS. It was shown in a study where bronchoalveolar lavage fluid was added to A549 cells that by

increasing the ratio of ω -3: ω -6 PUFAs, there was a decrease in levels of NF- κ B, COX-2 and PGE2, and an increase in release of IL-10 and PPAR γ [343].

It has been noted that there is a strong link between seasonality of low vitamin D levels and occurrence and prevalence of influenza in winter [80]. It has also been reported that a high percentage (>80%) of chronic obstructive pulmonary disease patients had low vitamin D levels [344]. The association between higher vitamin D levels and improved lung function has also been reported [345-347]. Moreover, it has been reported that vitamin D deficiency is associated with occurrence of respiratory disease and ensuing mortality [90,347-349].

The role of magnesium in lung function was discussed by de Baaij and coworkers [124], where magnesium was described as having three roles: a strong vasodilator and bronchodilator effect, regulation of the release of acetylcholine and histamine, and as an anti-inflammatory agent. Magnesium was therefore suggested as a useful treatment for asthma and chronic obstructive pulmonary disorder. Micke and co-workers [114] also discussed magnesium and lung function in some detail, with a similar analysis of the anticholinergic, antihistaminic and anti-inflammatory effects of magnesium.

7.5 Cancer

Cancer has been discussed in the context of essentiality of sufficiency of zinc, ω -3 PUFAs and vitamin D [1]. The opportunity of including magnesium as a further essential component in prevention and treatment of cancers is taken here, as magnesium is essential for the activation of vitamin D [122,123]. Magnesium, as discussed above, is also active in regulating the immune system and controlling untoward oxidative stress and inflammation [119,120] which are prevalent in the early development of cancers [350].

8. Discussion

COVID-19 and its virus SARS-CoV-2 have provided an ideal opportunity to reset the approach to prevention and treatment of non-communicable diseases, particularly those that occur predominantly in the aged. COVID-19 has been shown to be linked to comorbidities such as senescence occurring in the aged, obesity/diabetes which are more severe in the aged, and cardiovascular diseases and chronic pulmonary diseases which are more prevalent in the aged, as well as cancers. It is therefore opportune to examine carefully the prevention and treatment of COVID-19 and those diseases, with particular attention to those features and characteristics that are common to these diseases. The most outstanding common features are inflammation and overactivity of the innate and adaptive immune systems. Control of inflammation and the immune system are fundamentally dependent on sufficiency of the essential nutrients zinc, ω -3 PUFAs, vitamin D and magnesium.

This paper has been directed towards an appreciation of the benefits of having sufficiency of zinc, ω -3 PUFAs, vitamin D and magnesium. These four components are essential as they are natural to the normal functioning of cells and multiple other components of the human body. They are extremely safe when supplemented in a controlled manner. Control in the aged (eg, 65 years and older) can be maintained by annual analyses of their serum levels. This can

be achieved with government support, as well as by government supplies of supplements where necessary. The cost of this service to those over 65 would be small compared with the potential savings in hospitalization and critical care costs. As an example, a German estimate of the effect of supplementing vitamin D alone on the cost savings of cancer alone in Germany showed a cost saving of approximately &254 million per year with a prevention of almost 30,000 deaths to cancer per year [351].

Zinc, ω -3 PUFAs, vitamin D and magnesium are pleiotropic as they allow, and in fact boost, the functioning of granulocytes such as mast cells, neutrophils and eosinophils, as well as monocytes/macrophages, dendritic cells, T cells and B cells in normal conditions and when there are minor invasions of pathogens such as minor viral and bacterial infections. In contrast, zinc, ω -3 PUFAs, vitamin D and magnesium all act to suppress hyperinflammation and major disruptions of the immune system that occur when there is a significant invasion by viral or bacterial pathogens such as SARS-CoV-2 or non-communicable diseases such as diabetes, cardiovascular disease or chronic pulmonary disease. In these situations, zinc, ω -3 PUFAs, vitamin D and magnesium have the ability to suppress excessive inflammation and dysregulation of the immune system. These nutrients are therefore essential in all aspects; when present in sufficiency they are directed towards ensuring good health for humans at all times and for all ages. This is not normally the case with non-natural drugs that are prescribed for treatment of particular pathological conditions.

Vaccines are rarely 100% in their prevention of transmission and their prevention of humans contracting the particular disease; there are potential problems with mutations and diminution of their effectiveness. It is of note that vaccines perform their function through the adaptive immune system, whilst zinc, ω -3 PUFAs, vitamin D and magnesium affect both the innate and adaptive immune systems. Supplementation of the four nutrients in treatment of COVID-19 is therefore desirable, especially if this supplementation is beneficial in preventing or treating non-communicable diseases or reducing the adverse effects of ageing.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of competing interest

There is no conflict of interest.

References

- 1. Story MJ (2021). Zinc, ω -3 polyunsaturated fatty acids and vitamin D: An essential combination for prevention and treatment of cancers. Biochimie 181:100-1222.
- 2. "Global Status Report on Non-Communicable Diseases," WHO. 2010. https://www.who.int/nmh/publications/ncd report full en.pdf
- 3. Pecora F, Persico F, Argentiero A, Neglia C, Esposito S (2020). The role of micronutrients in support of the immune response against viral infections. Nutrients 12(10):E3198
- 4. Szabó Z, Marosvölgyi T, Szabó É, Bai P, Figler M, Verzár Z (2020). The potential beneficial effect of EPA and DHA supplementation managing cytokine storm in coronavirus disease. Front Physiol 11:752
- 5. Calder PC, Carr AC, Gombart AF, Eggersdorfer M (2020). Optimal nutritional status for a well-functioning immune system is an important factor to protect against viral infections. Nutrients 12(4):1181
- 6. Iddir M, Brito A, Dingeo G, Fernandez Del Campo SS, Samouda H, La Frano MR, et al (2020). Strengthening the immune system and reducing inflammation and oxidative stress through diet and nutrition: Considerations during the COVID-19 crisis. Nutrients 12(6):1562
- 7. Mammadova-Bach E, Braun A (2019). Zinc homeostasis in platelet-related diseases. Int J Mol Sci 20(21):5258
- 8. Prasad AS (2008). Zinc in human health: Effect of zinc on immune cells. Mol Med 14(5-6):353-7
- 9. Bao B, Prasad AS, Beck FW, Fitzgerald JT, Snell D, Bao GW, et al (2010). Zinc decreases C-reactive protein, lipid peroxidation, and inflammatory cytokines in elderly subjects: a potential implication of zinc as an atheroprotective agent. Am J Clin Nutr 91(6):1634-41
- Prasad AS, Beck FW, Bao B, Fitzgerald JT, Snell DC, Steinberg JD, et al (2007). Zinc supplementation decreases incidence of infections in the elderly: effect of zinc on generation of cytokines and oxidative stress. Am J Clin Nutr 85(3):837-44
- 11. Gao H, Dai W, Zhao L, Min J, Wang F (2018). The role of zinc and zinc homeostasis in macrophage function. J Immunol Res 2018:6872621
- Skalny AV, Rink L, Ajsuvakova OP, Aschner M, Gritsenko VA, Alekseenko SI, et al (2020). Zinc and respiratory tract infections: Perspectives for COVID-19 (Review). Int J Mol Med 46(1):17-26
- 13. Kloubert V, Rink L (2015). Zinc as a micronutrient and its preventive role of oxidative damage in cells. Food Funct 6(10):3195–3204
- 14. Prasad AS (2013). Discovery of human zinc deficiency: its impact on human health and disease. Adv Nutr 4(2):176-90
- 15. Shankar AH, Prasad AS (1998). Zinc and immune function: The biological basis of altered resistance to infection. Am J Clin Nutr 68(2 Suppl):447S-463S
- 16. Haase H, Rink L (2009). The immune system and the impact of zinc during aging. Immun Ageing 6:9
- 17. Mayor-Ibarguren A, Busca-Arenzana C, Robles-Marhuenda Á (2020). A hypothesis for the possible role of zinc in the immunological pathways related to COVID-19 infection. Front Immunol 11:1736
- 18. Haase H, Rink L (2014). Zinc signals and immune function. Biofactors 40(1):27-40

- 19. Gombart AF, Pierre A, Maggini S (2020). A review of micronutrients and the immune system-working in harmony to reduce the risk of infection. Nutrients 12(1):236
- 20. Meydani SN, Barnett JB, Dallal GE, Fine BC, Jacques PF, Leka LS, et al (2007). Serum zinc and pneumonia in nursing home elderly. Am J Clin Nutr 86(4):1167-73
- 21. Barnett JB, Hamer DH, Meydani SN (2010). Low zinc status: a new risk factor for pneumonia in the elderly? Nutr Rev 68(1):30–37
- 22. Braun LA, Ou R, Kure C, Trang A, Rosenfeldt F (2018). Prevalence of zinc deficiency in cardiac surgery patients. Heart Lung Circ 27(6):760–762
- 23. Derwand R, Scholz M (2020). Does zinc supplementation enhance the clinical efficacy of chloroquine/hydroxychloroquine to win today's battle against COVID-19? Med Hypotheses 142:109815
- 24. George MM, Subramanian Vignesh K, Landero Figueroa JA, Caruso JA, Deepe GS Jr (2016). Zinc induces dendritic cell tolerogenic phenotype and skews regulatory T cell-Th17 balance. J Immunol 197(5):1864-76
- 25. Prasad AS, Bao B, Beck FW, Sarkar FH (2011). Zinc-suppressed inflammatory cytokines by induction of A20-mediated inhibition of nuclear factor-κB. Nutrition 27(7-8):816-23
- 26. McGonagle D, Sharif K, O'Regan A, Bridgewood C (2020). The role of cytokines including interleukin-6 in COVID-19 induced pneumonia and macrophage activation syndrome-like disease. Autoimmun Rev 19(6):102537
- 27. Alexander J, Tinkov A, Strand TA, Alehagen U, Skalny A, Aaseth J (2020). Early nutritional interventions with zinc, selenium and vitamin D for raising anti-viral resistance against progressive COVID-19. Nutrients 12(8):2358
- Arentz S, Yang G, Goldenberg J, Beardsley J, Myers SP, Mertz D, et al (2020). Zinc for the prevention and treatment of SARS-CoV-2 and other acute viral respiratory infections: a rapid review. Adv Integr Med 7(4):252-260
- 29. Zabetakis I, Lordan R, Norton C, Tsoupras A (2020). COVID-19: The inflammation link and the role of nutrition in potential mitigation. Nutrients 12(5):1466
- 30. Stefanidou M, Maravelias C, Dona A, Spiliopoulou C (2006). Zinc: a multipurpose trace element. Arch Toxicol 80(1):1-9
- 31. te Velthuis AJ, van den Worm SH, Sims AC, Baric RS, Snijder EJ, van Hemert MJ (2010). Zn(2+) inhibits coronavirus and arterivirus RNA polymerase activity in vitro and zinc ionophores block the replication of these viruses in cell culture. PLoS Pathog 6(11):e1001176
- 32. Rajaiah R, Abhilasha KV, Shekar MA, Vogel SN, Vishwanath BS (2020). Evaluation of mechanisms of action of re-purposed drugs for treatment of COVID-19. Cell Immunol 358:104240
- 33. Doboszewska U, Wlaź P, Nowak G, Młyniec K (2020). Targeting zinc metalloenzymes in coronavirus disease 2019. Br J Pharmacol 177(21):4887-4898
- 34. Dabbagh-Bazarbachi H, Clergeaud G, Quesada IM, Ortiz M, O'Sullivan CK, Fernández-Larrea JB (2014). Zinc ionophore activity of quercetin and epigallocatechin-gallate: from Hepa 1-6 cells to a liposome model. J Agric Food Chem 62(32):8085-93
- 35. Brooks HA, Gardner D, Poyser JP, King TJ (1984). The structure and absolute stereochemistry of zincophorin (antibiotic M144255): a monobasic

carboxylic acid ionophore having a remarkable specificity for divalent cations. J Antibiot (Tokyo) 37(11):1501-4

- 36. Rizzo E (2020). Ivermectin, antiviral properties and COVID-19: a possible new mechanism of action. Naunyn Schmiedebergs Arch Pharmacol 393(7):1153-1156
- 37. ClinicalTrials.gov. https://clinicaltrials.gov/ct2/home
- 38. Pal A, Squitti R, Picozza M, Pawar A, Rongioletti M, Dutta AK, et al (2021). Zinc and COVID-19: Basis of current clinical trials. Biol Trace Elem Res Ahead of print
- 39. Kumar A, Kubota Y, Chernov M, Kasuya H (2020). Potential role of zinc supplementation in prophylaxis and treatment of COVID-19. Med Hypotheses 144:109848
- 40. Wessels I, Rolles B, Rink L (2020). The potential impact of zinc supplementation on COVID-19 pathogenesis. Front Immunol 11:1712
- 41. Roscioli E, Jersmann HP, Lester S, Badiei A, Fon A, Zalewski P, et al (2017). Zinc deficiency as a codeterminant for airway epithelial barrier dysfunction in an ex vivo model of COPD. Int J Chron Obstruct Pulmon Dis 12:3503-3510
- 42. Wessels I, Pupke JT, von Trotha KT, Gombert A, Himmelsbach A, Fischer HJ, et al (2020). Zinc supplementation ameliorates lung injury by reducing neutrophil recruitment and activity. Thorax 75(3):253-261
- 43. Finzi E (2020). Treatment of SARS-CoV-2 with high dose oral zinc salts: A report on four patients. Int J Infect Dis 99:307-309
- 44. Carlucci PM, Ahuja T, Petrilli C, Rajagopalan H, Jones S, Rahimian J (2020). Zinc sulfate in combination with a zinc ionophore may improve outcomes in hospitalized COVID-19 patients. J Med Microbiol 69(10):1228-1234
- 45. Jothimani D, Kailasam E, Danielraj S, Nallathambi B, Ramachandran H, Sekar P, et al (2020). COVID-19: Poor outcomes in patients with zinc deficiency. Int J Infect Dis 100:343-349
- 46. Razzaque MS (2020). COVID-19 pandemic: Can maintaining optimal zinc balance enhance host resistance? Tohoku J Exp Med 251(3):175-181
- 47. Simopoulos AP (2002). The importance of the ratio of omega-6/omega-3 essential fatty acids. Biomed Pharmacother 56(8):365-79
- 48. Simopoulos AP (2008). The importance of the omega-6/omega-3 fatty acid ratio in cardiovascular disease and other chronic diseases. Exp Biol Med (Maywood) 233(6):674-88
- 49. Blondeau N, Lipsky RH, Bourourou M, Duncan MW, Gorelick PB, Marini AM (2015). Alpha-linolenic acid: an omega-3 fatty acid with neuroprotective properties-ready for use in the stroke clinic? Biomed Res Int 2015:519830
- 50. Das UN (2020). Can bioactive lipids inactivate coronavirus (COVID-19)? Arch Med Res 51(3):282-286
- 51. Weill P, Plissonneau C, Legrand P, Rioux V, Thibault R (2020). May omega-3 fatty acid dietary supplementation help reduce severe complications in Covid-19 patients? Biochimie 179:275-280
- 52. Schiessel DL, Yamazaki RK, Kryczyk M, Coelho I, Yamaguchi AA, Pequito DC, et al (2015). a-Linolenic fatty acid supplementation decreases tumor growth and cachexia parameters in Walker 256 tumor-bearing rats. Nutr Cancer 67(5):839-46
- 53. Gutiérrez S, Svahn SL, Johansson ME (2019). Effects of omega-3 fatty acids on immune cells. Int J Mol Sci 20(20):5028

- 54. Serhan CN, Chiang N, Van Dyke TE (2008). Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators. Nat Rev Immunol 8(5):349-61
- 55. Rius B, López-Vicario C, González-Périz A, Morán-Salvador E, García-Alonso V, Clária J, et al (2012). Resolution of inflammation in obesity-induced liver disease. Front Immunol 3:257
- 56. Lu Y, Liu DX, Tam JP (2008). Lipid rafts are involved in SARS-CoV entry into Vero E6 cells. Biochem Biophys Res Commun 369(2):344–349
- 57. Li GM, Li YG, Yamate M, Li SM, Ikuta K (2007). Lipid rafts play an important role in the early stage of severe acute respiratory syndrome-coronavirus life cycle. Microbes Infect 9(1):96-102
- 58. Goc A, Niedzwiecki A, Rath M (2021). Polyunsaturated ω -3 fatty acids inhibit ACE2-controlled SARS-CoV-2 binding and cellular entry. Research Square Ahead of print
- 59. Fritsche K (2006). Fatty acids as modulators of the immune response. Annu Rev Nutr 26:45-73
- 60. Vedin I, Cederholm T, Freund Levi Y, Basun H, Garlind A, et al (2008). Effects of docosahexaenoic acid-rich n-3 fatty acid supplementation on cytokine release from blood mononuclear leukocytes: The OmegAD study. Am J Clin Nutr 87(6):1616-22
- 61. Kiecolt-Glaser JK, Belury MA, Andridge R, Malarkey WB, Hwang BS, Glaser R (2012). Omega-3 supplementation lowers inflammation in healthy middle-aged and older adults: a randomized controlled trial. Brain Behav Immun 26(6):988-95
- 62. Chiang KC, Chen TC (2009). Vitamin D for the prevention and treatment of pancreatic cancer. World J Gastroenterol 15(27):3349-54
- 63. Barragan M, Good M, Kolls JK (2015). Regulation of dendritic cell function by vitamin D. Nutrients 7(9):8127-51
- 64. Sassi F, Tamone C, D'Amelio P (2018). Vitamin D: Nutrient, hormone, and immunomodulator. Nutrients 10(11):1656
- 65. Xu Y, Baylink DJ, Chen CS, Reeves ME, Xiao J, Lacy C, et al (2020). The importance of vitamin D metabolism as a potential prophylactic, immunoregulatory and neuroprotective treatment for COVID-19. J Transl Med 18(1):322
- 66. Ohadian Moghadam S (2020). A review on currently available potential therapeutic options for COVID-19. Int J Gen Med 13:443-467
- 67. Munshi R, Hussein MH, Toraih EA, Elshazli RM, Jardak C, Sultana N, et al (2020). Vitamin D insufficiency as a potential culprit in critical COVID-19 patients. J Med Virol Ahead of print
- 68. Chaabouni M, Feki W, Chaabouni K, Kammoun S (2020). Vitamin D supplementation to prevent COVID-19 in patients with COPD: A research perspective. Adv Respir Med 88(4):364-365
- 69. Grant WB, Lahore H, McDonnell SL, Baggerly CA, French CB, Aliano JL, et al (2020). Evidence that vitamin D supplementation could reduce risk of influenza and COVID-19 infections and deaths. Nutrients 12(4):988
- 70. Rhodes JM, Subramanian S, Laird E, Kenny RA (2020). Editorial: low population mortality from COVID-19 in countries south of latitude 35 degrees North supports vitamin D as a factor determining severity. Aliment Pharmacol Ther 51(12):1434-1437
- 71. Mandal AKJ, Baktash V, Hosack T, Missouris CG (2020). Vitamin D status and COVID-19 in older adults. Aging Clin Exp Res 32(11):2425-2426

- 72. Dixon KM, Tongkao-On W, Sequeira VB, Carter SE, Song EJ, Rybchyn MS, Gordon-Thomson C, Mason RS (2013). Vitamin D and death by sunshine. Int J Mol Sci 14(1):1964-77
- 73. Garland CF, Garland FC, Gorham ED (2006). The role of vitamin D in cancer prevention. Am J Public Health 96(2):252–261
- 74. Macdonald HM (2013). Contributions of sunlight and diet to vitamin D status. Calcif Tissue Int 92(2):163-76
- 75. Dobnig H (2011). A review of the health consequences of the vitamin D deficiency pandemic. J Neurol Sci 311(1-2):15-8
- 76. Hoel DG, Berwick M, de Gruijl FR, Holick MF (2016). The risks and benefits of sun exposure 2016. Dermatoendocrinol 8(1):e1248325
- 77. van der Rhee H, de Vries E, Coomans C, van de Velde P, Coebergh JW (2016). For better or for worse? A review of positive and negative effects of sun exposure. Cancer Res Front 2(2):156-183
- Green RJ, Samy G, Miqdady MS, El-Hodhod M, Akinyinka OO, Saleh G, et al (2015). Vitamin D deficiency and insufficiency in Africa and the Middle East, despite year-round sunny days. S Afr Med J 105(7):603-5
- 79. Li H, Zhong X, Li W, Wang Q (2019). Effects of 1,25-dihydroxyvitamin D3 on experimental periodontitis and AhR/NF-κB/NLRP3 inflammasome pathway in a mouse model. J Appl Oral Sci 27:e20180713
- Bilezikian JP, Bikle D, Hewison M, Lazaretti-Castro M, Formenti AM, Gupta A, et al (2020). MECHANISMS IN ENDOCRINOLOGY: Vitamin D and COVID-19. Eur J Endocrinol 183(5):R133-R147
- 81. Simonson W (2020). Vitamin D and Coronavirus. Geriatr Nurs 41(4):496-497
- 82. Al-Ani M, Elemam NM, Hundt JE, Maghazachi AA (2020). Drugs for multiple sclerosis activate natural kKiller cells: Do they protect against COVID-19 infection? Infect Drug Resist 13:3243-3254
- 83. Adams KK, Baker WL, Sobieraj DM (2020). Myth busters: Dietary supplements and COVID-19. Ann Pharmacother 54(8):820-826
- 84. Ali N (2020). Role of vitamin D in preventing of COVID-19 infection, progression and severity. J Infect Public Health 13(10):1373-1380
- 85. Kara M, Ekiz T, Ricci V, Kara Ö, Chang KV, Özçakar L (2020). 'Scientific Strabismus' or two related pandemics: coronavirus disease and vitamin D deficiency. Br J Nutr 124(7):736-741
- 86. Allegra A, Tonacci A, Pioggia G, Musolino C, Gangemi S (2020). Vitamin deficiency as risk factor for SARS-CoV-2 infection: correlation with susceptibility and prognosis. Eur Rev Med Pharmacol Sci 24(18):9721-9738
- 87. Ilie PC, Stefanescu S, Smith L (2020). The role of vitamin D in the prevention of coronavirus disease 2019 infection and mortality. Aging Clin Exp Res 32(7):1195-1198
- 88. Annweiler C, Cao Z, Sabatier JM (2020). Point of view: Should COVID-19 patients be supplemented with vitamin D? Maruritas 140:24-26
- 89. Baktash V, Hosack T, Patel N, Shah S, Kandiah P, Van Den Abbeele K, et al (2021). Vitamin D status and outcomes for hospitalised older patients with COVID-19. Postgrad Med J Ahead of print
- 90. Carpagnano GE, Di Lecce V, Quaranta VN, Zito A, Buonamico E, Capozza E, et al (2020). Vitamin D deficiency as a predictor of poor prognosis in patients with acute respiratory failure due to COVID-19. J Endocrinol Invest 9:1-7
- 91. Hadizadeh F (2021). Supplementation with vitamin D in the COVID-19 pandemic? Nutr Rev 79(2):200-208

- 92. Kaufman HW, Niles JK, Kroll MH, Bi C, Holick MF (2020). SARS-CoV-2 positivity rates associated with circulating 25-hydroxyvitamin D levels. PLoS One 15(9):e0239252
- 93. Maghbooli Z, Sahraian MA, Ebrahimi M, Pazoki M, Kafan S, Tabriz HM, et al (2020). Vitamin D sufficiency, a serum 25-hydroxyvitamin D at least 30 ng/mL reduced risk for adverse clinical outcomes in patients with COVID-19 infection. PLoS One 15(9):e0239799
- 94. Merzon E, Tworowski D, Gorohovski A, Vinker S, Golan Cohen A, Green I, et al (2020). Low plasma 25(OH) vitamin D level is associated with increased risk of COVID-19 infection: an Israeli population-based study. FEBS J 287(17):3693-3702
- 95. Moozhipurath RK, Kraft L, Skiera B (2020). Evidence of protective role of Ultraviolet-B (UVB) radiation in reducing COVID-19 deaths. Sci Rep 10(1):17705
- 96. Weir EK, Thenappan T, Bhargava M, Chen Y (2020). Does vitamin D deficiency increase the severity of COVID-19? Clin Med (Lond) 20(4):e107-e108
- 97. Zemb P, Bergman P, Camargo CA Jr, Cavalier E, Cormier C, Courbebaisse M, et al (2020). Vitamin D deficiency and the COVID-19 pandemic. J Glob Antimicrob Resist 22:133-134
- 98. Entrenas Castillo M, Entrenas Costa LM, Vaquero Barrios JM, Alcalá Díaz JF, López Miranda J, Bouillon R, et al (2020). Effect of calcifediol treatment and best available therapy versus best available therapy on intensive care unit admission and mortality among patients hospitalized for COVID-19: A pilot randomized clinical study. J Steroid Biochem Mol Biol 203:105751
- 99. Tarazona-Santabalbina FJ, Cuadra L, Cancio JM, Carbonell FR, Garrote JMP, Casas-Herrero Á, Martínez-Velilla N, Serra-Rexach JA, Formiga F (2021). VitaminD supplementation for the prevention and treatment of COVID-19: a position statement from the Spanish Society of Geriatrics and Gerontology. Rev Esp Geriatr Gerontol Ahead of print
- 100. Lau FH, Majumder R, Torabi R, Saeg F, Hoffman R, Cirillo JD, et al (2021). Vitamin D insufficiency is prevalent in severe COVID-19. MedRxiv preprint Ahead of print
- 101. de Haan K, Groeneveld AB, de Geus HR, Egal M, Struijs A (2014). Vitamin D deficiency as a risk factor for infection, sepsis and mortality in the critically ill: systematic review and meta-analysis. Crit Care 18(6):660
- 102. Claro da Silva T, Hiller C, Gai Z, Kullak-Ublick GA (2016). Vitamin D3 transactivates the zinc and manganese transporter SLC30A10 via the Vitamin D receptor. J Steroid Biochem Mol Biol 163:77–87
- 103. McDonnell SL, Baggerly CA, French CB, Baggerly LL, Garland CF, Gorham ED, et al (2018). Breast cancer risk markedly lower with serum 25hydroxyvitamin D concentrations ≥60 vs <20 ng/ml (150 vs 50 nmol/L): Pooled analysis of two randomized trials and a prospective cohort. PLoS One 13(6):e0199265
- 104. McCullough ML, Zoltick ES, Weinstein SJ, Fedirko V, Wang M, Cook NR, et al (2019). Circulating vitamin D and colorectal cancer risk: An international pooling project of 17 cohorts. J Natl Cancer Inst 111(2):158-169
- 105. Chandler PD, Chen WY, Ajala ON, Hazra A, Cook N, Bubes V, et al (2020). Effect of vitamin D3 supplements on development of advanced cancer: A secondary analysis of the VITAL randomized clinical trial. JAMA Netw Open 3(11):e2025850

- 106. Moukayed M, Grant WB (2017). The roles of UVB and vitamin D in reducing risk of cancer incidence and mortality: A review of the epidemiology, clinical trials, and mechanisms. Rev Endocr Metab Disord 18(2):167-182
- 107. Ma Y, Johnson CS, Trump DL (2016). Mechanistic insights of vitamin D anticancer effects. Vitam Horm 100:395-431
- 108. Abu El Maaty MA, Wölfl S (2017). Effects of 1,25(OH)₂D₃ on cancer cells and potential applications in combination with established and putative anti-cancer agents. Nutrients 9(1):87
- 109. D'Avolio A, Avataneo V, Manca A, Cusato J, De Nicolò A, Lucchini R, et al (2020). 25-Hydroxyvitamin D concentrations are lower in patients with positive PCR for SARS-CoV-2. Nutrients 12(5):1359
- 110. Charoenngam N, Holick MF (2020). Immunologic Effects of vitamin D on human health and disease. Nutrients 12(7):E2097
- 111. Kumar R, Rathi H, Haq A, Wimalawansa SJ, Sharma A (2021). Putative roles of vitamin D in modulating immune response and immunopathology associated with COVID-19. Virus Res 292:198235
- 112. Dominguez LJ, Veronese N, Guerrero-Romero F, Barbagallo M (2021). Magnesium in infectious diseases in older people. Nutrients 13(1):180
- 113. Iotti S, Wolf F, Mazur A, Maier JA (2020). The COVID-19 pandemic: is there a role for magnesium? Hypotheses and perspectives. Magnes Res 33(2):21-27
- 114. Micke O, Vormann J, Kisters K (2021). Magnesium and COVID-19 Some further comments - A commentary on Wallace TC. Combating COVID-19 and building immune resilience: A potential role for magnesium nutrition? J Am Coll Nutr Ahead of print
- 115. Alamdari NM, Afaghi S, Rahimi FS, Tarki FE, Tavana S, Zali A, et al (2020). Mortality risk factors among hospitalized COVID-19 patients in a major referral center in Iran. Tohoku J Exp Med 252(1):73-84
- 116. Quilliot D, Bonsack O, Jaussaud R, Mazur A (2020). Dysmagnesemia in Covid-19 cohort patients: prevalence and associated factors. Magnes Res 33(4):114-122
- 117. Zeng HL, Yang Q, Yuan P, Wang X, Cheng L (2021). Associations of essential and toxic metals/metalloids in whole blood with both disease severity and mortality in patients with COVID-19. FASEB J 35(3):e21392
- 118. Costello RB, Nielsen F (2017). Interpreting magnesium status to enhance clinical care: key indicators. Curr Opin Clin Nutr Metab Care 20(6):504-511
- 119. Nielsen FH (2018). Magnesium deficiency and increased inflammation: current perspectives. J Inflamm Res 11:25-34
- 120. Tang CF, Ding H, Jiao RQ, Wu XX, Kong LD (2020). Possibility of magnesium supplementation for supportive treatment in patients with COVID-19. Eur J Pharmacol 886:173546
- 121. Wallace TC (2020). Combating COVID-19 and building immune resilience: A potential role for magnesium nutrition? J Am Coll Nutr 39(8):685-693
- 122. DiNicolantonio JJ, O'Keefe JH (2021). Magnesium and Vitamin D Deficiency as a Potential Cause of Immune Dysfunction, Cytokine Storm and Disseminated Intravascular Coagulation in covid-19 patients. Mo Med 118(1):68-73
- 123. Dai Q, Zhu X, Manson JE, Song Y, Li X, Franke AA, et al (2018). Magnesium status and supplementation influence vitamin D status and metabolism: results from a randomized trial. Am J Clin Nutr 108(6):1249-1258
- 124. de Baaij JH, Hoenderop JG, Bindels RJ (2015). Magnesium in man: implications for health and disease. Physiol Rev 95(1):1-46

- 125. Maier JA, Castiglioni S, Locatelli L, Zocchi M, Mazur A (2020). Magnesium and inflammation: Advances and perspectives. Semin Cell Dev Biol Ahead of print
- 126. Chacko SA, Song Y, Nathan L, Tinker L, de Boer IH, Tylavsky F, et al (2010). Relations of dietary magnesium intake to biomarkers of inflammation and endothelial dysfunction in an ethnically diverse cohort of postmenopausal women. Diabetes Care 33(2):304-10
- 127. Qu X, Jin F, Hao Y, Li H, Tang T, Wang H, et al (2013). Magnesium and the risk of cardiovascular events: a meta-analysis of prospective cohort studies. PLoS One 8(3):e57720
- 128. Zipeto D, Palmeira JDF, Argañaraz GA, Argañaraz ER (2020). ACE2/ADAM17/TMPRSS2 interplay may be the main risk factor for COVID-19. Front Immunol 11:576745
- 129. Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O, et al (2020). Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. Science 367(6483):1260-1263
- 130. Martín Giménez VM, Inserra F, Tajer CD, Mariani J, Ferder L, Reiter RJ, et al (2020). Lungs as target of COVID-19 infection: Protective common molecular mechanisms of vitamin D and melatonin as a new potential synergistic treatment. Life Sci 254:117808
- 131. Zhang H, Penninger JM, Li Y, Zhong N, Slutsky AS (2020). Angiotensinconverting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. Intensive Care Med 46(4):586–590
- 132. Hathaway D, Pandav K, Patel M, Riva-Moscoso A, Singh BM, Patel A, et al (2020). Omega 3 fatty acids and COVID-19: A comprehensive review. Infect Chemother 52(4):478-495
- 133. Aygun H (2020). Vitamin D can prevent COVID-19 infection-induced multiple organ damage. Naunyn Schmiedebergs Arch Pharmacol 393(7):1157-1160
- 134. Kumar D, Gupta P, Banerjee D (2020). Letter: does vitamin D have a potential role against COVID-19? Aliment Pharmacol Ther 52(2):409-411
- 135. Xu J, Yang J, Chen J, Luo Q, Zhang Q, Zhang H (2017). Vitamin D alleviates lipopolysaccharide-induced acute lung injury via regulation of the renin-angiotensin system. Mol Med Rep 16(5):7432-7438
- 136. Xiao D, Li X, Su X, Mu D, Qu Y (2020). Could SARS-CoV-2-induced lung injury be attenuated by vitamin D? Int J Infect Dis 102:196-202
- 137. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al (2020). A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature 579(7798):270-273
- 138. Biggs L, Yu C, Fedoric B, Lopez AF, Galli SJ, Grimbaldeston MA (2010). Evidence that vitamin D(3) promotes mast cell-dependent reduction of chronic UVB-induced skin pathology in mice. J Exp Med 207(3):455-63
- 139. Skrajnowska D, Bobrowska-Korczak B (2019). Role of zinc in immune system and anti-cancer defense mechanisms. Nutrients 11(10):2273
- 140. Mohan M, Cherian JJ, Sharma A (2020). Exploring links between vitamin D deficiency and COVID-19. PLoS Pathog 16(9):e1008874
- 141. Stone KD, Prussin C, Metcalfe DD (2010). IgE, mast cells, basophils, and eosinophils. J Allergy Clin Immunol 125(2 Suppl 2):S73-80
- 142. Kempuraj D, Selvakumar GP, Ahmed ME, Raikwar SP, Thangavel R, Khan A, et al (2020). COVID-19, mast cells, cytokine storm, psychological stress, and neuroinflammation. Neuroscientist 26(5-6):402-414

- 143. Graham AC, Temple RM, Obar JJ (2015). Mast cells and influenza A virus: association with allergic responses and beyond. Front Immunol 6:238
- 144. Wang X, Kulka M (2015). n-3 Polyunsaturated fatty acids and mast cell activation. J Leukoc Biol 97(5):859-71
- 145. Mukai K, Tsai M, Saito H, Galli SJ (2018). Mast cells as sources of cytokines, chemokines, and growth factors. Immunol Rev 282(1):121-150
- 146. Kawakami T, Ando T, Kimura M, Wilson BS, Kawakami Y (2009). Mast cells in atopic dermatitis. Curr Opin Immunol 21(6):666-78
- 147. Baeke F, Takiishi T, Korf H, Gysemans C, Mathieu C (2010). Vitamin D: modulator of the immune system. Curr Opin Pharmacol 10(4):482-96
- 148. Meftahi GH, Jangravi Z, Sahraei H, Bahari Z (2020). The possible pathophysiology mechanism of cytokine storm in elderly adults with COVID-19 infection: the contribution of "inflame-aging". Inflamm Res 69(9):825-839
- 149. Jafarzadeh A, Chauhan P, Saha B, Jafarzadeh S, Nemati M (2020). Contribution of monocytes and macrophages to the local tissue inflammation and cytokine storm in COVID-19: Lessons from SARS and MERS, and potential therapeutic interventions. Life Sci 257:118102
- 150. Harihar1an A, Hakeem AR, Radhakrishnan S, Reddy MS, Rela M (2020). The role and therapeutic potential of NF-kappa-B pathway in severe COVID-19 patients. Inflammopharmacology 7:1–10
- 151. Wensveen FM, Valentić S, Šestan M, Turk Wensveen T, Polić B (2015). The "Big Bang" in obese fat: Events initiating obesity-induced adipose tissue inflammation. Eur J Immunol 45(9):2446-56
- 152. Blanchard C, Rothenberg ME (2009). Biology of the eosinophil. Adv Immunol 101:81-121
- 153. Hogan SP, Rosenberg HF, Moqbel R, Phipps S, Foster PS, Lacy P, Kay AB, Rothenberg ME (2008). Eosinophils: biological properties and role in health and disease. Clin Exp Allergy 38(5):709-50
- 154. Davoine F, Lacy P (2014). Eosinophil cytokines, chemokines, and growth factors: emerging roles in immunity. Front Immunol 5:570
- 155. Pacha O, Sallman MA, Evans SE (2020). COVID-19: a case for inhibiting IL-17? Nat Rev Immunol 20(6):345-346
- 156. Aggarwal N, Korenbaum E, Mahadeva R, Immenschuh S, Grau V, Dinarello CA, et al (2016). α -Linoleic acid enhances the capacity of α -1 antitrypsin to inhibit lipopolysaccharide-induced IL-1 β in human blood neutrophils. Mol Med 22:680-93
- 157. Mantovani A, Cassatella MA, Costantini C, Jaillon S (2011). Neutrophils in the activation and regulation of innate and adaptive immunity. Nat Rev Immunol 11(8):519-31
- 158. Tecchio C, Micheletti A, Cassatella MA (2014). Neutrophil-derived cytokines: facts beyond expression. Front Immunol 5:508
- 159. Lau YL, Peiris JS, Law HK (2012). Role of dendritic cells in SARS coronavirus infection. Hong Kong Med J 18 Suppl 3:28-30
- 160. Shibabaw T (2020). Inflammatory Cytokine: IL-17A Signaling pathway in patients present with COVID-19 and current treatment strategy. J Inflamm Res 13:673-680
- 161. Marone G, Columbo M, de Paulis A, Cirillo R, Giugliano R, Condorelli M (1986). Physiological concentrations of zinc inhibit the release of histamine from human basophils and lung mast cells. Agents Actions 18(1-2):103-6

- 162. Feltis BN, Elbaz A, Wright PF, Mackay GA, Turney TW, Lopata AL (2015). Characterizing the inhibitory action of zinc oxide nanoparticles on allergictype mast cell activation. Mol Immunol 66(2):139-46
- 163. Gueck T, Seidel A, Fuhrmann H (2003). Effects of essential fatty acids on mediators of mast cells in culture. Prostaglandins Leukot Essent Fatty Acids 68(5):317-22
- 164. Park BK, Park S, Park JB, Park MC, Min TS, Jin M (2013). Omega-3 fatty acids suppress Th2-associated cytokine gene expressions and GATA transcription factors in mast cells. J Nutr Biochem 24(5):868-76
- 165. Yip KH, Kolesnikoff N, Yu C, Hauschild N, Taing H, Biggs L, et al (2014). Mech,anisms of vitamin D₃ metabolite repression of IgE-dependent mast cell activation. J Allergy Clin Immunol 133(5):1356-64
- 166. Liu ZQ, Li XX, Qiu SQ, Yu Y, Li MG, Yang LT, et al (2017). Vitamin D contributes to mast cell stabilization. Allergy 72(8):1184-1192
- 167. Zhao JW, Ping JD, Wang YF, Liu XN, Li N, Hu ZL, et al (2020). Vitamin D suppress the production of vascular endothelial growth factor in mast cell by inhibiting PI3K/Akt/p38 MAPK/HIF-1a pathway in chronic spontaneous urticaria. Clin Immunol 215:108444
- 168. Takemoto S, Yamamoto A, Tomonaga S, Funaba M, Matsui T (2013). Magnesium deficiency induces the emergence of mast cells in the liver of rats. J Nutr Sci Vitaminol (Tokyo) 59(6):560-3
- 169. Ohbori K, Fujiwara M, Ohishi A, Nishida K, Uozumi Y, Nagasawa K (2017). Prophylactic oral administration of magnesium ameliorates dextran sulfate sodium-induced colitis in mice through a decrease of colonic accumulation of P2X7 receptor-expressing mast cells. Biol Pharm Bull 40(7):1071-1077
- 170. Haase H, Rink L (2007). Signal transduction in monocytes: the role of zinc ions. Biometals 20(3-4):579-85
- 171. Lu H, Xin Y, Tang Y, Shao G (2012). Zinc suppressed the airway inflammation in asthmatic rats: effects of zinc on generation of eotaxin, MCP-1, IL-8, IL-4, and IFN-γ. Biol Trace Elem Res 150(1-3):314-21
- 172. Cho E, Park Y (2016). Association between serum fatty acid composition and innate immune markers in healthy adults. Nutr Res Pract 10(2):182-7
- 173. Snodgrass RG, Huang S, Namgaladze D, Jandali O, Shao T, Sama S, et al (2016). Docosahexaenoic acid and palmitic acid reciprocally modulate monocyte activation in part through endoplasmic reticulum stress. J Nutr Biochem 32:39-45
- 174. Sugimoto J, Romani AM, Valentin-Torres AM, Luciano AA, Ramirez Kitchen CM, Funderburg N, et al (2012). Magnesium decreases inflammatory cytokine production: a novel innate immunomodulatory mechanism. J Immunol 188(12):6338-46
- 175. Sun L, Li X, Xu M, Yang F, Wang W, Niu X (2020). In vitro immunomodulation of magnesium on monocytic cell toward anti-inflammatory macrophages. Regen Biomater 7(4):391-401
- 176. Aude-Garcia C, Dalzon B, Ravanat JL, Collin-Faure V, Diemer H, Strub JM, et al (2016). A combined proteomic and targeted analysis unravels new toxic mechanisms for zinc oxide nanoparticles in macrophages. J Proteomics 134:174-185
- 177. Zhao G, Etherton TD, Martin KR, Vanden Heuvel JP, Gillies PJ, West SG, et al (2005). Anti-inflammatory effects of polyunsaturated fatty acids in THP-1 cells. Biochem Biophys Res Commun 336(3):909-17
- 178. Rao Z, Zhang N, Xu N, Pan Y, Xiao M, Wu J, et al (2017). 1,25-Dihydroxyvitamin D inhibits LPS-induced high-mobility group box 1

(HMGB1) secretion via targeting the NF-E2-related factor 2hemeoxygenase-1-HMGB1 pathway in macrophages. Front Immunol 8:1308

- 179. Helming L, Böse J, Ehrchen J, Schiebe S, Frahm T, Geffers R, et al (2005). 1alpha,25-Dihydroxyvitamin D3 is a potent suppressor of interferon gamma-mediated macrophage activation. Blood 106(13):4351-8
- 180. Karkeni E, Morin SO, Bou Tayeh B, Goubard A, Josselin E, Castellano R, et al (2019). Vitamin D controls tumor growth and CD8+ T cell infiltration in breast cancer. Front Immunol 10:1307
- 181. Turner DL, Ford WR, Kidd EJ, Broadley KJ, Powell C (2017). Effects of nebulised magnesium sulphate on inflammation and function of the guineapig airway. Eur J Pharmacol 801:79-85
- 182. Hu T, Xu H, Wang C, Qin H, An Z (2018). Magnesium enhances the chondrogenic differentiation of mesenchymal stem cells by inhibiting activated macrophage-induced inflammation. Sci Rep 8(1):3406
- 183. Finamore A, Massimi M, Conti Devirgiliis L, Mengheri E (2008). Zinc deficiency induces membrane barrier damage and increases neutrophil transmigration in Caco-2 cells. J Nutr 138(9):1664-70
- 184. Sakakibara Y, Sato S, Kawashima Y, Someya Y, Shirato K, Tachiyashiki K, et al (2011). Different recovery responses from dietary zinc-deficiency in the distribution of rat granulocytes. J Nutr Sci Vitaminol (Tokyo) 57(2):197-201
- 185. Yoshida S, Yasutomo K, Watanabe T (2016). Treatment with DHA/EPA ameliorates atopic dermatitis-like skin disease by blocking LTB4 production. J Med Invest 63(3-4):187-91
- 186. Chang YF, Hou YC, Pai MH, Yeh SL, Liu JJ (2017). Effects of ω -3 polyunsaturated fatty acids on the homeostasis of CD4+ T cells and lung injury in mice with polymicrobial sepsis. J Parenter Enteral Nutr 41(5):805-814
- 187. Akbas EM, Gungor A, Ozcicek A, Akbas N, Askin S, Polat M (2016). Vitamin D and inflammation: evaluation with neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio. Arch Med Sci 12(4):721-7
- 188. Tang Y, Liu J, Yan Y, Fang H, Guo C, Xie R, et al (2018). 1,25dihydroxyvitamin-D3 promotes neutrophil apoptosis in periodontitis with type 2 diabetes mellitus patients via the p38/MAPK pathway. Medicine (Baltimore) 97(52):e13903
- 189. Li W, Wu X, Yu J, Ma C, Zhuang P, Zeng J, Zhang J, Deng G, Wang Y (2019). Magnesium sulfate attenuates lipopolysaccharides-induced acute lung injury in mice. Chin J Physiol 62(5):203-209
- 190. Kitamura H, Morikawa H, Kamon H, Iguchi M, Hojyo S, Fukada T, et al (2006). Toll-like receptor-mediated regulation of zinc homeostasis influences dendritic cell function. Nat Immunol 7(9):971-7
- 191. Shumilina E, Xuan NT, Schmid E, Bhavsar SK, Szteyn K, Gu S, et al (2010). Zinc induced apoptotic death of mouse dendritic cells. Apoptosis 15(10):1177-86
- 192. Zeyda M, Säemann MD, Stuhlmeier KM, Mascher DG, Nowotny PN, Zlabinger GJ, et al (2005). Association between serum fatty acid composition and innate immune markers in healthy adults. J Biol Chem 280(14):14293-301
- 193. Kong W, Yen JH, Ganea D (2011). Docosahexaenoic acid prevents dendritic cell maturation, inhibits antigen-specific Th1/Th17 differentiation and

suppresses experimental autoimmune encephalomyelitis. Brain Behav Immun 25(5):872-82

- 194. Vanherwegen AS, Eelen G, Ferreira GB, Ghesquière B, Cook DP, Nikolic T, et al (2019). Vitamin D controls the capacity of human dendritic cells to induce functional regulatory T cells by regulation of glucose metabolism. J Steroid Biochem Mol Biol 187:134-145
- 195. Piemonti L, Monti P, Sironi M, Fraticelli P, Leone BE, Dal Cin E, et al (2000). Vitamin D3 affects differentiation, maturation, and function of human monocyte-derived dendritic cells. J Immunol 164(9):4443-51
- 196. Libako P, Miller J, Nowacki W, Castiglioni S, Maier JA, Mazur A (2015). Extracellular Mg concentration and Ca blockers modulate the initial steps of the response of Th2 lymphocytes in co-culture with macrophages and dendritic cells. Eur Cytokine Netw 26(1):1-9
- 197. Richter M, Bonneau R, Girard MA, Beaulieu C, Larivée P (2003). Zinc status modulates bronchopulmonary eosinophil infiltration in a murine model of allergic inflammation. Chest 123(3 Suppl):446S
- 198. Lang C, Murgia C, Leong M, Tan LW, Perozzi G, Knight D, et al (2007). Antiinflammatory effects of zinc and alterations in zinc transporter mRNA in mouse models of allergic inflammation. Am J Physiol Lung Cell Mol Physiol 292(2):L577-84
- 199. Tanigai T, Ueki S, Kihara J, Kamada R, Yamauchi Y, Sokal A, et al (2012). Docosahexaenoic acid exerts anti-inflammatory action on human eosinophils through peroxisome proliferator-activated receptor-independent mechanisms. Int Arch Allergy Immunol 158(4):375-86
- 200. Snyman JR, de Sommers K, Steinmann MA, Lizamore DJ (1997). Effects of calcitriol on eosinophil activity and antibody responses in patients with schistosomiasis. Eur J Clin Pharmacol 52(4):277-80
- 201. Souto Filho JTD, de Andrade AS, Ribeiro FM, Alves PAS, Simonini VRF (2018). Impact of vitamin D deficiency on increased blood eosinophil counts. Hematol Oncol Stem Cell Ther 11(1):25-29
- 202. Hungerford GF, Karson EF (1960). The eosinophilia of magnesium deficiency. Blood 16(5):1642-1650
- 203. Prasad AS (2000). Effects of zinc deficiency on Th1 and Th2 cytokine shifts. J Infect Dis 182 Suppl 1:S62-8
- 204. Bao B, Thakur A, Li Y, Ahmad A, Azmi AS, Banerjee S, et al (2012). The immunological contribution of NF-κB within the tumor microenvironment: a potential protective role of zinc as an anti-tumor agent. Biochim Biophys Acta Reviews on Cancer 1825(2):160-72
- 205. Chang HH, Chen CS, Lin JY (2008). Dietary perilla oil inhibits proinflammatory cytokine production in the bronchoalveolar lavage fluid of ovalbumin-challenged mice. Lipids 43(6):499-506
- 206. Mizota T, Fujita-Kambara C, Matsuya N, Hamasaki S, Fukudome T, Goto H, et al (2009). Effect of dietary fatty acid composition on Th1/Th2 polarization in lymphocytes. J Parenter Enteral Nutr 33(4):390-6
- 207. Zhang P, Smith R, Chapkin RS, McMurray DN (2005). Dietary (n-3) polyunsaturated fatty acids modulate murine Th1/Th2 balance toward the Th2 pole by suppression of Th1 development. J Nutr 135(7):1745-51
- 208. Chung HS, Park CS, Hong SH, Lee S, Cho ML, Her YM, et al (2013). Effects of magnesium pretreatment on the levels of T helper cytokines and on the severity of reperfusion syndrome in patients undergoing living donor liver transplantation. Magnes Res 26(2):46-55

- 209. Kitabayashi C, Fukada T, Kanamoto M, Ohashi W, Hojyo S, Atsumi T, et al (2010). Zinc suppresses Th17 development via inhibition of STAT3 activation. Int Immunol 22(5):375-86
- 210. Lee H, Kim B, Choi YH, Hwang Y, Kim DH, Cho S, et al (2015). Inhibition of interleukin-1β-mediated interleukin-1 receptor-associated kinase 4 phosphorylation by zinc leads to repression of memory T helper type 17 response in humans. Immunology 146(4):645-56
- 211. Rosenkranz E, Maywald M, Hilgers RD, Brieger A, Clarner T, Kipp M, et al (2016). Induction of regulatory T cells in Th1-/Th17-driven experimental autoimmune encephalomyelitis by zinc administration. J Nutr Biochem 29:116-23
- 212. Monk JM, Hou TY, Turk HF, Weeks B, Wu C, McMurray DN, et al (2012). Dietary n-3 polyunsaturated fatty acids (PUFA) decrease obesity-associated Th17 cell-mediated inflammation during colitis. PLoS One 7(11):e49739
- 213. Han SC, Koo DH, Kang NJ, Yoon WJ, Kang GJ, Kang HK, et al (2015). Docosahexaenoic acid alleviates atopic dermatitis by generating Tregs and IL-10/TGF-β-modified macrophages via a TGF-β-dependent mechanism. J Invest Dermatol 135(6):1556-64
- 214. Huang CH, Hou YC, Pai MH, Yeh CL, Yeh SL (2017). Dietary ω -6/ ω -3 PUFA ratios affect the homeostasis of Th/Treg cells in mice with dextran sulfate sodium-induced colitis. J Parenter Enteral Nutr 41(4):647-56
- 215. Daniel C, Sartory NA, Zahn N, Radeke HH, Stein JM (2008). Immune modulatory treatment of trinitrobenzene sulfonic acid colitis with calcitriol is associated with a change of a T helper (Th) 1/Th17 to a Th2 and regulatory T cell profile. J Pharmacol Exp Ther 324(1):23-33
- 216. Tang J, Zhou R, Luger D, Zhu W, Silver PB, Grajewski RS, et al (2009). Calcitriol suppresses antiretinal autoimmunity through inhibitory effects on the Th17 effector response. J Immunol 182(8):4624-32
- 217. Rosenkranz E, Metz CH, Maywald M, Hilgers RD, Weßels I, Senff T, et al (2016). Zinc supplementation induces regulatory T cells by inhibition of Sirt-1 deacetylase in mixed lymphocyte cultures. Mol Nutr Food Res 60(3):661-71
- 218. Maywald M, Meurer SK, Weiskirchen R, Rink L (2017). Zinc supplementation augments TGF- β 1-dependent regulatory T cell induction. Mol Nutr Food Res 61(3):1600493
- 219. Carlsson JA, Wold AE, Sandberg AS, Östman SM (2015). The polyunsaturated fatty acids arachidonic acid and docosahexaenoic acid induce mouse dendritic cells maturation but reduce T-cell responses in vitro. PLoS One 10(11):e0143741
- 220. Lian M, Luo W, Sui Y, Li Z, Hua J (2015). Dietary n-3 PUFA protects mice from Con A induced liver injury by modulating regulatory T cells and PPARγ expression. PLoS One 10(7):e0132741
- 221. Onodera T, Fukuhara A, Shin J, Hayakawa T, Otsuki M, Shimomura I (2017). Eicosapentaenoic acid and 5-HEPE enhance macrophage-mediated Treg induction in mice. Sci Rep 7(1):4560
- 222. Guillot X, Semerano L, Saidenberg-Kermanac'h N, Falgarone G, Boissier MC (2010). Vitamin D and inflammation. Joint Bone Spine 77(6):552-7
- 223. Gorman S, Geldenhuys S, Judge M, Weeden CE, Waithman J, Hart PH (2016). Dietary vitamin D increases percentages and function of regulatory T cells in the skin-draining lymph nodes and suppresses dermal inflammation. J Immunol Res 2016:1426503

- 224. Fisher SA, Rahimzadeh M, Brierley C, Gration B, Doree C, Kimber CE, et al (2019). The role of vitamin D in increasing circulating T regulatory cell numbers and modulating T regulatory cell phenotypes in patients with inflammatory disease or in healthy volunteers: A systematic review. PLoS One 14(9):e0222313
- 225. Summersgill H, England H, Lopez-Castejon G, Lawrence CB, Luheshi NM, Pahle J, (2014). Zinc depletion regulates the processing and secretion of IL- 1β . Cell Death Dis 5:e1040
- 226. Fan Y, Zhang X, Yang L, Wang J, Hu Y, Bian A, et al (2017). Zinc inhibits high glucose-induced NLRP3 inflammasome activation in human peritoneal mesothelial cells. Mol Med Rep 16(4):5195-5202
- 227. Williams-Bey Y, Boularan C, Vural A, Huang NN, Hwang IY, Shan-Shi C, et al (2014). Omega-3 free fatty acids suppress macrophage inflammasome activation by inhibiting NF-κB activation and enhancing autophagy. PLoS One 9(6):e97957
- 228. De Boer AA, Monk JM, Liddle DM, Hutchinson AL, Power KA, Ma DW, et al (2016). Fish-oil-derived n-3 polyunsaturated fatty acids reduce NLRP3 inflammasome activity and obesity-related inflammatory cross-talk between adipocytes and CD11b(+) macrophages. J Nutr Biochem 34:61-72
- 229. Kumar N, Gupta G, Anilkumar K, Fatima N, Karnati R, Reddy GV, et al (2016). 15-Lipoxygenase metabolites of a-linolenic acid, [13-(S)-HPOTrE and 13-(S)-HOTrE], mediate anti-inflammatory effects by inactivating NLRP3 inflammasome. Sci Rep 6::31649
- 230. Zhang JH, Chen YP, Yang X, Li CQ (2017). Vitamin D3 levels and NLRP3 expression in murine models of obese asthma: association with asthma outcomes. Braz J Med Biol Res 51(1):e6841
- 231. Lu L, Lu Q, Chen W, Li J, Li C, Zheng Z (2018). Vitamin D 3 protects against diabetic retinopathy by inhibiting high-glucose-induced activation of the ROS/TXNIP/NLRP3 inflammasome pathway. J Diabetes Res 2018:8193523
- 232. Dai Y, Zhang J, Xiang J, Li Y, Wu D, Xu J (2019). Calcitriol inhibits ROS-NLRP3-IL-1β signaling axis via activation of Nrf2-antioxidant signaling in hyperosmotic stress stimulated human corneal epithelial cells. Redox Biol 21:101093
- 233. Chang CY, Shih HJ, Huang IT, Tsai PS, Chen KY, Huang CJ (2019). Magnesium sulfate mitigates the progression of monocrotaline pulmonary hypertension in rats. Int J Mol Sci 20(18):4622
- 234. Li C, Chen M, He X, Ouyang D (2021). A mini-review on ion fluxes that regulate NLRP3 inflammasome activation. Acta Biochim Biophys Sin (Shanghai) 53(2):131-139
- 235. von Bülow V, Dubben S, Engelhardt G, Hebel S, Plümäkers B, Heine H, (2007). Zinc-dependent suppression of TNF-alpha production is mediated by protein kinase A-induced inhibition of Raf-1, I kappa B kinase beta, and NF-kappa B. J Immunol 179(6):4180-6
- 236. Novak TE, Babcock TA, Jho DH, Helton WS, Espat NJ (2003). NF-kappa B inhibition by omega -3 fatty acids modulates LPS-stimulated macrophage TNF-alpha transcription. Am J Physiol Lung Cell Mol Physiol 284(1):L84-9
- 237. Rogero MM, Leão MC, Santana TM, Pimentel MVMB, Carlini GCG, da Silveira TFF, et al (2020). Potential benefits and risks of omega-3 fatty acids supplementation to patients with COVID-19. Free Radic Biol Med 156:190-199

- 238. Yu XP, Bellido T, Manolagas SC (1995). Down-regulation of NF-kappa B protein levels in activated human lymphocytes by 1,25-dihydroxyvitamin D3. Proc Natl Acad Sci USA 92(24):10990-4
- 239. Cohen-Lahav M, Shany S, Tobvin D, Chaimovitz C, Douvdevani A (2006). Vitamin D decreases NFkappaB activity by increasing IkappaBalpha levels. Nephrol Dial Transplant 21(4):889-97
- 240. Wöbke TK, Sorg BL, Steinhilber D (2014). Vitamin D in inflammatory diseases. Front Physiol 5:244
- 241. Su NY, Peng TC, Tsai PS, Huang CJ (2013). Phosphoinositide 3-kinase/Akt pathway is involved in mediating the anti-inflammation effects of magnesium sulfate. J Surg Res 185(2):726-32
- 242. Ren J, Chung SH (2007). Anti-inflammatory effect of alpha-linolenic acid and its mode of action through the inhibition of nitric oxide production and inducible nitric oxide synthase gene expression via NF-kappaB and mitogen-activated protein kinase pathways. J Agric Food Chem 55(13):5073-80
- 243. Xie N, Zh,ang W, Li J, Liang H, Zhou H, Duan W, et al (2011). a-Linolenic acid intake attenuates myocardial ischemia/reperfusion injury through antiinflammatory and anti-oxidative stress effects in diabetic but not normal rats. Arch Med Res 42(3):171-81
- 244. Bi X, Li F, Liu S, Jin Y, Zhang X, Yang T, et al (2017). ω -3 polyunsaturated fatty acids ameliorate type 1 diabetes and autoimmunity. J Clin Invest 127(5):1757-1771
- 245. Choi M, Park H, Cho S, Lee M (2013). Vitamin D3 supplementation modulates inflammatory responses from the muscle damage induced by high-intensity exercise in SD rats. Cytokine 63(1):27-35
- 246. Bessler H, Djaldetti M (2012). 1a,25-Dihydroxyvitamin D3 modulates the interaction between immune and colon cancer cells. Biomed Pharmacother 66(6):428-32
- 247. Lysandropoulos AP, Jaquiéry E, Jilek S, Pantaleo G, Schluep M, Du Pasquier RA (2011). Vitamin D has a direct immunomodulatory effect on CD8+ T cells of patients with early multiple sclerosis and healthy control subjects. J Neuroimmunol 233(1-2):240-4
- 248. Lucisano S, Arena A, Stassi G, Iannello D, Montalto G, Romeo A, et al (2015). Role of paricalcitol in modulating the immune response in patients with renal disease. Int J Endocrinol 2015:765364
- 249. Han F, Xu L, Huang Y, Chen T, Zhou T, Yang L (2018). Magnesium sulphate can alleviate oxidative stress and reduce inflammatory cytokines in rat placenta of intrahepatic cholestasis of pregnancy model. Arch Gynecol Obstet 298(3):631-638
- 250. Schubert C, Guttek K, Grüngreiff K, Thielitz A, Bühling F, Reinhold A, et al (2014). Oral zinc aspartate treats experimental autoimmune encephalomyelitis. Biometals 27(6):1249-62
- 251. Cippitelli M, Santoni M (1998). Vitamin D3 : a transcriptional modulator of the interferon- gene. Eur J Immunol 28:3017-30
- 252. Sharifi A, Vahedi H, Nedjat S, Rafiei H, Hosseinzadeh-Attar MJ (2019). Effect of single-dose injection of vitamin D on immune cytokines in ulcerative colitis patients: a randomized placebo-controlled trial. APMIS 127(10):681-687
- 253. Carvalho JTG, Schneider M, Cuppari L, Grabulosa CC, T Aoike D, Q Redublo BM, et al (2017). Cholecalciferol decreases inflammation and improves

vitamin D regulatory enzymes in lymphocytes in the uremic environment: A randomized controlled pilot trial. PLoS One 12(6):e0179540

- 254. Bao S, Liu MJ, Lee B, Besecker B, Lai JP, Guttridge DC, et al (2010). Zinc modulates the innate immune response in vivo to polymicrobial sepsis through regulation of NF-kappaB. Am J Physiol Lung Cell Mol Physiol 298(6):L744-54
- 255. Tyagi A, Kumar U, Reddy S, Santosh VS, Mohammed SB, Ehtesham NZ, et al (2012). Attenuation of colonic inflammation by partial replacement of dietary linoleic acid with a-linolenic acid in a rat model of inflammatory bowel disease. Br J Nutr 108(9):1612-22
- 256. Liu YH, Li XY, Chen CY, Zhang HM, Kang JX (2015). Omega-3 fatty acid intervention suppresses lipopolysaccharide-induced inflammation and weight loss in mice. Mar Drugs 13(2):1026-36
- 257. Cui C, Xu P, Li G, Qiao Y, Han W, Geng C, et al (2019). Vitamin D receptor activation regulates microglia polarization and oxidative stress in spontaneously hypertensive rats and angiotensin II-exposed microglial cells: Role of renin-angiotensin system. Redox Biol 26:101295
- 258. Chang YY, Kao MC, Lin JA, Chen TY, Cheng CF, Wong CS, Tzeng IS, Huang CJ (2018). Effects of MgSO4 on inhibiting Nod-like receptor protein 3 inflammasome involve decreasing intracellular calcium. J Surg Res 221:257-265
- 259. Ozen M, Xie H, Shin N, Al Yousif G, Clemens J, McLane MW, et al (2020). Magnesium sulfate inhibits inflammation through P2X7 receptors in human umbilical vein endothelial cells. Pediatr Res 87(3):463-471
- 260. Uzzo RG, Crispen PL, Golovine K, Makhov P, Horwitz EM, Kolenko VM (2006). Diverse effects of zinc on NF-kappaB and AP-1 transcription factors: implications for prostate cancer progression. Carcinogenesis 27(10):1980-90
- 261. Orrù B, Szekeres-Bartho J, Bizzarri M, Spiga AM, Unfer V (2020). Inhibitory effects of Vitamin D on inflammation and IL-6 release. A further support for COVID-19 management?. Eur Rev Med Pharmacol Sci 24(15):8187-8193
- 262. Silberstein M (2020). Vitamin D: A simpler alternative to tocilizumab for trial in COVID-19?. Med Hypotheses 140:109767
- 263. Jiang J, Chen Q, Chen X, Li J, Li S, Yang B (2020). Magnesium sulfate ameliorates sepsis-induced diaphragm dysfunction in rats via inhibiting HMGB1/TLR4/NF-κB pathway. Neuroreport 31(12):902-908
- 264. Rani V, Verma Y, Rana K, Rana SVS (2018). Zinc oxide nanoparticles inhibit dimethylnitrosamine induced liver injury in rat. Chem Biol Interact 295:84-92
- 265. Kong W, Yen JH, Vassiliou E, Adhikary S, Toscano MG, Ganea D (2010). Docosahexaenoic acid prevents dendritic cell maturation and in vitro and in vivo expression of the IL-12 cytokine family. Lipids Health Dis 9:12
- 266. Benson AA, Toh JA, Vernon N, Jariwala SP (2012). The role of vitamin D in the immunopathogenesis of allergic skin diseases. Allergy 67:296–301
- 267. Martínez-Moreno J, Hernandez JC, Urcuqui-Inchima S (2020). Effect of high doses of vitamin D supplementation on dengue virus replication, Toll-like receptor expression, and cytokine profiles on dendritic cells. Mol Cell Biochem 464(1-2):169-180
- 268. Reda R, Abbas AA, Mohammed M, El Fedawy SF, Ghareeb H, El Kabarity RH, et al (2015). The interplay between zinc, vitamin D and, IL-17 in patients with chronic hepatitis C liver disease. J Immunol Res 2015:846348

- 269. Cantorna MT, Snyder L, Lin YD, Yang L (2015). Vitamin D and 1,25(OH)2D regulation of T cells. Nutrients 7(4):3011-21
- 270. Schardey J, Globig AM, Janssen C, Hofmann M, Manegold P, Thimme R, et al (2019). Vitamin D inhibits pro-inflammatory T cell function in patients with inflammatory bowel disease. J Crohns Colitis 13(12):1546-1557
- 271. Muroi M, Tanamoto K (2015). Zinc- and oxidative property-dependent degradation of pro-caspase-1 and NLRP3 by ziram in mouse macrophages. Toxicol Lett 235(3):199-205
- 272. Kong J, Grando SA, Li YC (2006). Regulation of IL-1 family cytokines IL-1alpha, IL-1 receptor antagonist, and IL-18 by 1,25-dihydroxyvitamin D3 in primary keratinocytes. J Immunol 176(6):3780-7
- 273. Yang FX, Hou L, Wen WL, Shen XL, Feng NY, Ma RX, et al (2020). Role of zinc sulphate in immune regulation in Artemisia annua pollen-challenged P815 mastocytoma cells. Immunol Invest 49(6):622-631
- 274. Fletcher P, Hamilton RF, Buford M, Postma B, Pestka JJ, Holian A (2019). Comparing docosahexaenoic acid as a prophylactic treatment for acute and chronic particle-exposed Balb/c mice. L Immunol 202 (1 Suppl):117.6
- 275. Mohammadi-Kordkhayli M, Ahangar-Parvin R, Azizi SV, Nemati M, Shamsizadeh A, Khaksari M, et al (2015). Vitamin D modulates the expression of IL-27 and IL-33 in the central nervous system in experimental autoimmune encephalomyelitis (EAE). Iran J Immunol 12(1):35-49
- 276. Monk JM, Liddle DM, Brown MJ, Zarepoor L, De Boer AA, Ma DW, et al (2016). Anti-inflammatory and anti-chemotactic effects of dietary flaxseed oil on CD8(+) T cell/adipocyte-mediated cross-talk. Mol Nutr Food Res 60(3):621-30
- 277. Han H, Qiu F, Zhao H, Tang H, Li X, Shi D (2017). Dietary flaxseed oil prevents Western-type diet-induced nonalcoholic fatty liver disease in apolipoprotein-E knockout mice. Oxid Med Cell Longev 2017:3256241
- 278. Song J, Jing Z, Hu W, Yu J, Cui X (2017). α-Linolenic acid inhibits receptor activator of NF-κB ligand induced (RANKL-induced) osteoclastogenesis and prevents inflammatory bone loss via downregulation of nuclear factorkappaB-inducible nitric oxide synthases (NF-κB-iNOS) signaling pathways. Med Sci Monit 23:5056-5069
- 279. Spigoni V, Lombardi C, Cito M, Picconi A, Ridolfi V, Andreoli R, et al (2014). N-3 PUFA increase bioavailability and function of endothelial progenitor cells. Food Func 5(8):1881-90
- 280. Zhang J, McCullough PA, Tecson KM (2020). Vitamin D deficiency in association with endothelial dysfunction: Implications for patients with COVID-19. Rev Cardiovasc Med 21(3):339-344
- 281. Thota C, Farmer T, Garfield RE, Menon R, Al-Hendy A (2013). Vitamin D elicits anti-inflammatory response, inhibits contractile-associated proteins, and modulates Toll-like receptors in human myometrial cells. Reprod Sci 20(4):463-75
- 282. Sanchez-Niño MD, Bozic M, Córdoba-Lanús E, Valcheva P, Gracia O, Ibarz M, et al (2012). Beyond proteinuria: VDR activation reduces renal inflammation in experimental diabetic nephropathy. Am J Physiol Renal Physiol 302(6):F647-57
- 283. Beloosesky R, Khatib N, Ginsberg Y, Anabosy S, Shalom-Paz E, Dahis M, et al (2016). Maternal magnesium sulfate fetal neuroprotective effects to the fetus: inhibition of neuronal nitric oxide synthase and nuclear factor kappa-

light-chain-enhancer of activated B cells activation in a rodent model. Am J Obstet Gynecol 215(3):382.e1-6

- 284. Slinko S, Piraino G, Hake PW, Ledford JR, O'Connor M, Lahni P, et al (2014). Combined zinc supplementation with proinsulin C-peptide treatment decreases the inflammatory response and mortality in murine polymicrobial sepsis. Shock 41(4):292-300
- 285. Lund AS, Hasselbalch AL, Gamborg M, Skogstrand K, Hougaard DM, Heitmann BL, et al (2013). N-3 polyunsaturated fatty acids, body fat and inflammation. Obes Facts 6(4):369-79
- 286. Boontanrart M, Hall SD, Spanier JA, Hayes CE, Olson JK (2016). Vitamin D3 alters microglia immune activation by an IL-10 dependent SOCS3 mechanism. J Neuroimmunol 292:126-36
- 287. Wang LJ, Wang MQ, Hu R, Yang Y, Huang YS, Xian SX, et al (2017). Effect of zinc supplementation on maintenance hemodialysis patients: A systematic review and meta-analysis of 15 randomized controlled trials. Biomed Res Int 2017:1024769
- 288. Khorsandi H, Nikpayam O, Yousefi R, Parandoosh M, Hosseinzadeh N, Saidpour A, et al (2019). Zinc supplementation improves body weight management, inflammatory biomarkers and insulin resistance in individuals with obesity: a randomized, placebo-controlled, double-blind trial. Diabetol Metab Syndr 11:101
- 289. Hajji M, Khedher R, Mrad M, Bassem HM, Rafrafi N, Chouchi S, et al (2021). Effects of zinc supplementation on serum copper to zinc and CRP to albumin ratios in hemodialysis patients. J Med Biochem 40(2):193-198
- 290. Thota RN, Rosato JI, Burrows TL, Dias CB, Abbott KA, Martins RN, et al (2020). Docosahexaenoic acid-rich fish oil supplementation reduces kinase associated with insulin resistance in overweight and obese midlife adults. Nutrients 12(6):1612
- 291. Kasemy ZA, Hathout HM, Omar ZA, Samir MA, Bahbah WA (2020). Effect of Omega-3 supplements on quality of life among children on dialysis: A prospective cohort study. Medicine (Baltimore) 99(40):e22240
- 292. AbuMweis S, Abu Omran D, Al-Shami I, Jew S (2021). The ratio of eicosapentaenoic acid to docosahexaenoic acid as a modulator for the cardio-metabolic effects of omega-3 supplements: A meta-regression of randomized clinical trials. Complement Ther Med 57:102662
- 293. Cheshmazar E, Hosseini AF, Yazdani B, Razmpoosh E, Zarrati M (2020). Effects of vitamin D supplementation on omentin-1 and spexin levels, inflammatory parameters, lipid profile, and anthropometric indices in obese and overweight adults with vitamin D deficiency under low-calorie diet. Evid Based Complement Alternat Med 2020:3826237
- 294. Zhao JF, Li BX, Zhang Q (2021). Vitamin D improves levels of hormonal, oxidative stress and inflammatory parameters in polycystic ovary syndrome: a meta-analysis study. Ann Palliat Med 10(1):169-183
- 295. Mazidi M, Rezaie P, Banach M (2018). Effect of magnesium supplements on serum C-reactive protein: a systematic review and meta-analysis. Arch Med Sci 14(4):707-716
- 296. Aster I, Barth LM, Rink L, Wessels I (2019). Alterations in membrane fluidity are involved in inhibition of GM-CSF-induced signaling in myeloid cells by zinc. J Trace Elem Med Biol 54:214-220
- 297. Jensen KN, Omarsdottir SY, Reinhardsdottir MS, Hardardottir I, Freysdottir J (2020). Docosahexaenoic acid modulates NK cell effects on neutrophils and their crosstalk. Front Immunol 11:570380

- 298. Ibrahim A, Mbodji K, Hassan A, Aziz M, Boukhettala N, Coëffier M, et al (2011). Anti-inflammatory and anti-angiogenic effect of long chain n-3 polyunsaturated fatty acids in intestinal microvascular endothelium. Clin Nutr 30(5):678-87
- 299. Yusupov E, Li-Ng M, Pollack S, Yeh JK, Mikhail M, Aloia JF (2010). Vitamin D and serum cytokines in a randomized clinical trial. Int J Endocrinol 2010:305054
- 300. Konya V, Czarnewski P, Forkel M, Rao A, Kokkinou E, Villablanca EJ, et al (2018). Vitamin D downregulates the IL-23 receptor pathway in human mucosal group 3 innate lymphoid cells. J Allergy Clin Immunol 141(1):279-292
- 301. Towers TL, Staeva TP, Freedman LP (1999). A two-hit mechanism for vitamin D3-mediated transcriptional repression of the granulocyte-macrophage colony-stimulating factor gene: vitamin D receptor competes for DNA binding with NFAT1 and stabilizes c-Jun. Mol Cell Biol 19(6):4191-9
- 302. Tobler A, Gasson J, Reichel H, Norman AW, Koeffler HP (1987). Granulocyte-macrophage colony-stimulating factor. Sensitive and receptormediated regulation by 1,25-dihydroxyvitamin D3 in normal human peripheral blood lymphocytes. J Clin Invest 79(6):1700-5
- 303. Li G, Fan Y, Lai Y, Han T, Li Z, Zhou P, et al (2020). Coronavirus infections and immune responses. J Med Virol 92(4):424-432
- 304. Li X, Geng M, Peng Y, Meng L, Lu S (2020). Molecular immune pathogenesis and diagnosis of COVID-19. J Pharm Anal 10(2):102-108
- 305. Ferrucci L, Fabbri E (2018). Inflammageing: chronic inflammation in ageing, cardiovascular disease, and frailty. Nat Rev Cardiol 15(9):505-522
- 306. Fiorino S, Gallo C, Zippi M, Sabbatani S, Manfredi R, Moretti R, et al (2020). Cytokine storm in aged people with CoV-2: possible role of vitamins as therapy or preventive strategy. Aging Clin Exp Res 32(10):2115-2131
- 307. Marazuela M, Giustina A, Puig-Domingo M (2020). Endocrine and metabolic aspects of the COVID-19 pandemic. Rev Endocr Metab Disord 21(4):495-507
- 308. Rai V, Agrawal DK (2017). Role of vitamin D in cardiovascular diseases. Endocrinol Metab Clin North Am 46(4):1039–1059
- 309. Williams JW, Huang LH, Randolph GJ (2019). Cytokine circuits in cardiovascular disease. Immunity 50(4):941-954
- 310. Koelman L, Pivovarova-Ramich O, Pfeiffer AFH, Grune T, Aleksandrova K (2019). Cytokines for evaluation of chronic inflammatory status in ageing research: Reliability and phenotypic characterisation. Immun Ageing 16:11
- 311. Janssen CI, Kiliaan AJ (2014). Long-chain polyunsaturated fatty acids (LCPUFA) from genesis to senescence: The influence of LCPUFA on neural development, aging, and neurodegeneration. Prog Lipid Res 53:1-17
- 312. Barbagallo M, Belvedere M, Dominguez LJ (2009). Magnesium homeostasis and aging. Magnes Res 22(4):235-46
- 313. Morgante G, Troìa L, De Leo V (2020). Coronavirus Disease 2019 (SARS-CoV-2) and polycystic ovarian disease: Is there a higher risk for these women? J Steroid Biochem Mol Biol 205:105770
- 314. Wang EW, Siu PM, Pang MY, Woo J, Collins AR, Benzie IFF (2017). Vitamin D deficiency, oxidative stress and antioxidant status: only weak association seen in the absence of advanced age, obesity or pre-existing disease. Br J Nutr 118(1):11–16

- 315. Olechnowicz J, Tinkov A, Skalny A, Suliburska J (2018). Zinc status is associated with inflammation, oxidative stress, lipid, and glucose metabolism. J Physiol Sci 68(1):19-31
- 316. Fukunaka A, Fujitani Y (2018). Role of zinc homeostasis in the pathogenesis of diabetes and obesity. Int J Mol Sci 19(2):476
- 317. Mossink JP (2020). Zinc as nutritional intervention and prevention measure for COVID-19 disease. BMJ Nutr Prev Health 3(1):111-117
- 318. Chabosseau P, Rutter GA (2016). Zinc and diabetes. Arch Biochem Biophys 611:79-85
- 319. Simopoulos AP (2016). An Increase in the Omega-6/Omega-3 Fatty Acid Ratio Increases the Risk for Obesity. Nutrients 8(3):128
- 320. Torrinhas RS, Calder PC, Waitzberg DL (2020). Response to Bistrian BR. Parenteral fish-oil emulsions in critically ill COVID-19 emulsions. J Parenter Enteral Nutr 44(7):1169-1170
- 321. Rafiq S, Jeppesen PB (2018). Body Mass Index, Vitamin D, and Type 2 Diabetes: A Systematic Review and Meta-Analysis. Nutrients 10(9):1182
- 322. Walsh JS, Bowles S, Evans AL (2017). Vitamin D in obesity. Curr Opin Endocrinol Diabetes Obes 24(6):389–394
- 323. Hyppönen E, Boucher BJ (2018). Adiposity, vitamin D requirements, and clinical implications for obesity-related metabolic abnormalities. Nutr Rev 76(9):678–692
- 324. Yao Y, Zhu L, He L, Duan Y, Liang W, Nie Z, et al (2015). A meta-analysis of the relationship between vitamin D deficiency and obesity. Int J Clin Exp Med 8(9):14977-84
- 325. Li X, Liu Y, Zheng Y, Wang P, Zhang Y (2018). The effect of vitamin D supplementation on glycemic control in type 2 diabetes patients: A systematic review and meta-analysis. Nutrients 10(3):375
- 326. Mauss D, Jarczok MN, Hoffmann K, Thomas GN, Fischer JE (2015). Association of vitamin D levels with type 2 diabetes in older working adults. Int J Med Sci 12(5):362-8
- 327. Nielsen FH (2010). Magnesium, inflammation, and obesity in chronic disease. Nutr Rev 68(6):333-40
- 328. Madjid M, Safavi-Naeini P, Solomon SD, Vardeny O (2020). Potential effects of coronaviruses on the cardiovascular system: A review. JAMA Cardiol 5(7):831-840
- 329. Radenkovic D, Chawla S, Pirro M, Sahebkar A, Banach M (2020). Cholesterol in relation to COVID-19: Should we care about it? J Clin Med 9(6):1909
- 330. Choi S, Cui C, Luo Y, Kim SH, Ko JK, Huo X, et al (2018). Selective inhibitory effects of zinc on cell proliferation in esophageal squamous cell carcinoma through Orai1. FASEB J 32(1):404-416
- 331. Jurowski K, Szewczyk B, Nowak G, Piekoszewski W (2014). Biological consequences of zinc deficiency in the pathomechanisms of selected diseases. J Biol Inorg Chem 19(7):1069-79
- 332. Calder PC (2017). Omega-3 fatty acids and inflammatory processes: From molecules to man. Biochem Soc Trans 45(5):1105-1115
- 333. Calder PC (2013). Omega-3 polyunsaturated fatty acids and inflammatory processes: nutrition or pharmacology?. Br J Clin Pharmacol 75(3):645-62
- 334. Darwesh AM, Sosnowski DK, Lee TY, Keshavarz-Bahaghighat H, Seubert JM (2019). Insights into the cardioprotective properties of n-3 PUFAs against ischemic heart disease via modulation of the innate immune system. Chem Biol Interact 308:20-44

- 335. Kris-Etherton PM, Harris WS, Appel LJ; Nutrition Committee (2003). Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. Arterioscler Thromb Vasc Biol 23(2):e20-30
- 336. Lee JH, O'Keefe JH, Lavie CJ, Harris WS (2009). Omega-3 fatty acids: Cardiovascular benefits, sources and sustainability. Nat Rev Cardiol 6(12):753-8
- 337. Mozaffarian D, Lemaitre RN, King IB, Song X, Huang H, Sacks FM, Rimm EB, Wang M, Siscovick DS (2013). Plasma phospholipid long-chain ω -3 fatty acids and total and cause-specific mortality in older adults: A cohort study. Ann Intern Med 158(7):515-25
- 338. Darwesh AM, Bassiouni W, Sosnowski DK, Seubert JM (2020). Can N-3 polyunsaturated fatty acids be considered a potential adjuvant therapy for COVID-19-associated cardiovascular complications? Pharmacol Ther 219:107703
- 339. Paschou SA, Kosmopoulos M, Nikas IP, Spartalis M, Kassi E, Goulis DG, et al (2019). The impact of obesity on the association between vitamin D deficiency and cardiovascular disease. Nutrients 11(10):2458
- 340. Porto CM, Silva VL, da Luz JSB, Filho BM, da Silveira VM (2018). Association between vitamin D deficiency and heart failure risk in the elderly. ESC Heart Fail 5(1):63–74
- 341. Hughes DA, Norton R (2009). Vitamin D and respiratory health. Clin Exp Immunol 158(1):20-5
- 342. Liu LC, Voors AA, van Veldhuisen DJ, van der Veer E, Belonje AM, Szymanski MK, et al (2011). Vitamin D status and outcomes in heart failure patients. Eur J Heart Fail 13(6):619-25
- 343. Cotogni P, Trombetta A, Muzio G, Maggiora M, Canuto RA (2015). The omega-3 fatty acid docosahexaenoic acid modulates inflammatory mediator release in human alveolar cells exposed to bronchoalveolar lavage fluid of ARDS patients. Biomed Res Int 2015:642520
- 344. Mekov E, Slavova Y, Tsakova A, Genova M, Kostadinov D, Minchev D, et al (2015). Vitamin D deficiency and insufficiency in hospitalized COPD patients. PLoS One 10(6):e0129080
- 345. Martineau AR, Jolliffe DA, Hooper RL, Greenberg L, Aloia JF, Bergman P, et al (2017). Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. BMJ 356:i6583
- 346. Moghaddassi M, Pazoki M, Salimzadeh A, Ramim T, Alipour Z (2018). Association of serum level of 25-hydroxy vitamin D deficiency and pulmonary function in healthy individuals. ScientificWorldJournal 2018:3860921
- 347. Mulrennan S, Knuiman M, Walsh JP, Hui J, Hunter M, Divitini M, et al (2018). Vitamin D and respiratory health in the Busselton Healthy Ageing Study. Respirology 23(6):576-582
- 348. Brenner H, Holleczek B, Schöttker B (2020). Vitamin D insufficiency and deficiency and mortality from respiratory diseases in a cohort of older adults: Potential for limiting the death toll during and beyond the COVID-19 pandemic?. Nutrients 12(8):2488
- 349. Ilyas M, Agussalim A, Megawati M, Massi N, Djaharuddin I, Bakri S, et al (2019). Relationship between vitamin D level and serum TNF-a concentration on the severity of chronic obstructive pulmonary disease. Open Access Maced J Med Sci 7(14):2298-2304

- 350. Patterson WL 3rd, Georgel PT (2014). Breaking the cycle: the role of omega-3 polyunsaturated fatty acids in inflammation-driven cancers. Biochem Cell Biol 92(5):321-8
- 351. Niedermaier T, Gredner T, Kuznia S, Schöttker B, Mons U, Brenner H (2021). Vitamin D supplementation to the older adult population in Germany has the cost-saving potential of preventing almost 30 000 cancer deaths per year. Mol Oncol Ahead of print

Journal Pre-proof

Highlights

Title: Essential sufficiency of zinc, ω -3 polyunsaturated fatty acids, vitamin D and magnesium for prevention and treatment of COVID-19, diabetes, cardiovascular diseases, lung diseases and cancer

- Zinc, $\omega\text{-}3$ PUFAs, vitamin D and magnesium are inexpensive, safe and essential, especially for the aged
- These nutrients inhibit viral entry, viral proliferation, inflammation and immune dysfunction
- These nutrients will assist in preventing and alleviating COVID-19 and associated comorbidities
- Levels of these nutrients should be checked annually and supplemented for sufficiency in the aged
- Costs of checking and provision of supplements should be at government expense for the aged

ournal