

## Original Article

**Cite this article:** Kjeldsen EW, Thomassen JQ, Rasmussen KL, Nordestgaard BG, Tybjaerg-Hansen A, Frikke-Schmidt R (2022). Adherence to dietary guidelines and risk of dementia: a prospective cohort study of 94 184 individuals. *Epidemiology and Psychiatric Sciences* **31**, e71, 1–10. <https://doi.org/10.1017/S2045796022000567>

Received: 12 January 2022

Revised: 15 August 2022

Accepted: 12 September 2022

### Key words:


Ageing; diet; lipids; non-Alzheimer's dementia

### Author for correspondence:

Ruth Frikke-Schmidt,

E-mail: [ruth.frikke-schmidt@regionh.dk](mailto:ruth.frikke-schmidt@regionh.dk)

# Adherence to dietary guidelines and risk of dementia: a prospective cohort study of 94 184 individuals

E. W. Kjeldsen<sup>1,2</sup>, J. Q. Thomassen<sup>1</sup>, K. L. Rasmussen<sup>1,2</sup>, B. G. Nordestgaard<sup>2,3,4</sup>, A. Tybjaerg-Hansen<sup>1,2,3</sup> and R. Frikke-Schmidt<sup>1,2,3</sup> 

<sup>1</sup>Department of Clinical Biochemistry, Copenhagen University Hospital – Rigshospitalet, Blegdamsvej 9, 2100 Copenhagen, Denmark; <sup>2</sup>Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Blegdamsvej 3, 2200 Copenhagen, Denmark; <sup>3</sup>The Copenhagen General Population Study, Copenhagen University Hospital – Herlev and Gentofte, Borgmester Ib Juuls Vej 1, 2730 Herlev, Denmark and <sup>4</sup>Department of Clinical Biochemistry, Copenhagen University Hospital – Herlev and Gentofte, Borgmester Ib Juuls Vej 1, 2730 Herlev, Denmark

## Abstract

**Aims.** Recent estimates suggest that 40% of dementia cases could be avoided by treating recognised cardiovascular risk factors such as hypertension, diabetes, smoking and physical inactivity. Whether diet is associated with dementia remains largely unknown. We tested if low adherence to established dietary guidelines is associated with elevated lipids and lipoproteins and with increased risk of Alzheimer's disease and non-Alzheimer's dementia – a dementia subtype with a high frequency of cardiovascular risk factors.

**Methods.** We used the prospective Copenhagen General Population Study including 94 184 individuals with dietary information and free of dementia at baseline. Mean age at study entry was 58 years, and 55% ( $N = 51\,720$ ) were women and 45% ( $N = 42\,464$ ) were men. Adherence to dietary guidelines was grouped into low, intermediate and high adherence based on food frequency questionnaires. Main outcomes were non-Alzheimer's dementia and Alzheimer's disease.

**Results.** Low-density lipoprotein cholesterol, non-high-density lipoprotein cholesterol and plasma triglyceride levels were higher in individuals with intermediate and low adherence to dietary guidelines compared with individuals with high adherence (all  $p$  for trends  $<0.001$ ). Age and sex-adjusted hazard ratios (HRs) for non-Alzheimer's dementia *v.* individuals with high adherence were 1.19 (95% confidence interval 0.97–1.46) for intermediate adherence, and 1.54 (1.18–2.00) for low adherence. Corresponding HRs in multivariable-adjusted models including *APOE* genotype were 1.14 (0.92–1.40) and 1.35 (1.03–1.79). These relationships were not observed in individuals on lipid-lowering therapy.

**Conclusions.** Low adherence to national dietary guidelines is associated with an atherogenic lipid profile and with increased risk of non-Alzheimer's dementia – the subtype of dementia with a high frequency of vascular risk factors. This study suggests that implementation of dietary guidelines associated with an anti-atherogenic lipid profile could be important for prevention of non-Alzheimer's dementia.

## Introduction

Dementia is a devastating disease currently affecting the activity of daily life in 50 million people (World Health Organization, 2017). Due to ageing of populations worldwide, the number is estimated to grow to 152 million by 2050 (World Health Organization, 2017). No curative treatment or medication to effectively halt dementia exist (World Health Organization, 2017). Therefore, it is of utmost importance to establish and act on modifiable risk factors to implement strategies to prevent or delay dementia onset. Approximately 40% of dementia cases are considered caused by modifiable vascular risk factors such as hypertension, diabetes, smoking and physical inactivity. This leaves a substantial potential for prevention (Livingston *et al.*, 2020), where diet could play a central role in reducing occurrence of vascular risk factors. Although the contribution of diet to death is recognised to be substantial globally (Abbas *et al.*, 2020), the association with dementia remains largely unknown (Livingston *et al.*, 2020).

Recently, the Global Burden of Disease report showed that dietary risk was a major contributor to attributable deaths (Abbas *et al.*, 2020). Additionally diet quality was the fifth leading risk factor for disability due to cardiovascular disease, diabetes, kidney diseases and neoplasms (Abbas *et al.*, 2020). Furthermore, current diets in most countries worldwide are too unhealthy and dietary changes as recommended by national guidelines could lower rates of premature mortality (Springmann *et al.*, 2020). Adherence to a healthy diet could therefore have a promising

© The Author(s), 2022. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

potential for improving public health but also for preventing dementia. A systematic review concluded that a Mediterranean diet may contribute to better cognitive function, but that there is a need for larger epidemiological studies taking more factors such as the presence of comorbidities into account (Petersson and Philippou, 2016). Furthermore, adherence to a healthy diet by mainly using unsaturated fat, eating fruit and plenty of vegetables, eating more fish, choosing whole grain foods and consuming only moderate amounts of sugar and foods high in salt content, is associated with lower risk of the dementia subtype Alzheimer's disease (Féart *et al.*, 2009; Morris *et al.*, 2015; Haring *et al.*, 2016; Van Den Brink *et al.*, 2019). Individuals with vascular and unspecified dementia have a high frequency of cardiovascular risk factors. These diagnoses are often grouped into a combined dementia subtype, called non-Alzheimer's dementia. It is, however, unknown whether diet is associated with this type of dementia (Perez *et al.*, 2012; Wu and Sun, 2017; Akbaraly *et al.*, 2019; Barbaresco *et al.*, 2020; McKenzie *et al.*, 2021). Consequently, it is important to explore the gaps in knowledge and examine the association of adherence to established dietary guidelines with risk of non-Alzheimer's dementia and Alzheimer's disease in a large population-based study with long follow-up.

The aim of this study was to investigate the relationship between adherence to established dietary guidelines and lipids and lipoproteins and risk of Alzheimer's disease and non-Alzheimer's dementia. Adherence to established dietary guidelines was assessed by a simple food frequency questionnaire in a large prospective cohort, The Copenhagen General Population Study (CGPS), including 94 184 individuals aged 20–100 years at baseline.

## Materials and methods

The study was approved by institutional review boards and a Danish Ethical Committee (H-KF-01-144/01) and was conducted according to the Declaration of Helsinki (revision 2000). All individuals were white and of Danish descent. Written informed consent was obtained from all individuals.

### Participants

The CGPS was initiated in 2003–2015 (online Supplementary Figs S1 and S2) with follow-up examinations starting in 2015 (CGPS2) (Jørgensen *et al.*, 2014; Rasmussen *et al.*, 2019; Juul Rasmussen *et al.*, 2020). To reflect the adult Danish population aged 20–100 years, individuals were selected randomly based on the Danish Civil Registration System. Data were obtained at baseline (study inclusion) from a self-administered questionnaire reviewed together with an investigator on the day of attendance, a physical examination and from blood samples including DNA extraction. Information on diet, vital status and disease status was available for 94 184 individuals who were followed for a median of 9 years. A table of baseline characteristics for individuals included and excluded from this study is provided in online Supplementary Table S1. Individuals participating in the study were by linkage to the Danish registries followed with information on diagnosis codes, emigration status and information from the causes of death registry. We excluded 137 individuals with a dementia diagnosis at baseline.

### Dietary assessment and dietary categories

Dietary habits were assessed from a short food frequency questionnaire (FFQ) which was a part of the extensive overall

questionnaire used in CGPS (online Supplementary Table S2). The short FFQ was developed in collaboration with nutrition specialists and included important items based on the Danish dietary guidelines to assess overall dietary habits (Ewers *et al.*, 2021). This kind of categorisation showed strong associations with cardiovascular disease and mortality (Ewers *et al.*, 2021). The FFQ focused on selected key items of the Danish food-based dietary guidelines, i.e. the recommendations to: (1) eat less saturated fat, (2) eat fruit and plenty of vegetables, (3) eat more fish, (4) eat less sugar and (5) eat foods with less salt (The Official Dietary Guidelines – Good for Health and Climate, 2020). The FFQ specifically explored: (1) dietary fat quality in cold and warm meals (saturated fats: butter, butter-based blends and hard margarines; unsaturated fats: soft margarines and vegetable oils); (2) usual intake frequencies (from almost never to several daily servings) of fruit, vegetables, fish, sucrose-sweetened beverages, cold meat cuts like sausages and pâtés for open sandwiches and fast food (Ewers *et al.*, 2021). The FFQ did not cover intake of foods directly related to the remaining recommendations of the Danish food-based dietary guidelines advocating to choose whole grain foods, to choose low-fat dairy products, to eat low-fat meat and meat products or to drink water. Nor did the FFQ include assessment of portion sizes.

For consistency between studies, we used an identical set-up as in a previous study that used the same short FFQ; we classified our FFQ questions into three levels of importance (from A to C) for an overall healthy dietary pattern (Ewers *et al.*, 2021). Class A questions focused on major contributors to the dietary macronutrient composition, specifically dietary fat quality (fats in cold and warm meals: unsaturated *v.* saturated fat) and dietary fibre content (fruit:  $\geq 3$  *v.*  $< 3$  weekly servings and vegetables:  $\geq 3$  *v.*  $< 3$  weekly servings). Class B questions elucidated intake of specific foods considered healthy (fish:  $\geq 3$  *v.*  $< 1$  weekly servings), or unhealthy (sugar sweetened beverages:  $< 0.5$  *v.*  $\geq 1$  l/week). Class C questions focused on foods rich in salt (cold meat cuts like sausages and pâtés for open sandwiches:  $< 5$  *v.*  $\geq 7$  weekly servings and fast foods:  $< 1$  *v.*  $\geq 1$  weekly servings) (Ewers *et al.*, 2021).

Based on the FFQ, individuals were divided into three predefined categories ranging from high to low adherence to current dietary guidelines. High adherence: all class A, B and C answers in agreement with current guidelines; or all A, B and C answers in agreement with current guidelines except for either one class B answer in disagreement with guidelines or one or two class C answers in disagreement with guidelines. Intermediate adherence: individuals between high and low adherence categories. Low adherence: three or four class A answers in disagreement with current guidelines (Fig. 1 and online Supplementary Table S3).

We included individuals participating in both CGPS and CGPS2 ( $n = 17\ 506$ ) in an analysis to estimate changes in dietary patterns from baseline to second examination follow-up. A total of 12 800 individuals filled out FFQs at both CGPS and CGPS2. The FFQ included the same questions at both visits. The median time interval between first examination and follow-up was 10 years.


Description of clinical endpoints, laboratory analyses, genotyping, covariates and statistical analyses is provided in the online Supplementary materials, Methods.

## Results

Baseline characteristics of 94 184 individuals stratified by three groups of adherence to dietary guidelines are shown in Table 1.

## Classification of dietary assessment

Question Priority	Dietary food component	In agreement with guidelines	In disagreement with guidelines
A	Fats for cold meals	Unsaturated	Saturated
A	Fats for warm meals	Unsaturated	Saturated
A	Vegetables, weekly servings	≥3	≤2
A	Fruit, weekly servings	≥3	≤2
B	Fish, weekly servings	≥3	≤1
B	Sugar-sweetened beverages, L/week	<0.5	≥1
C	Cold meat cuts, weekly servings	<5	≥7
C	Fast food, weekly servings	<1	≥1



## Categorization of adherence to dietary guidelines

High	Intermediate	Low
All class A, B and C in agreement with guidelines.  All class A in agreement AND one class B OR one to two class C in disagreement with guidelines.	Individuals between high and low adherence.	Three or four class A in disagreement with guidelines.

**Fig. 1.** Dietary assessment in 94 184 individuals from the CGPS. FFQ questions were classified into three levels of importance (from A to C) according to an overall healthy dietary pattern. Based on the FFQ, individuals were divided into three predefined categories ranging from high to low adherence to current dietary guidelines. See 'Methods' for further details. FFQ, Food Frequency Questionnaire.

**Table 1.** Baseline characteristics of 94 184 individuals grouped according to degree of adherence to Danish dietary guidelines

Baseline characteristics ( <i>N</i> = 94 184)	Adherence to dietary guidelines		
	High	Intermediate	Low
Number, no. (%)	19 679 (20.9)	65 903 (70.0)	8602 (9.0)
Age, years (median, IQR)	59.6 (50.8–67.4)	57.3 (47.4–67.0)	60.1 (49.0–70.2)
Men, no. (%)	6592 (33.5)	30 222 (45.9)	5650 (65.7)
Low household income, no. (%)	6177 (31.4)	22 706 (34.5)	4483 (52.1)
Education <8 years, no. (%)	1431 (7.3)	5839 (8.9)	1506 (17.5)
Body mass index, kg/m <sup>2</sup> (median, IQR)	25.4 (23.1–28.1)	25.5 (23.2–28.4)	(23.5–29.1)
Physical inactivity in leisure time, no. (%)	7925 (40.3)	31 732 (48.2)	5542 (64.4)
Cumulative tobacco consumption, pack-years (median, IQR)	12 (5–24)	15 (6–30)	29 (15–44)
High alcohol consumption, no. (%)	3215 (16.3)	11 055 (16.8)	1945 (22.6)
Hypertension, no. (%)	11 917 (60.6)	39 161 (59.4)	5669 (65.9)
Diabetes mellitus, no. (%)	935 (4.8)	2592 (3.9)	409 (4.8)
Lipid-lowering therapy, no. (%)	2999 (15.2)	7477 (11.4)	1009 (11.7)

IQR, interquartile range.

Low education corresponds to completion of primary school or less. See Methods (covariates) in the online Supplementary materials for details.

The mean age of participants at study inclusion was 58. Fifty-five per cent (51 720/94 184) were women and 45% (42 464/94 184) were men. Analyses were conducted in men and women combined as no interaction between sex and exposure in the association with dementia was observed ( $p$ -value = 0.12). Twenty-one per cent of individuals had high adherence to dietary guidelines, 70% had intermediate adherence and 9% had low adherence. Individuals in the group with low adherence to dietary guidelines were more frequently men, had shorter education, lower household income, higher body mass index, were less physically active, had a higher tobacco and alcohol consumption, had more frequently hypertension, and treatment with lipid-lowering therapy was less frequent.

### Changes in diet over time

The frequency of individuals who did not change their diet was 8382 individuals out of 12 800 (65%). A total of 4336 (34%) individuals changed their diet to either one group higher or lower in adherence to dietary guidelines, whereas only 72 individuals (0.6%) changed two groups: from high to low or from low to high adherence to dietary guidelines. The specific number of individuals for each diet group either increasing or decreasing their adherence to dietary guidelines is detailed in online Supplementary Table S5. The distribution of individuals in the three dietary categories was largely similar at baseline and at follow-up indicating no major changes in overall diets.

### Lipids and lipoproteins

Levels of low-density lipoprotein (LDL) cholesterol, non-high-density lipoprotein (HDL) cholesterol and plasma triglycerides increased stepwise for individuals from high over intermediate to low adherence to dietary guidelines, while HDL cholesterol levels decreased ( $p$ -values ranging from  $8 \times 10^{-227}$  to  $3 \times 10^{-8}$ ) (Fig. 2).

### Risk of dementia

Age and sex-adjusted hazard ratios (HRs) for non-Alzheimer's dementia increased stepwise with lower adherence to dietary guidelines (Fig. 3). Compared with individuals with high adherence, age and sex-adjusted HRs were 1.19 (95% confidence interval 0.97–1.46) for individuals with intermediate adherence and 1.54 (1.18–2.00) for individuals with low adherence. Corresponding multivariable-adjusted HRs were 1.15 (0.94–1.41) and 1.35 (1.03–1.78) while corresponding HRs when further adjusting for *APOE* genotype were 1.14 (0.92–1.40) and 1.35 (1.03–1.79). In contrast, adherence to dietary guidelines was not associated with risk of Alzheimer's disease. Results were similar for both non-Alzheimer's dementia and Alzheimer's disease in a sensitivity analysis where individuals with less than 2 years of follow-up were excluded (online Supplementary Fig. S3). Furthermore, when additionally adjusting for ischaemic heart disease and ischaemic cerebrovascular disease, or when using imputed variables based on all covariates and end-points, results were similar (online Supplementary Figs S4 and S5).


### Cardiovascular and all-cause mortality as positive controls

Age and sex-adjusted HRs for cardiovascular mortality *v.* individuals with high adherence to dietary guidelines were 1.21 (1.07–1.38) for intermediate and 1.95 (1.67–2.28) for low adherence (online Supplementary Fig. S6). Corresponding HRs for all-cause mortality were 1.26 (1.19–1.33) and 1.97 (1.83–2.12), respectively. Results were similar after multivariable adjustment.

### Interaction with lipid-lowering therapy

The  $p$ -value for interaction between dietary groups and lipid-lowering therapy on risk of non-Alzheimer's dementia was 0.04. No other interactions were detected for the remaining covariates (all  $p > 0.05$ ) (online Supplementary Fig. S7). Consequently, we performed an analysis stratified by lipid-lowering therapy (Fig. 4). For individuals who were not treated with lipid-lowering





**Fig. 2.** Plasma levels of lipids and lipoproteins as a function of adherence to dietary guidelines in 94 184 individuals from the CGPS. Geometric mean  $\pm$  standard errors of the mean are given for triglycerides; arithmetic mean  $\pm$  standard errors of the mean are given for LDL cholesterol, HDL cholesterol and non-HDL cholesterol. LDL cholesterol, low-density lipoprotein cholesterol; HDL cholesterol, high-density lipoprotein cholesterol.

therapy multivariable-adjusted HRs for non-Alzheimer's dementia were 1.39 (1.07–1.81) in individuals with intermediate adherence and 1.63 (1.18–2.25) in individuals with low adherence to

dietary guidelines. Corresponding HRs for individuals treated with lipid-lowering therapy were 0.78 (0.55–1.11) and 0.90 (0.50–1.60), respectively.


## Discussion

In this prospective cohort study of 94 184 individuals from the Danish general population, low adherence to national dietary guidelines was associated with an atherogenic plasma lipid profile and with increased risk of non-Alzheimer's dementia, an association that did not appear to be present in individuals treated with lipid-lowering therapy. These findings are novel and may be clinically important as they suggest that non-Alzheimer's dementia can potentially be prevented by well-established interventions such as adhering to national dietary guidelines and by lipid-lowering treatment.

To our knowledge this is the first study to show an association between low adherence to dietary guidelines and increased risk of non-Alzheimer's dementia in Denmark as well as globally. Previous cross-sectional or prospective studies investigating the association between diet and risk of dementia have primarily observed an association between a healthy diet and less cognitive decline or a lower risk of Alzheimer's disease (Féart *et al.*, 2009; Morris *et al.*, 2015; Haring *et al.*, 2016; Van Den Brink *et al.*, 2019). These studies often focused on single nutrients or foods, did not adjust for all known risk factors and did not take *APOE* genotype into account (Cao *et al.*, 2016; Van Den Brink *et al.*, 2019; Barbaresko *et al.*, 2020; Limongi *et al.*, 2020; Zhang *et al.*, 2021). Evidence of an association with all-cause dementia has been inconsistent (Wu and Sun, 2017; Akbaraly *et al.*, 2019; Van Den Brink *et al.*, 2019; McKenzie *et al.*, 2021), and findings on vascular dementia are sparse (Perez *et al.*, 2012). Randomised controlled trials including dietary interventions show results in favour of a healthy diet with higher cognition, however, with no measures on hard endpoints such as Alzheimer's disease or vascular dementia (Martínez-Lapiscina *et al.*, 2013; Valls-Pedret *et al.*, 2015). Non-Alzheimer's dementia is a dementia subtype associated with cardiovascular risk factors; the present findings on low adherence to dietary guidelines and non-Alzheimer's dementia firmly support that diet may play an important role in vascular health including cerebrovascular health and dementia with vascular pathology. The World Health Organization states that a 'healthy and balanced diet should be recommended to all adults', as it has the potential to prevent cognitive decline both directly and through other potential risk factors or diseases (WHO Guidelines, 2019).

During the past few years dietary research has changed from focusing on the role of single nutrients or food items to the role of dietary patterns like the Mediterranean diet, the Dietary Approaches to Stop Hypertension (DASH) or the Mediterranean-DASH Intervention for Neurodegenerative Delay (MIND) diet (Pistollato *et al.*, 2018). Previous studies have frequently investigated these specific diets and their associations with dementia, which are all characterised by a high intake of vegetables, fruits and unsaturated fats with modest amounts of alcohol and animal foods. In general, observational studies have found that these diets are associated with lower risk of Alzheimer's disease and less cognitive decline (Van Den Brink *et al.*, 2019). The current study did not use one of these established methods to characterise dietary patterns, however, the dietary assessment focused on very similar aspects with only minor discrepancies.


Dietary risk is one of the leading causes of attributable deaths in the Global Burden of Disease 2019 report, and diet quality is



**Fig. 3.** Risk of non-Alzheimer's dementia and Alzheimer's disease according to dietary groups in 94 184 individuals from the CGPS. Multivariable adjustment was for age, sex, household income, education, pack-years, alcohol consumption, physical activity, body mass index, diabetes, hypertension, lipid-lowering therapy, LDL cholesterol, HDL cholesterol and triglycerides. *APOE* genotype, apolipoprotein E  $\epsilon 2/\epsilon 3/\epsilon 4$  genotype; CI, confidence interval; HDL cholesterol, high-density lipoprotein cholesterol; LDL cholesterol, low-density lipoprotein cholesterol.

the fifth leading risk factor for disability mainly due to cardiovascular disease, diabetes, kidney diseases and neoplasms (Abbatati *et al.*, 2020). In addition, our study suggests an increased risk of non-Alzheimer's dementia by non-adherence to dietary guidelines. Interestingly, a study published in 2020 found that current diets in most countries worldwide are too unhealthy and that national dietary guidelines should be reformed and better implemented (Springmann *et al.*, 2020). The study showed that dietary changes as those recommended by dietary guidelines could be associated with substantially reduced premature mortality (Springmann *et al.*, 2020). This would be by following the Mediterranean diet, the DASH, the MIND diet or more specifically by increasing consumption of healthy foods such as fruits, vegetables and legumes; reducing consumption of unhealthy foods, such as sugar, unrefined grains, salt, highly processed foods and red meat and choosing unsaturated fats rather than saturated fats (Willett *et al.*, 2019). There are numerous ways of

implementing food-based dietary guidelines e.g. by updating policies to incentivise better adherence to dietary guidelines for instance by investment in health promotion campaigns and programmes, regulation of agricultural strategies, changes in consumer behaviour, educating families on healthy diets and taxation of unhealthy food items (Sassi *et al.*, 2018; Willett *et al.*, 2019; Springmann *et al.*, 2020). Furthermore, as this study shows a clear clustering of unfavourable cardiovascular risk factors in individuals with low adherence to dietary guidelines the potential to target health strategies towards specific subpopulations is evident. Moreover, updating national food-based dietary guidelines could be beneficial for meeting global sustainability goals (Springmann *et al.*, 2020) in line with the EAT Lancet Commission and the United Nations Sustainable Development Goals 2030 Agenda (United Nations; Willett *et al.*, 2019). Adopting a sustainable diet rich in plant-based foods, low in animal source foods and highly processed foods would not only



**Fig. 4.** Risk of non-Alzheimer's dementia according to dietary groups stratified by lipid-lowering therapy in 94 184 individuals from the CGPS. Multivariable adjustment was for age, sex, household income, education, pack-years, alcohol consumption, physical activity, body mass index, diabetes, hypertension, LDL cholesterol, HDL cholesterol and triglycerides. *APOE* genotype, apolipoprotein E  $\epsilon 2/\epsilon 3/\epsilon 4$  genotype; CI, confidence interval; HDL cholesterol, high-density lipoprotein cholesterol; LDL cholesterol, low-density lipoprotein cholesterol.

benefit human health but also the environment (Willett *et al.*, 2019). A huge potential for prevention exists, unfortunately, it is not currently met. More attention towards effective dietary strategies to reduce disease and mortality risk is imperative.

A plausible mechanism for our findings could be that non-Alzheimer's dementia and atherosclerotic cardiovascular disease share pathogenesis through atherosclerosis. In this study, individuals with low adherence to dietary guidelines were characterised by an atherogenic lipid profile with high levels of LDL cholesterol, non-HDL cholesterol and plasma triglycerides as a marker of triglyceride-rich lipoproteins, all causally related to atherosclerotic cardiovascular disease (Nordestgaard and Varbo, 2014; Ference *et al.*, 2017; Mach *et al.*, 2020). This lipid profile is most probably caused by primarily using saturated fat, eating more fast food, eating more sweets and fewer fruits and vegetables. Consistent evidence from genetic studies, prospective cohort studies, Mendelian randomisation studies and randomised controlled trials demonstrate that LDL causes atherosclerotic cardiovascular disease (Ference *et al.*, 2017). LDL is the most abundant atherogenic lipoprotein in plasma and a key deliverer of cholesterol to the arterial wall (Borén *et al.*, 2020). The likelihood of LDLs being kept inside the arterial intima leading to the establishment of atherosclerotic plaques advances when concentrations of circulating LDL particles increase (Skälén *et al.*, 2002). Retention and subsequent modification of the LDL particle can elicit inflammatory processes and oxidative changes which influence the rate of plaque growth and the tendency to plaque disruption (Borén *et al.*, 2020). Furthermore, due to their cholesterol content, triglyceride-rich lipoproteins are causally associated with atherosclerotic cardiovascular disease, as shown repeatedly in observational and genetic studies (Nordestgaard and Varbo, 2014; Hegele *et al.*, 2020). When triglyceride concentrations are mild to moderately elevated (2–10 mmol/L), triglyceride-rich lipoproteins are sufficiently small to go through the arterial wall and therefore prone to preferential entrapment, and consequently accumulation causing atherosclerosis (Nordestgaard and Varbo, 2014). Our findings are supported by a recent report showing an association between moderate hypertriglyceridaemia and increased risk of non-Alzheimer's dementia in the general population (Nordestgaard *et al.*, 2021). Importantly, in the present study we also show that the increasingly higher risk of non-Alzheimer's dementia with non-adherence to dietary guidelines was not observed in individuals on lipid-lowering therapy, indirectly suggesting that statins potentially may be used to prevent non-Alzheimer's dementia. Due to the observational nature of our study, these findings need to be tested in a randomised controlled trial before any conclusion on treatment benefits can be drawn. Other possible mechanisms for the present findings include the hypothesised effects of oxidative stress and neuroinflammation on risk of dementia (Kinney *et al.*, 2018; Luo *et al.*, 2020). Several risk factors including cardiovascular factors and metabolic diseases such as diabetes, often caused by unhealthy diets, are correlated with an immune response, also in the brain (Kinney *et al.*, 2018). Systemic inflammation can lead to damage of the blood-brain barrier (BBB) which allows entry of immune cells into the brain. Moreover, impaired integrity of the BBB and thus increased BBB permeability can lead to excessive activation of microglial cells, which as a result release proinflammatory and reactive oxygen species.

Limitations of our study include the assessment of diet which was self-reported. Diets often change over time secondary to medications, influences from surroundings or diagnosis of a medical

condition. However, when we analysed a subset of individuals who both participated at baseline and at follow-up examinations and fulfilled the FFQ at both visits, we found that the majority of individuals remained in the same group of adherence to dietary guidelines. Only a small fraction (<1%) changed dietary adherence group from either high to low or low to high. This indicates that our dietary instrument is a valid exposure over time. The FFQ did not take all national dietary advice into account, as for instance intake of dietary fibre or portion sizes, which therefore prevents us from investigating the impacts of macronutrients or adjusting for energy intake. However, the included questions have previously shown to be a sufficient proxy for an individual's dietary habits (Ewers *et al.*, 2021). Furthermore, a previous Danish study compared an extensive FFQ of 198 items with a short FFQ, like the one in this study (Toft *et al.*, 2007). They found that the short FFQ reflected dietary quality adequately and could be used as a tool to classify dietary patterns. Whether a change in adherence to dietary guidelines either to the better or worse was associated with Alzheimer's disease or non-Alzheimer's dementia could not be determined due to a limited number of cases in the subset. This important issue warrants further investigation in future prospective studies with repeated assessment of diet during follow-up. Non-response bias cannot be excluded, as individuals who did not respond to the FFQ, and thus excluded from this study, were less healthy and had lower socioeconomic status than those who did respond. This could limit the generalisability of the study. Study participants were white and from an ethnically homogenous population, which may also limit the generalisability of our findings. However, since the Danish national dietary guidelines are similar to guidelines of other countries, we would expect that our findings apply to other populations (Wilson *et al.*, 2019; Mach *et al.*, 2020). Finally, our findings for those treated with lipid-lowering therapy need to be confirmed in independent studies, preferably in randomised controlled trials. Strengths of the present study are addressed in the online Supplementary materials.

Low adherence to national dietary guidelines is associated with an atherogenic lipid profile and with an increased risk of non-Alzheimer's dementia – the subtype of dementia with a high frequency of vascular risk factors. The present study suggests that implementation of national dietary guidelines associated with an anti-atherogenic lipid profile could be of importance for prevention of non-Alzheimer's dementia and for improved cognitive health in the general population.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S2045796022000567>

**Acknowledgements.** We thank staff and participants in the Copenhagen General Population Study.

**Author contributions.** EWK and RF-S: study concept and design, validation of underlying data, drafting the manuscript. EWK, BGN, AT-H and RF-S: acquisition of data. EWK, JQT, KLR and RF-S: statistical analysis, and analysis and interpretation of data. BGN, AT-H and RF-S: obtained funding, and administrative, technical and material support. All authors: critical revision of the manuscript, important intellectual content and final approval for submission. RF-S: study supervision and accountable for all aspects of the work.

**Financial support.** This work was supported by the Lundbeck Foundation (grant no. R278-2018-804) and the Danish Heart Foundation.

**Conflict of interest.** BGN reports consultancies and talks sponsored by AstraZeneca, Sanofi, Regeneron, Akcea, Amgen, Kowa, Denka Seiken,



Amarin, Novartis, Novo Nordisk, Esperion and Silence Therapeutics. Other authors declare that they have no conflict of interest.

**Ethical standards.** The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2000.

**Availability of data and materials.** Danish Law does not allow transfer of these data. Upon reasonable request to the corresponding author, the steering committee of the CGPS will evaluate whether data access through direct collaboration can be granted.

## References

Abbatfati C, Abbas KM, Abbasi-Kangevari M, Abd-Allah F, Abdelalim A, Abdollahi M, Abdollahpour I, Abegaz KH, Abolhassani H, Abovans V, Abreu LG, Abrigo MRM, Abualhasan A, Abu-Raddad LJ, Abushouk AI, Adabi M, Adekanmbi V, Adeoye AM, Adetokunboh OO, Adham D, Advani SM, Afshin A, Agarwal G, Aghamir SMK, Agrawal A, Ahmad T, Ahmadi K, Ahmadi M, Ahmadi H, Ahmed MB, Akalu TY, Akinyemi RO, Akinyemiju T, Akombi B, Akunna CJ, Alahdab F, Al-Aly Z, Alam K, Alam S, Alam T, Alanezi FM, Alanzi TM, Alemu BW, Alhabib KF, Ali M, Ali S, Alicandro G, Alinia C, Alipour V, Alizade H, Aljunid SM, Alla F, Allebeck P, Almasi-Hashiani A, Al-Mekhlafi HM, Alonso J, Altirkawi KA, Amini-Rarani M, Amiri F, Amugsi DA, Ancuceanu R, Anderlini D, Anderson JA, Andrei CL, Andrei T, Angus C, Anjomshoa M, Ansari F, Ansari-Moghaddam A, Antonazzo IC, Antonio CAT, Antony CM, Antriyandarti E, Anvari D, Anwer R, Appiah SCY, Arabloo J, Arab-Zozani M, Aravkin AY, Ariani F, Armoon B, Årnlöv J, Arzani A, Asadi-Aliabadi M, Asadi-Pooya AA, Ashbaugh C, Assmus M, Atafar Z, Atnafu DD, Atout MMDW, Ausloos F, Ausloos M, Ayala Quintanilla BP, Ayano G, Ayanore MA, Azari S, Azarian G, Azene ZN, Badawi A, Badiye AD, Bahrani MA, Bakhshaei MH, Bakhtiari A, Bakkannavar SM, Baldasseroni A, Ball K, Ballew SH, Balzi D, Banach M, Banerjee SK, Bante AB, Baraki AG, Barker-Collo SL, Bärnighausen TW, Barrero LH, Barthelemy CM, Barua L, Basu S, Baune BT, Bayati M, Becker JS, Bedi N, Beghi E, Béjot Y, Bell ML, Bennitt FB, Bensenor IM, Berhe K, Berman AE, Bhagavathula AS, Bhageerathy R, Bhala N, Bhandari D, Bhattacharyya K, Bhutta ZA, Bijani A, Bikbov B, Bin Sayeed MS, Biondi A, Birihane BM, Bisignano C, Biswas RK, Bitew H, Bohlouli S, Bohluli M, Boon-Dooley AS, Borges G, Borzi AM, Borzouei S, Bosetti C, Boufous S, Braithwaite D, Brauer M, Breitborde NJK, Breitner S, Brenner H, Briant PS, Briko AN, Briko NI, Britton GB, Bryazka D, Bumgarner BR, Burkart K, Burnett RT, Burugina Nagaraja S, Butt ZA, Caetano Dos Santos FL, Cahill LE, Cámera LA, Campos-Nonato IR, Cárdenas R, Carreras G, Carrero JJ, Carvalho F, Castaldelli-Maia JM, Castañeda-Orjuela CA, Castelpietra G, Castro F, Causey K, Cederroth CR, Cercy KM, Cerin E, Chandan JS, Chang KL, Charlson FJ, Chattu VK, Chaturvedi S, Cherbuin N, Chimed-Ochir O, Cho DY, Choi JY, Christensen H, Chu DT, Chung MT, Chung SC, Cicuttini FM, Ciobanu LG, Cirillo M, Classen TKD, Cohen AJ, Compton K, Cooper OR, Costa VM, Cousin E, Cowden RG, Cross DH, Cruz JA, Dahlawi SMA, Damasceno AAM, Damiani G, Dandona L, Dandona R, Dangel WJ, Danielsson AK, Dargan PI, Darwesh AM, Daryani A, Das JK, Das Gupta Rajat, das Neves J, Dávila-Cervantes CA, Davitciu DV, De Leo D, Degenhardt L, DeLang M, Dellavalle RP, Demeke FM, Demoz GT, Demisie DG, Denova-Gutiérrez E, Derveniz N, Dhungana GP, Dianatinasab M, Dias da Silva D, Diaz D, Dibaji Forooshani ZS, Djalalinia S, Do HT, Dokova K, Dorostkar F, Doshmangir L, Driscoll TR, Duncan BB, Duraes AR, Eagan AW, Edvardsson D, El Nahas N, El Sayed I, El Tantawi M, Elbarazi I, Elgendy IY, El-Jaafary SI, Elyazar IRE, Emmons-Bell S, Erskine HE, Eskandarieh S, Esmailnejad S, Esteghamati A, Estep K, Etemadi A, Etisso AE, Fanzo J, Farahmand M, Fareed M, Faridnia R, Farioli A, Faro A, Faruque M, Farzadfar F, Fattahi N, Fazlzadeh M, Feigin VL, Feldman R, Fereshtehjad SM, Fernandes E, Ferrara G, Ferrari AJ, Ferreira ML, Filip I, Fischer F,

Fisher JL, Flor LS, Foigt NA, Folyan MO, Fomenkov AA, Force LM, Foroutan M, Franklin RC, Freitas M, Fu W, Fukumoto T, Furtado JM, Gad MM, Gakidou E, Gallus S, Garcia-Basteiro AL, Gardner WM, Geberemariam BS, Ayalew Gebreslassie AAA, Geremew A, Gershberg Hayoon A, Gething PW, Ghadimi M, Ghadiri K, Ghaffarifar F, Ghafourifard M, Ghamari F, Ghashghaee A, Ghiasvand H, Ghith N, Gholamian A, Ghosh R, Gill PS, Ginindza TG, Giussani G, Gnedovskaya E V., Goharinezhad S, Gopalani SV, Gorini G, Goudarzi H, Goulart AC, Greaves F, Grivna M, Grosso G, Gubari MIM, Gugnani HC, Guimarães RA, Guled RA, Guo G, Guo Y, Gupta Rajeev, Gupta T, Haddock B, Hafezi-Nejad N, Hafiz A, Haj-Mirzaian Arvin, Haj-Mirzaian Arya, Hall BJ, Halvaei I, Hamadeh RR, Hamidi S, Hammer MS, Hankey GJ, Haririan H, Haro JM, Hasaballah AI, Hasan MM, Hasanpoor E, Hashi A, Hassanipour S, Hassankhani H, Havmoeller RJ, Hay SI, Hayat K, Heidari G, Heidari-Soureshjani R, Henrikson HJ, Herbert ME, Herteliu C, Heydarpour F, Hird TR, Hoek HW, Holla R, Hoogar P, Hosgood HD, Hossain N, Hosseini M, Hosseinzadeh M, Hostiuc M, Hostiuc S, Househ M, Hsairi M, Hsieh VCR, Hu G, Hu K, Huda TM, Humayun A, Huynh CK, Hwang BF, Iannucci VC, Ibitoye SE, Ikeda N, Ikuta KS, Ilesanmi OS, Ilic IM, Ilic MD, Inbaraj LR, Ippolito H, Iqbal U, Irvani SSN, Irvine CMS, Islam MM, Islam SMS, Iso H, Ivers RQ, Iwu CCD, Iwu CJ, Iyamu IO, Jaafari J, Jacobsen KH, Jafari H, Jafarinaia M, Jahani MA, Jakovljevic M, Jalilian F, James SL, Janjani H, Javaheri T, Javidnia J, Jeemon P, Jenabi E, Jha RP, Jha V, Ji JS, Johansson L, John O, John-Akinola YO, Johnson CO, Jonas JB, Joukar F, Jozwiak JJ, Jürisson M, Kabir A, Kabir Z, Kalani H, Kalani R, Kalankesh LR, Kalhor R, Kanchan T, Kapoor N, Matin BK, Karch A, Karim MA, Kassa GM, Katikireddi SV, Kayode GA, Kazemi Karyani A, Keiyoro PN, Keller C, Kemmer L, Kendrick PJ, Khalid N, Khammarnia M, Khan EA, Khan M, Khatab K, Khater MM, Khatib MN, Khayamzadeh M, Khazaei S, Kielling C, Kim YJ, Kimokoti RW, Kisa A, Kisa S, Kivimäki M, Knibbs LD, Knudsen AKS, Kocarnik JM, Kochhar S, Kopec JA, Korshunov VA, Koul PA, Koyanagi A, Kraemer MUG, Krishan K, Krohn KJ, Kromhout H, Kuate Defo B, Kumar GA, Kumar V, Kurmi OP, Kusuma D, La Vecchia C, Lacey B, Lal DK, Lalloo R, Lallukka T, Lami FH, Landires I, Lang JJ, Langan SM, Larsson AO, Lasrado S, Lauriola P, Lazarus JV, Lee PH, Lee SWH, Legrand KE, Leigh J, Leonardi M, Lescinsky H, Leung J, Levi M, Li S, Lim LL, Linn S, Liu Shiwei, Liu Simin, Liu Y, Lo J, Lopez AD, Lopez JCF, Lopukhov PD, Lorkowski S, Lotufo PA, Lu A, Lugo A, Maddison ER, Mahasha PW, Mahdavi MM, Mahmoudi M, Majeed A, Maleki A, Maleki S, Malekzadeh R, Malta DC, Mamun AA, Manda AL, Manguerra H, Mansour-Ghanaei F, Mansouri B, Mansournia MA, Mantilla Herrera AM, Maravilla JC, Marks A, Martin RV., Martini S, Martins-Melo FR, Masaka A, Masoumi SZ, Mathur MR, Matsushita K, Maulik PK, McAlinden C, McGrath JJ, McKee M, Mehndiratta MM, Mehri F, Mehta KM, Memish ZA, Mendoza W, Menezes RG, Mengesha EW, Merete A, Mereta ST, Meretoja A, Meretoja TJ, Mestrovic T, Miazgowski B, Miazgowski T, Michalek IM, Miller TR, Mills EJ, Mini GK, Miri M, Mirica A, Mirrahimov EM, Mirzaei H, Mirzaei M, Mirzaei R, Mirzaei-Alavijeh M, Misganaw AT, Mithra P, Moazen B, Mohammad DK, Mohammad Y, Mohammad Gholi Mezerji N, Mohammadian-Hafshejani A, Mohammadifard N, Mohammadpourhodki R, Mohammed AS, Mohammed H, Mohammed JA, Mohammed S, Mokdad AH, Molokhia M, Monasta L, Mooney MD, Moradi G, Moradi M, Moradi-Lakeh M, Moradzadeh R, Moraga P, Morawska L, Morgado-Da-Costa J, Morrison SD, Mospour A, Mosser JF, Mouodi S, Mousavi SM, Khaneghah AM, Mueller UO, Mukhopadhyay S, Mullany EC, Musa KI, Muthupandian S, Nabhan AF, Naderi M, Nagarajan AJ, Nagel G, Naghavi M, Naghshtabrizi B, Naimzada MD, Najafi F, Nangia V, Nansseu JR, Naserbakht M, Nayak VC, Negoj I, Ngunjiri JW, Nguyen CT, Nguyen HLT, Nguyen M, Nigatu YT, Nikbakhsh R, Nixon MR, Nnaji CA, Nomura S, Norrving B, Noubiap JJ, Nowak C, Nunez-Samudio V, Oancea B, Odell CM, Ogbo FA, Oh IH, Okunga EW, Oladnabi M, Olagunju AT, Olusanya BO, Olusanya JO, Omer MO, Ong KL, Onwujekwe OE, Orpana HM, Ortiz A, Osarenotor O, Osei FB, Ostroff SM, Otoi A, Otstavnov N, Otstavnov SS, Øverland S, Owolabi



- MO, Mahesh PA, Padubidri JR, Palladino R, Panda-Jonas S, Pandey A, Parry CDH, Pasovic M, Pasupula DK, Patel SK, Pathak M, Patten SB, Patton GC, Toroudi HP, Peden AE, Pennini A, Pepito VCF, Peprah EK, Pereira DM, Pesudovs K, Pham HQ, Phillips MR, Piccinelli C, Pilz TM, Piradov MA, Pirsaeheb M, Plass D, Polinder S, Polkinghorne KR, Pond CD, Postma MJ, Pourjafar H, Pourmalek F, Poznańska A, Prada SI, Prakash V, Pribadi DRA, Pupillo E, Syed ZQ, Rabiee M, Rabiee N, Radfar A, Rafiee A, Raggi A, Rahman MA, Rajabpour-Sanati A, Rajati F, Rakovac I, Ram P, Ramezanzadeh K, Ranabhat CL, Rao PC, Rao SJ, Rashedi V, Rathi P, Rawaf DL, Rawaf S, Rawal L, Rawassizadeh R, Rawat R, Razo C, Redford SB, Reiner RC, Reitsma MB, Remuzzi G, Renjith V, Renzaho AMN, Resnikoff S, Rezaei Negar, Rezaei Nima, Rezapour A, Rhinehart PA, Riahi SM, Ribeiro DC, Ribeiro D, Rickard J, Rivera JA, Roberts NLS, Rodríguez-Ramírez S, Roever L, Ronfani L, Room R, Roshandel G, Roth GA, Rothenbacher D, Rubagotti E, Rwegerera GM, Sabour S, Sachdev PS, Saddik B, Sadeghi E, Sadeghi M, Saeedi R, Saeedi Moghaddam S, Safari Y, Safi S, Safiri S, Sagar R, Sahebkar A, Sajadi SM, Salam N, Salamati P, Salem H, Salem MR, Salimzadeh H, Salman OM, Salomon JA, Samad Z, Samadi Kafil H, Sambala EZ, Samy AM, Sanabria J, Sánchez-Pimienta TG, Santomauro DF, Santos IS, Santos JV, Santric-Milicevic MM, Saraswathy SYI, Sarmiento-Suárez R, Sarrafzadegan N, Sartorius B, Sarveazad A, Sathian B, Sathish T, Sattin D, Saxena S, Schaeffer LE, Schiavolin S, Schlaich MP, Schmidt MI, Schutte AE, Schwebel DC, Schwendicke F, Senbeta AM, Senthilkumaran S, Sepanlou SG, Serdar B, Serre ML, Shadid J, Shafaat O, Shahabi S, Shaheen AA, Shaikh MA, Shalash AS, Shams-Beyranvand M, Shamsizadeh M, Sharafi K, Sheikh A, Sheikhtaheeri A, Shibuya K, Shield KD, Shigematsu M, Shin JLL, Shin MJ, Shiri R, Shirkoobi R, Shuval K, Siabani S, Sierpinski R, Sigfusdottir ID, Sigurvinsdottir R, Silva JP, Simpson KE, Singh JA, Singh P, Skiadaresi E, Skou ST, Skryabin VY, Smith EUR, Soheili A, Soltani S, Soofi M, Sorensen RJD, Soriano JB, Sorrie MB, Soshnikov S, Soyiri IN, Spencer CN, Spotin A, Sreeramreddy CT, Srinivasan V, Stanaway JD, Stein C, Stein DJ, Steiner C, Stockfelt L, Stokes MA, Straif K, Stubbs JL, Sufiyan MB, Suleria HAR, Suliankatchi Abdulkader R, Sulo G, Sultan I, Tabarés-Seisdedos R, Tabb KM, Tabuchi T, Taherkhani A, Tajdini M, Takahashi K, Takala JS, Tamiru AT, Taveira N, Tehrani-Banihashemi A, Temsah MH, Tesema GA, Tessema ZT, Thurston GD, Titova MV, Tohidnik HR, Tonelli M, Topor-Madry R, Topouzis F, Torre AE, Touvier M, Tovani-Palone MR, Tran BX, Travillian R, Tsatsakis A, Tudor Car IT, Tyrovolas S, Uddin R, Umeokonkwo CD, Unnikrishnan B, Upadhyay E, Vacante M, Valdez PR, van Donkelaar A, Vasankari TJ, Vasseghian Y, Veisani Y, Venketasubramanian N, Violante FS, Vlassov V, Vollset SE, Vos T, Vukovic R, Waheed Y, Wallin MT, Wang Y, Wang YP, Watson A, Wei J, Wei MYW, Weintraub RG, Weiss J, Werdecker A, West JJ, Westerman R, Whisnant JL, Whiteford HA, Wiens KE, Wolfe CDA, Wozniak SS, Wu AM, Wu J, Wulf Hanson S, Xu G, Xu R, Yadgir S, Yahyazadeh Jabbari SH, Yamagishi K, Yaminfirooz M, Yano Y, Yaya S, Yazdi-Feyzabadi V, Yeheyis TY, Yilgwan CS, Yilma MT, Yip P, Yonemoto N, Younis MZ, Younker TP, Yousefi B, Yousefi Z, Yousefinezhadi T, Yousuf AY, Yu C, Yusefzadeh H, Moghadam TZ, Zamani M, Zamanian M, Zandian H, Zastrozhin MS, Zhang Y, Zhang ZJ, Zhao JT, Zhao XJG, Zhao Y, Zheng P, Zhou M, Ziapour A, Zimsen SRM, Lim SS and Murray CJL (2019) Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the global burden of disease study 2019. *The Lancet* 396, 1223–1249.
- Akbaraly TN, Singh-Manoux A, Dugravot A, Brunner EJ, Kivimäki M and Sabia S (2019) Association of midlife diet with subsequent risk for dementia. *JAMA* 321, 957–968.
- Barbaresco J, Lellmann AW, Schmidt A, Lehmann A, Amini AM, Egert S, Schlesinger S and Nöthlings U (2020) Dietary factors and neurodegenerative disorders: an umbrella review of meta-analyses of prospective studies. *Advances in Nutrition* 11, 1161–1173.
- Borén J, John Chapman M, Krauss RM, Packard CJ, Bentzon JF, Binder CJ, Daemen MJ, Demer LL, Hegele RA, Nicholls SJ, Nordestgaard BG, Watts GF, Bruckert E, Fazio S, Ference BA, Graham I, Horton JD, Landmesser U, Laufs U, Masana L, Pasterkamp G, Raal FJ, Ray KK, Schunkert H, Taskinen MR, van de Sluis B, Wiklund O, Tokgozlu L, Catapano AL and Ginsberg HN (2020) Low-density lipoproteins cause atherosclerotic cardiovascular disease: pathophysiological, genetic, and therapeutic insights: a consensus statement from the European atherosclerosis society consensus panel. *European Heart Journal* 41, 2313–2330.
- Cao L, Tan L, Wang HF, Jiang T, Zhu XC, Lu H, Tan MS and Yu JT (2016) Dietary patterns and risk of dementia: a systematic review and meta-analysis of cohort studies. *Molecular Neurobiology* 53, 6144–6154.
- Ewers B, Marott JL, Schnohr P, Nordestgaard BG and Marckmann P (2021) Non-adherence to established dietary guidelines associated with increased mortality: the Copenhagen General Population study. *European Journal of Preventive Cardiology* 28, 1259–1268.
- Féart C, Samieri C, Rondeau V, Amieva H, Portet F, Dartigues JF, Scarmeas N and Barberger-Gateau P (2009) Adherence to a Mediterranean diet, cognitive decline, and risk of dementia. *JAMA* 302, 638–648.
- Ference BA, Ginsberg HN, Graham I, Ray KK, Packard CJ, Bruckert E, Hegele RA, Krauss RM, Raal FJ, Schunkert H, Watt GF, Borén J, Fazio S, Horton JD, Masana L, Nicholls SJ, Nordestgaard BG, Van De Sluis B, Taskinen MR, Tokgozlu L, Landmesser U, Laufs U, Wiklund O, Stock JK, Chapman MJ and Catapano AL (2017) Low-density lipoproteins cause atherosclerotic cardiovascular disease. 1. Evidence from genetic, epidemiologic, and clinical studies. A consensus statement from the European atherosclerosis society consensus panel. *European Heart Journal* 38, 2459–2472.
- Haring B, Wu C, Mossavar-Rahmani Y, Snelaelar L, Brunner R, Wallace RB, Neuhauser ML and Wassertheil-Smolter S (2016) No association between dietary patterns and risk for cognitive decline in older women with 9-year follow-up: data from the women's health initiative memory study. *Journal of the Academy of Nutrition and Dietetics* 116, 921–930.
- Hegele RA, Borén J, Ginsberg HN, Arca M, Averna M, Binder CJ, Calabresi L, Chapman MJ, Cuchel M, von Eckardstein A, Frikke-Schmidt R, Gaudet D, Hovingh GK, Kronenberg F, Lütjohann D, Parhofer KG, Raal FJ, Ray KK, Remaley AT, Stock JK, Stroes ES, Tokgozlu L, Catapano AL, von Eckardstein A and Rohrer L (2020) Rare dyslipidaemias, from phenotype to genotype to management: a European atherosclerosis society task force consensus statement. *The Lancet Diabetes and Endocrinology* 8, 50–67.
- Jørgensen AB, Frikke-Schmidt R, Nordestgaard BG and Tybjaerg-Hansen A (2014) Loss-of-function mutations in APOC3 and risk of ischemic vascular disease. *New England Journal of Medicine* 371, 32–41.
- Juul Rasmussen I, Rasmussen KL, Nordestgaard BG, Tybjaerg-Hansen A and Frikke-Schmidt R (2020) Impact of cardiovascular risk factors and genetics on 10-year absolute risk of dementia: risk charts for targeted prevention. *European Heart Journal* 41, 4024–4033.
- Kinney JW, Bemiller SM, Murtishaw AS, Leisgang AM, Salazar AM and Lamb BT (2018) Inflammation as a central mechanism in Alzheimer's disease. *Alzheimer's and Dementia: Translational Research and Clinical Interventions* 4, 575–590.
- Limongi F, Siviero P, Bozanic A, Noale M, Veronese N and Maggi S (2020) The effect of adherence to the Mediterranean diet on late-life cognitive disorders: a systematic review. *Journal of the American Medical Directors Association* 21, 1402–1409.
- Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S, Brayne C, Burns A, Cohen-Mansfield J, Cooper C, Costafreda SG, Dias A, Fox N, Gitlin LN, Howard R, Kales HC, Kivimäki M, Larson EB, Ogunniyi A, Orgeta V, Ritchie K, Rockwood K, Sampson EL, Samus Q, Schneider LS, Selbæk G, Teri L and Mukadam N (2020) Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *The Lancet* 396, 413–446.
- Luo J, Mills K, le Cessie S, Noordam R and van Heemst D (2020) Ageing, age-related diseases and oxidative stress: what to do next? *Ageing Research Reviews* 57, 100982.
- Mach F, Baigent C, Catapano AL, Koskinas KC, Casula M, Badimon L, Chapman MJ, De Backer GG, Delgado V, Ference BA, Graham IM, Halliday A, Landmesser U, Mihaylova B, Pedersen TR, Riccardi G, Richter DJ, Sabatine MS, Taskinen MR, Tokgozlu L, Wiklund O, Mueller C, Drexel H, Aboyans V, Corsini A, Doehner W, Farnier M, Gigante B, Kayikcioglu M, Krstacic G, Lambrinou E, Lewis BS, Masip J, Moulin P, Petersen S, Petronio AS, Piepoli MF, Pinto X, Raber L, Ray KK, Reiner Z, Riesen WF, Roffi M, Schmid JP, Shlyakhto E, Simpson IA, Stroes E, Sudano I, Tselepis AD, Viigimaa M, Vindis C,

- Vonbank A, Vrablik M, Vrsalovic M, Gomez JLZ, Collet JP, Windecker S, Dean V, Fitzsimons D, Gale CP, Grobbee DE, Halvorsen S, Hindricks G, Iung B, Jüni P, Katus HA, Leclercq C, Lettino M, Merkely B, Sousa-Uva M, Touyz RM, Nibouche D, Zelveian PH, Siostrzonek P, Najafov R, Van De Borne P, Pojskic B, Postadzhiyan A, Kypris L, Spinar J, Larsen ML, Eldin HS, Strandberg TE, Ferrières J, Agladze R, Laufs U, Rallidis L, Bajnok L, Gudjonsson T, Maher V, Henkin Y, Gulizia MM, Mussagaliyeva A, Bajraktari G, Kerimkulova A, Latkovskis G, Hamoui O, Slapikas R, Visser L, Dingli P, Ivanov V, Boskovic A, Nazzi M, Visseren F, Mitevska I, Retterstøl K, Jankowski P, Fontes-Carvalho R, Gaita D, Ezhov M, Foscoli M, Giga V, Pella D, Fras Z, De Isla LP, Hagström E, Lehmann R, Abid L, Ozdogan O, Mitchenko O and Patel RS (2020) 2019 ESC/EAS guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *European Heart Journal* **41**, 111–188.
- Martínez-Lapiscina EH, Clavero P, Toledo E, Estruch R, Salas-Salvadó J, San Julián B, Sanchez-Tainta A, Ros E, Valls-Pedret C and Martínez-González MÁ (2013) Mediterranean diet improves cognition: the PREDIMED-NAVARRA randomised trial. *Journal of Neurology, Neurosurgery and Psychiatry* **84**, 1318–1325.
- McKenzie BL, Harris K, Peters SAE, Webster J and Woodward M (2021) The association of energy and macronutrient intake with all-cause mortality, cardiovascular disease, and dementia: findings from 120,963 women and men in the UK Biobank. *British Journal of Nutrition* **127**, 1858–1867.
- Morris MC, Tangney CC, Wang Y, Sacks FM, Bennett DA and Aggarwal NT (2015) MIND diet associated with reduced incidence of Alzheimer's disease. *Alzheimer's & Dementia* **11**, 1007–1014.
- Nordestgaard BG and Varbo A (2014) Triglycerides and cardiovascular disease. *The Lancet* **384**, 626–635.
- Nordestgaard IT, Christoffersen M, Afzal S, Nordestgaard BG, Tybjaerg-Hansen A and Frikke-Schmidt R (2021) Triglycerides as a shared risk factor between dementia and atherosclerotic cardiovascular disease: a study of 125,727 individuals. *Clinical Chemistry* **67**, 245–255.
- Perez L, Heim L, Sherzai A, Jaceldo-Siegl K and Sherzai A (2012) Nutrition and vascular dementia. *Journal of Nutrition, Health and Aging* **16**, 319–324.
- Petersson SD and Philippou E (2016) Mediterranean diet, cognitive function, and dementia: a systematic review of the evidence. *Advances in Nutrition* **7**, 889–904.
- Pistollato F, Iglesias RC, Ruiz R, Aparicio S, Crespo J, Lopez LD, Manna PP, Giampieri F and Battino M (2018) Nutritional patterns associated with the maintenance of neurocognitive functions and the risk of dementia and Alzheimer's disease: a focus on human studies. *Pharmacological Research* **131**, 32–43.
- Rasmussen KL, Tybjaerg-Hansen A, Nordestgaard BG and Frikke-Schmidt R (2019) Plasma levels of apolipoprotein E, APOE genotype, and all-cause and cause-specific mortality in 105 949 individuals from a white general population cohort. *European Heart Journal* **40**, 2813–2824.
- Sassi F, Belloni A, Mirelman AJ, Suhrcke M, Thomas A, Salti N, Vellakkal S, Visarathvong C, Popkin BM and Nugent R (2018) Equity impacts of price policies to promote healthy behaviours. *The Lancet* **391**, 2059–2070.
- Skälén K, Gustafsson M, Knutsen Rydberg E, Hultén LM, Wiklund O, Innerarity TL and Boren J (2002) Subendothelial retention of atherogenic lipoproteins in early atherosclerosis. *Nature* **417**, 750–754.
- Springmann M, Spajic L, Clark MA, Poore J, Herforth A, Webb P, Rayner M and Scarborough P (2020) The healthiness and sustainability of national and global food based dietary guidelines: modelling study. *BMJ* **370**, m2322.
- The Official Dietary Guidelines – Good for Health and Climate (in Danish) (2020) Denmark: Ministry of Food Agriculture and Fisheries.
- Toft U, Kristoffersen LH, Lau C, Borch-Johnsen K and Jørgensen T (2007) The dietary quality score: validation and association with cardiovascular risk factors: the Inter99 study. *European Journal of Clinical Nutrition* **61**, 270–278.
- United Nations. The Sustainable Development Agenda – United Nations Sustainable Development, 29 June 2021. Available at <https://www.un.org/sustainabledevelopment/development-agenda/>.
- Valls-Pedret C, Sala-Vila A, Serra-Mir M, Corella D, De La Torre R, Martínez-González MÁ, Martínez-Lapiscina EH, Fitó M, Pérez-Heras A, Salas-Salvadó J, Estruch R and Ros E (2015) Mediterranean diet and age-related cognitive decline: a randomized clinical trial. *JAMA Internal Medicine* **175**, 1094–1103.
- Van Den Brink AC, Brouwer-Brolsma EM, Berendsen AAM and Van De Rest O (2019) The Mediterranean, dietary approaches to stop hypertension (DASH), and Mediterranean-DASH intervention for neurodegenerative delay (MIND) diets are associated with less cognitive decline and a lower risk of Alzheimer's disease – a review. *Advances in Nutrition* **10**, 1040–1065.
- Willett W, Rockström J, Loken B, Springmann M, Lang T, Vermeulen S, Garnett T, Tilman D, DeClerck F, Wood A, Jonell M, Clark M, Gordon LJ, Fanzo J, Hawkes C, Zurayk R, Rivera JA, De Vries W, Majele Sibanda L, Afshin A, Chaudhary A, Herrero M, Agustina R, Branca F, Lartey A, Fan S, Crona B, Fox E, Bignet V, Troell M, Lindahl T, Singh S, Cornell SE, Srinath Reddy K, Narain S, Nishtar S and Murray CJL (2019) Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *The Lancet* **393**, 447–492.
- Wilson PWF, Polonsky TS, Miedema MD, Khera A, Kosinski AS and Kuvin JT (2019) Systematic review for the 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APHA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Journal of the American College of Cardiology* **73**, 3210–3227.
- World Health Organization (2017) *Global Action Plan on the Public Health Response to Dementia*. Geneva, Switzerland: World Health Organization.
- World Health Organization Guidelines (2019) *Risk Reduction of Cognitive Decline and Dementia*. Geneva, Switzerland: World Health Organization.
- Wu L and Sun D (2017) Adherence to Mediterranean diet and risk of developing cognitive disorders: an updated systematic review and meta-analysis of prospective cohort studies. *Scientific Reports* **7**, 41317.
- Zhang H, Greenwood DC, Risch HA, Bunce D, Hardie LJ and Cade JE (2021) Meat consumption and risk of incident dementia: cohort study of 493,888 UK Biobank participants. *American Journal of Clinical Nutrition* **114**, 175–184.