

Journal Pre-proof

Unlocking Insights: Navigating COVID-19 Challenges and Emulating Future Pandemic Resilience Strategies with Strengthening Natural Immunity

Sunil J. Wimalawansa



PII: S2405-8440(24)10722-0

DOI: <https://doi.org/10.1016/j.heliyon.2024.e34691>

Reference: HLY 34691

To appear in: *HELIYON*

Received Date: 6 March 2024

Revised Date: 17 June 2024

Accepted Date: 15 July 2024

Please cite this article as: Unlocking Insights: Navigating COVID-19 Challenges and Emulating Future Pandemic Resilience Strategies with Strengthening Natural Immunity *HELIYON*, <https://doi.org/10.1016/j.heliyon.2024.e34691>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2024 Published by Elsevier Ltd.

Perspective—Heliyon

Unlocking Insights: Navigating COVID-19 Challenges and Emulating Future Pandemic Resilience Strategies with Strengthening Natural Immunity

Sunil J. Wimalawansa*

retired Professor of Medicine, Endocrinology, and Nutrition
Moratuwa, Sri Lanka

suniljw@hotmail.com; 0000-0003-1096-8595

*Correspondence: B14 G2, De Soyza Flats, Moratuwa, Sri Lanka

Sunil.wimalawasa@rutgers.edu (retired from Rutgers University):

Preferred: suniljw@hotmail.com

Running Head: “SARS-CoV—2 vaccines and Clinical Outcomes”

Perspective

Unlocking Insights: Navigating COVID-19 Challenges and Emulating Future Pandemic Resilience Strategies with Strengthening Natural Immunity

Abstract:

The original COVID-19 vaccines, developed against SARS-CoV-2, initially mitigated hospitalizations. Bivalent vaccine boosters were used widely during 2022-23, but the outbreaks persisted. Despite this, hospitalizations, mortality, and outbreaks involving dominant mutants like Alpha and Delta increased during winters when the population's vitamin D levels were at their lowest. Notably, 75% of human immune cell/system functions, including post-vaccination adaptive immunity, rely on adequate circulatory vitamin D levels. Consequently, hypovitaminosis compromises innate and adaptive immune responses, heightening susceptibility to infections and complications. COVID-19 vaccines primarily target SARS-CoV-2 Spike proteins, thus offering limited protection through antibodies. mRNA vaccines, such as those for COVID-19, fail to generate secretory/mucosal immunity-like IgG responses, rendering them ineffective in halting viral spread. Additionally, mutations in the SARS-CoV-2 binding domain reduce immune recognition by vaccine-derived antibodies, leading to immune evasion by mutant viruses like Omicron variants. Meanwhile, the repeated administration of bivalent boosters intended to enhance efficacy resulted in the immunoparesis of recipients. As a result, relying solely on vaccines for outbreak prevention became less effective. Dominant variants exhibit increased affinity to angiotensin-converting enzyme receptor-2, enhancing infectivity but reducing virulence. Meanwhile, spike protein-related viral mutations do not impact the potency of widely available, repurposed early therapies, like vitamin D and ivermectin. With the re-emergence of COVID-19 and impending coronaviral pandemics, regulators and health organizations should proactively consider approval and strategic use of cost-effective adjunct therapies mentioned above to counter the loss of vaccine efficacy against emerging variants and novel coronaviruses and eliminate serious adverse effects. Timely implementation of these strategies could reduce morbidity, mortality, and healthcare costs and provide a rational approach to address future epidemics and pandemics. This perspective critically reviews relevant literature, providing insights, justifications, and viewpoints into how the scientific community and health authorities can leverage this knowledge cost-effectively.

Keywords: 25(OH)D; COVID-19; Immune system; Mutations; Omicron; SARS-CoV-2

Lay Abstract:

The original COVID-19 vaccines effectively reduced hospitalizations but faced major challenges during winter outbreaks, leading to increased mortality, particularly in individuals with low vitamin D levels. Adequate vitamin D is critical for most human immune functions, including post-vaccination adaptive immunity. Hypovitaminosis compromises immunity and

heightens susceptibility to infections. COVID-19 mRNA vaccines predominantly target SARS-CoV-2 Spike proteins, providing limited antibody protection. Moreover, it provides no mucosal immunity, causing a failure to prevent the spread of the virus. In addition, mutations in the SARS-CoV-2, like Omicron, reduce vaccine recognition by antibodies (i.e., immune evasion). Besides, repeated bivalent vaccine boosters weaken the immune system.

Consequently, exclusive reliance on vaccines for outbreak prevention became ineffective. In contrast, Spike protein mutations do not impact the efficacy of repurposed (already approved generic) early therapies like vitamin D and ivermectin. Regulators and health organizations should consider cost-effective adjunct therapies to address emerging variants and enhance vaccine efficacy in the future. Timely implementation has the potential to reduce morbidity, mortality, and healthcare costs, providing a rational, cost-effective approach with no adverse effects for future pandemics.

1.0 Introduction:

The rapid development of messenger RNA (mRNA)-technology-based COVID-19 vaccines within six months in 2020 marked a significant technological milestone during the COVID pandemic.¹ ten-billion-dollar grant from United States taxpayers facilitated its developments. Traditionally, vaccines aim to halt disease transmission. However, mRNA-based COVID-19 vaccines, while effective in reducing hospitalizations, did not prevent the spread of SARS-CoV-2.^{2,3} Breakthrough infections in fully vaccinated individuals result in peak viral loads comparable to those in unvaccinated individuals, allowing both groups to continue transmitting SARS-CoV-2 to their contacts.² The Omicron variants, characterized by multiple mutations in the receptor binding domain, exhibit significant immunological differences from the original SARS-CoV-2.⁴ These variants demonstrate reduced lethality but with increased infectivity (high R_0),⁵ and immune evasion against vaccine-derived antibodies.

Despite the three years of the declared pandemic,⁶ the potential for coronaviruses to mutate and evade immunity has been well-known for years.⁷ Global health authorities, however, primarily advocated reliance on vaccines to control pandemic outbreaks without offering broader strategies or recommending alternative adjunct therapies for SARS-CoV-2 control and eradication.^{6,8} The focus has been primarily on patented COVID-19 vaccines, antiviral agents, and monoclonal antibodies authorized under Emergency Use Authorization (EUA).⁹

In the USA and other countries, patented agents for COVID-19 received swift authorization from the Food and Drug Administration (FDA). Notably, one critical legal requirement for EUA approval and maintenance is the absence of alternative agents for disease prevention or treatment.⁹ The approvals of COVID-19 vaccines and antiviral agents under EUA were expedited and released to the market without undergoing rigorous efficacy and longer-term adverse effect testing.^{9,10} Despite the prompt approval of the patented COVID-19 vaccines, anti-viral agents, and monoclonal antibodies by regulators, a global urgency to identify and approve a genuinely compelling, cost-effective agent against SARS-CoV-2 failed to materialize.¹¹⁻¹³

The paper delves into aspects concerning COVID-19 outbreaks in industrialized countries and explores alternate options, specifically the potential use of vitamin D for future pandemic control. The author underscores concern (unlike traditional vaccines) regarding the

predominant reliance on vaccines, pointing out their limitations in preventing infection and transmission. The potential use of globally available cost-effective repurposed agents, such as vitamin D and ivermectin, is evaluated. The manuscript also emphasizes the significance of recognizing natural immunity and advocates for considering alternative therapeutic approaches for the future.

A thorough search was conducted across various research databases using keywords related to COVID-19 vaccines, mRNA vaccines, complications, immune evasion, vaccine efficacy, alternative therapies, vitamin D, and ivermectin. Combining keywords narrowed the number of relevant manuscripts to a manageable quantity. The search encompassed databases such as PubMed, Medline, Web of Science, and EMBASE, focusing on clinical studies, randomized controlled clinical trials (RCTs), prospective clinical studies, as well as original and review articles, following a methodology similar to that employed for systematic or narrative reviews.¹⁴ References were selected based on their relevance to the topic and incorporated into the manuscript after reviewing them.

1.1 Fundamentals of COVID vaccines

The mRNA and adenovirus vector-based COVID vaccines were designed based on the viral Spike protein sequences. mRNA-based immunizations utilize single-stranded mRNA to encode the desired antigen. The objective was to prompt the recipient's immune system to produce viral Spike proteins of SARS-CoV-2, comprising 1273 amino acids, to develop antibodies against the virus.¹⁵ In different vaccines, the modified Spike protein structures or selected portions of the viral sequences are not identical. As a result, the antigen expression and antibodies generated from mRNA and genetic adenovirus-vector vaccines differ.¹⁵

COVID-19 vaccines were developed using different platforms, including live attenuated viruses, inactivated vaccines, and non-replicating viral vectors—RNA and DNA. mRNA vaccines were created using DNA templates encoding the Spike protein. These mRNA particles are encapsulated in a lipoprotein-based carrier to safeguard them from degradation and facilitate rapid cell entry.¹⁶ mRNA molecules enter cells and undergo translation upon injection via the intramuscular route.¹⁷ Once inside host cells, mRNA generates peptides with Spike protein sequences recognized by the body as foreign antigens, activating humoral responses and prompting B-cells to transform into memory cells. Viral antigens are presented on antigen-presenting cells, triggering immune responses. Upon secondary exposure to a Spike protein antigen, memory B cells activate protective mechanisms, including heightened synthesis of neutralizing antibodies.¹⁸

The S1 subunit of the SARS-CoV-2 spike protein contains the receptor binding domain (RBD) along with the N-terminal domain, which is responsible for binding it to the angiotensin-converting enzyme-2 (ACE-2). The S2 subunit houses the fusion and transmembrane proteins, enabling viral attachment to the host cell.¹⁹ After the virus enters host cells, it undergoes uncoating and initiates the transcription and translation of mRNA, leading to viral assembly. The resulting exocytosis of viral particles from host cells into the bloodstream allows them to infect other cells.

Unlike vaccines developed through traditional approaches, the mRNA vaccines (i.e., Pfizer and Moderna) and adenovirus vector-based COVID-19 vaccine (J&J) did not

incorporate core viral elements or the complete inactivated virus.^{20,21} Immunity induced through traditional methods uses core components, which tend to have a broader spectrum and longer-lasting immune protection.^{22,23} In contrast, the immunity triggered by the mRNA COVID-19 vaccine is more focused, targeting a part of the Spike protein sequence. Consequently, mRNA vaccines do not produce antibodies against co-components, like nucleocapsids and co-proteins.^{24,25}

Random mutations occur, affecting the binding domains of the SARS-CoV-2 genome, which may modify the affinity to the membrane-bound ACE-2 receptor. Such mutations can modify the virulence and the potential for immune evasion.²⁴ However, dominant variants increase infectiousness as indicated by higher R_0 values (replication rate) while exhibiting reduced lethality and developing an increased ability to evade immune detection.²⁶ The virulence of the newer strains of SARS-CoV-2 strain is comparable to influenza,²⁷ resulting in a decreased need for hospitalization. Additionally, over the past years, the widespread availability of free Rapid Antigen (RAT) kits (intended for screening tools and not to make diagnosis) has led to a sizable portion of COVID-19 testing being conducted at home or health-related stores. Consequently, positive SARS-CoV-2 data might not have been consistently captured in general reporting or hospital statistics, potentially causing authorities to overlook or ignore outbreaks.

1.2 Basic mechanisms involved with gene-directed vaccines

The fundamental mechanism underlying mRNA vaccine technology involves delivering nucleic acid molecules that encode the target antigen, specifically the coding region of the Spike sequences of SARS-CoV-2. These strategically designed sequences utilize host cells to produce foreign mRNA, expressing and amplifying the intended antigen—producing spike protein sequences through ribosomes.²⁸ This intricate process entails synthesizing antigen sequences into corresponding proteins, with some degrading into smaller antigenic peptides by proteasomes.²⁵ However, the reported critical adverse effects and the dispersion of Spike protein-related mRNA, as suggested by mRNA scatter, indicate that the distribution of this material is not limited to immune cells.^{29,30}

Once in circulation, antigens interact with immune cells that recognize antigens, such as Toll-like receptor-4, which is distributed extensively in the body. In contrast to traditional protein-based vaccines, mRNA (or DNA) conveys the message to host cells to generate and express Spike proteins on their membrane. The soluble form of these proteins is then shed into circulation to alert the immune system.⁵ Consequently, a broad range of cells in the body is exposed to the Spike protein, leading to the reported adverse effects.²⁴

The antigenic peptides are presented to CD8+ cytotoxic T cells via the major histocompatibility complex (MHC) class I molecular pathway as endogenous antigens, activating cell-mediated immune responses.²⁵ Additionally, these antigenic peptides, derived from toxic Spike proteins generated by mRNA, are released into the extracellular environment, including the circulation.³¹ They are taken up by antigen-presenting cells (APCs) and presented to CD4+ T cells through MHC class II molecules as exogenous antigens. This initiates cellular immune responses through cytokines, activating B cells to produce antibodies and triggering humoral immune responses.³²

1.3 The development of COVID-19 vaccines and new antiviral agents

Novel COVID vaccines against the SARS-CoV-2 virus were developed swiftly, circumventing standard regulatory and safety requirements through EUA.³¹ Launching them in late 2020 was considered a significant achievement.³³ The development of this innovative technology for SARS-CoV-2 vaccines was financially supported by the United States taxpayer grant to Big Pharma, benefitting multiple major pharmaceutical companies capable of large-scale vaccine manufacturing.³⁴ Additional financial subsidies were extended in 2021 to develop antiviral agents. In addition, pharmaceutical companies received special exemptions and fast-track approvals (under the EUA Act) from the FDA to develop, test, and approve these vaccines.⁹

The fallacy emerged when vaccine companies and leading health authorities asserted that these vaccines prevented SARS-CoV-2 transmission and propagation while withholding primary data. The primary reason most governments purchased expensive COVID-19 vaccines and people believed this later became a falsehood. This contributed to a broader public distrust and misconceptions about vaccination and directions by health organizations like the WHO.³⁵ Large health agencies propagated vaccines as the sole solution to the pandemic, prompting many governments to coerce and mandate them.³⁶ Despite knowing about previous coronaviral infections, vaccine companies failed to design COVID-19 vaccines properly. Instead, they expedited their production by taking shortcuts in vaccine development and testing.³¹ This further added to the public distrust. The failure to address and mitigate adverse outcomes proactively and disclose critical information that these vaccines do not prevent infections further eroded their credibility and trust, which later affected their uptake.^{25,37}

The EUA and the favorable pharmaceutical agreements with the FDA allowed Big Pharma to bypass standard safety checks in COVID-19 vaccine trials, which would have otherwise taken several years.³⁸ Conversely, regulators and leading health authorities chose not to authorize using already approved, generic, cost-effective, repurposed agents.³⁹ Instead, they penalized physicians who used them. Relaxing of regulations includes otherwise mandatory toxicity studies (such as precluding genotoxicity studies), relaxation of standards of conducting and gathering data, the number of RCTs to confirm their efficacy, and the need for rigorous short-term and long-term adverse effect testing and reporting.³¹

Based on the EUA, regulators also lowered the threshold for experimental and untested methods in generating mRNA vaccines and human testing, later expanding and replicating them to new antiviral agents and monoclonal antibodies.⁴⁰ It became evident that if regulators were to approve any other agent for preventing or treating SARS-CoV-2, the temporary EUA approval of vaccines and antiviral agents would be revoked automatically—a critical reason for the non-approval of any generic agents for the prevention or treatment of SARS-CoV-2.

1.4 Regulatory categorization of COVID-19 vaccines

According to the definitions provided by the European Union (Section IV), New Zealand, and other regulatory bodies, Pfizer-BioNTech and Moderna's mRNA vaccines were categorized as gene therapies.⁴¹ Additionally, AstraZeneca and J&J's adenoviral (nucleic acid) vector (injections) were identified as modified gene therapies.⁴¹ These vaccines were designed

to synthesize large quantities of viral Spike proteins *in vivo* in the recipients, subsequently stimulating memory and plasma cells and leading to the generation of antibodies.⁴² Therefore, it is unsurprising that the adverse effects on vulnerable people following the SARS-CoV-2 virus and COVID-19 vaccines are similar.

1.5 Glimpse of virus and vaccine-related adverse effects

With billions of dollars in grant funding and impunity against adverse effects, Big Pharma was incentivized by the FDA to bring the mRNA vaccine to market quickly.³⁵ Besides, based on agreements and urgency, regulators increased tolerance of big pharma companies taking shortcuts and maintained the secrecy of the approval process.³¹ In conjunction with the FDA, Big Pharma refused to share methodology, process, vaccine-related primary RCT data, and adverse effects, using the pretense of patent protection. The vacuum of data availability to independent scientists from pharma and the FDA was striking and unprecedented.⁴³ Additionally, the subsequent acceptance of bivalent COVID-19 vaccines was without RCT (data), as well as the vaccination of children and young adults.⁴⁴ Vaccine companies provided neither safety data related to these groups nor supporting tangible benefits vs. potential harm.⁴⁵⁻⁴⁷

COVID-19 vaccines have been associated with broader and serious adverse effects,^{45,48} primarily affecting the cardiovascular and pulmonary systems.⁴⁹⁻⁵¹ Similar adverse complications have been reported following SARS-CoV-2 infection.^{48,52,53} They also linked to sudden death syndrome,⁵⁴ a multisystem inflammatory syndrome in adults⁵⁵ and children,⁵⁶ as well as adverse outcomes in pregnancy,⁵⁷ myocarditis,^{56,58} pericarditis,⁴⁶ clotting abnormalities (hemagglutinin),⁵⁹ and various other adverse effects in cardiological, hematological,⁶⁰ dermatological,^{61,62} ocular,⁶³ and neurological systems.^{53,64}

Inference: The unexpected and broad range of spike protein-mediated adverse effects began to be reported after the approval of COVID-19 vaccines⁵². These severe adverse effects from COVID-19 vaccines affirmed insufficient testing or withholding of adverse reaction data and insufficient follow-up duration before their broader clinical use was approved.⁶⁵ Data confirmed systemic adverse effects (e.g., pulmonary, cardiovascular, and neural effects) stem from disseminating toxic Spike Proteins throughout the body. A phenomenon was observed following SARS-CoV-2 infection and after COVID-19 vaccines.^{52,53,65}

1.6 The progress of vaccines and their efficacy

With a broader uptake, WHO and government sponsorships, vaccines showed promise in early 2021 by decreasing the severity of SARS-CoV-2 infection, resulting in fewer hospitalizations. However, no reduction in deaths has been demonstrated.³³ However, towards the end of 2021, vaccine efficacy declined. In response, mRNA companies such as Pfizer and Moderna developed a hybrid solution—creating bivalent vaccines designed to cover the antigenic sites of emerging Omicron variants.^{66,67} Despite assurances from vaccine pharmaceutical companies and leading health authorities regarding high efficacy and safety and the emphasis on vaccine reliance, opting to rely solely on vaccines to overcome the pandemic appears to have been a mistake.⁶⁸ Such has also led to vaccine hesitancy.⁶⁸

The diminishing efficacy of vaccines, especially bivalent boosters, coupled with the lack of an alternative plan or effective armamentarium, failed to control COVID-19 with

increasing virus outbreaks in 2022, noticeable in industrialized countries with high vaccine uptake.⁶⁹ Despite optimistic expectations, outbreaks persisted regardless of the vaccination status in the community.⁷⁰ This article critically examines the advantages and disadvantages of COVID-19 vaccines and the potential alternative adjunct therapies based on publicly available information. The data suggested that despite widespread vaccination efforts, the incidence of infections and hospitalizations from SARS-CoV-2 experienced a resurgence and continued, particularly during the winter of 2022 and 2023.^{35,53,71}

There is a cyclic decline in the serum 25(OH)D concentration population during winter.⁷² This is attributed to a lack of sunlight (i.e., exposure to UVB rays) and inadequate dietary intake and supplementation.^{73,74} Hypovitaminosis D weakens the immune system,⁵ increasing the vulnerability to respiratory viruses: flu viruses, SARS-CoV-2, etc. These viruses spread primarily through airborne mechanisms.^{75,76} This phenomenon explains the seasonal escalations of respiratory viral illnesses like flu, respiratory syncytial virus, and SARS-CoV-2 during winter (Figure 1).⁷⁷ people were infected predominantly in indoor gatherings (inhalation of air with high viral density), and dry air facilitated the spreading of the virus. This behavior change following prolonged curfews and lockdowns had made hypovitaminosis D worse.^{77,78} The failure to effectively control SARS-CoV-2 in 2020/21 and mismanaged pandemic guidance resulted in COVID-19 becoming endemic by early 2023.⁷⁹

< Figure 1 >




Figure 1: Winter-associated low serum 25(OH)D levels in the population increase susceptibility to respiratory viral infections, including those caused by SARS-CoV-2 (modified from Wimalawansa, SJ, 2022).⁷⁷

2.0 SARS-CoV-2 mutations, vaccine uptake, and viral transmissions

The efficacy of COVID-19 vaccines resulted in fewer hospitalizations from SARS-CoV-2 in early 2021 and faded by 2022, primarily due to emerging viral mutations and their

escape from neutralizing antibodies.³³ Since both mRNA and adenoviral vector COVID-19 vaccines failed to generate mucosal immunity.^{2,80,81} This was particularly notable as SARS-CoV-2 viruses and their variants predominantly spread via airborne transmission.^{2,82} As a result, there were no significant differences in infectivity rates between vaccinated and unvaccinated individuals.⁸¹ Furthermore, new variants like Delta and Omicron continually acquired new Spike protein mutations,⁸³ rendering them more contagious than the original SARS-CoV-2,^{2,84} but less lethal,⁸⁵ aiding the spread of the virus.

2.1 Viral mutations, immune evasion, and lack of effect on viral transmission

Evolving dominant mutated virus increased the affinity of ACE-2-RBD of the virus, enhancing the infectivity.⁸⁶ Some viruses become dominant, others develop immune evasion capabilities,^{6,7} rendering them resistant to vaccines,¹⁰ and antiviral agents.⁸⁷ New Omicron variants have higher R_0 values (the average number of people getting infected from a person),²⁶ indicating increased infectiousness,⁸⁸ but generally, they became less lethal than the original SARS-CoV-2 and the Delta variant.^{26,76} The loss of vaccine effectiveness due to mutagenesis and the resultant immune evasion and disordered and/or conflicted policies have contributed to the persistence of SARS-CoV-2 outbreaks in 2021 and 2022, becoming endemic. The following sections discuss these aspects and propose cost-effective measures to overcome outbreaks.

2.2 Viral loads and SARS-CoV-2 transmission

As previously discussed, several studies have highlighted the absence of a significant difference in SARS-CoV-2 nasal viral loads between vaccinated and unvaccinated individuals.^{2,64,89} This includes the presence of the virus in the nasopharynx, even in asymptomatic cases.^{2,80,81} Consequently, fully vaccinated and unvaccinated individuals contribute equally to the spread of the disease.⁸⁹ While some variations in reporting exist due to technological differences, the consensus is that, unlike traditional vaccines, COVID-19 vaccines do not prevent disease transmission—from human to human. As a result, mandates for vaccinations have been deemed irrelevant⁹⁰ and unjustified.⁹¹

The above points are supported by studies conducted in Israel, showing COVID-19 breakthrough transmission from fully vaccinated healthcare workers to patients.^{91,92} Various studies have consistently demonstrated no significant difference in the peak viral titers in the upper airways and the culturable viruses between vaccinated and unvaccinated individuals.^{2,80,81,92} This evidence challenges the justification for policies mandating compulsory vaccination and the dismissal of unvaccinated healthcare workers,⁹⁰ frontline workers, and military personnel.^{2,80}

2.3 Disparity to access vaccines—Industrial nations vs. developing countries

Global distribution of COVID-19 vaccines faced a significant disparity. Wealthier nations secured preferential status through multi-billion-dollar pre-marketing contracts and pre-payments by mRNA vaccine companies—Pfizer and Moderna—with payments to mRNA vaccine companies. This resulted in receiving the vaccine supplies first.⁹³ In contrast, the rollout of vaccines to developing countries faced delays of several months.⁹⁴ Many of these countries struggled to secure a smaller percentage of vaccines needed.

Lack of funds prevented them from purchasing expensive COVID-19 vaccines or antiviral agents in sufficient quantities, even for their vulnerable populations. In addition, under public pressure, some of these nations had to compete and spend more than the market price to acquire COVID-19 vaccines.⁹⁵ This was escalated because of the lack of raw material supply from the West to manufacture vaccines in countries like India.⁹⁶ This disparity underscores the challenges in ensuring equitable access to vaccines worldwide.⁹⁵

The focus of big pharma on enhancing upfront profits created major challenges for developing nations, as they lacked the capacity for large pre-payments in (USD) foreign currencies, making it difficult for them to secure delivery contracts.⁹⁵ This neglects over four billion people in developing nations without having adequate access to vaccines.⁹⁶ This may also contribute to the failure of the development of herd immunity and SARS-CoV-2 infection becoming endemic.⁹³ and allow vaccine pharma to raise prices for further vaccines. Consequently, areas like Africa and certain Asian countries were left behind with very low vaccination rates.⁹⁴ Ironically, these regions that failed to secure mRNA-based COVID-19 vaccines experienced the lowest mortality from SARS-CoV-2.⁹⁷ The health emergency declared by the World Health Organization (WHO) was managed haphazardly, leading to non-transparent industry practices and creating loopholes that allowed politicians,⁹⁰ especially in developing countries, to exploit the market for personal gains.

Inference: The mistaken premise that the COVID-19 vaccine would stop propagating SARS-CoV-2 infection drove developing countries to spend precious funds earmarked for essential needs like food and medicine to prioritize vaccine purchases. This stretched their limited foreign reserves, compromising the purchase of essential goods and oil, leading to increased food insecurity.⁹⁴ The diversion of resources towards vaccine procurement, often at exorbitant prices, was influenced by commissions and additional profits for those involved.⁹³ This financial strain compromised essential services in emerging economies and developing countries, highlighting the detrimental impact of poor policy and public health decisions affecting the well-being of their populations.^{96,98}

Developing countries, such as Burundi, Haiti, Chad, Ethiopia, Cameroon, Tanzania, Mali, etc., with lower vaccination rates (ranging from 0.5 to 5% of the population), experienced notably lower hospitalization and death rates from SARS-CoV-2.^{93,96} Whereas, countries like Sri Lanka, faced challenges after paying double the market rate to obtain COVID-19 vaccines. Poor fiscal policies and decision-making led to the dwindling of their foreign reserves and devaluation of currencies. The resulting exchange crisis was exacerbated by unwise decisions, such as banning fertilizer and pesticide importation, significantly reducing crop output and leading to severe food shortages.⁹⁵ This crisis, marked by severe shortages of essential goods like food, medicine, and fuel, ultimately led to Sri Lanka declaring bankruptcy in 2022.^{99,100}

2.4 Securing COVID-19 vaccines and government mandates to increase uptake

Despite efforts in industrialized countries to boost vaccination uptakes through coercive measures like imposing mandates and passports and restrictions for unvaccinated people, viral outbreaks persisted, as in most industrialized countries, including the USA.^{90,101} Inherent limitations of COVID-19 vaccines, the inability to stimulate mucosal immunity,⁵¹ played a

significant role in the viral spread and the failure to achieve herd immunity and eventually eliminate SARS-CoV-2.^{2,80,81}

Relying solely on vaccines at any cost, many administrations enforced vaccine mandates and extended curfews based on the misconception that using vaccination alone will prevent the transmission of SARS-CoV-2.⁹⁰ This approach neglected cost-effective generics, repurposed drugs, and essential traditional public health disease prevention measures. Additionally, governments using their agencies and health authorities utilized Big Tech and mass media to discredit and discourage the use of already approved and widely available repurposed agents, including vitamin D and ivermectin.^{102,103}

Inference: The lack of transparency, information censorship, and restricting access to scientific publishing prevented healthcare workers and the public from being adequately informed about the benefits and drawbacks of vaccines and alternative therapies,¹⁰¹ along with their efficacy and adverse effects.^{104,105} The focus was shifted toward a new vaccination round, using bivalent boosters designed for new mutants. However, these booster vaccines were approved without conducting RCTs.¹⁰⁶ Despite these efforts by prominent health organizations and Western governments, public skepticism increased with a parallel reduction of vaccine uptake in 2022.⁷⁰

2.5 Generation of bivalent vaccines and regulatory framework

In early 2022, vaccine companies acknowledged the fading efficacy of the original COVID-19 vaccine, prompting the development of hybrid bivalent vaccines. These were marketed as an innovative solution against the Omicron variants.^{1,106} For 100 vaccines, pharma emphasized the similarity in the manufacturing process to bypass the need for new RCTs. With little transparency and a lack of credible safety data, including genotoxicity and medium- to longer-term adverse effects, these were promptly approved by the FDA and other regulatory authorities.^{2,45,122,123,91,101} Regulatory approval for mRNA-based booster/bivalent vaccines, monoclonal antibodies, and antiviral agents was swiftly granted under the Public Health Emergency of International Concern²¹ declared by WHO.^{107,108} This hasty expedited approval process facilitated rapid market penetration influenced by governmental measures like booster vaccine mandates.

Inference: The untransparent collaboration between Big Pharma and regulators, such as the FDA, led to a critical error by overlooking efficacy and safety testing for bivalent vaccines.⁶⁶ Approval for commercialization without adequate testing, safety data, and RCT data was a significant regulatory oversight akin to coercing parents to vaccinate children. Subsequent data revealed that bivalent vaccine boosters had only about a third of the efficacy of the original COVID-19 vaccines, but others disagree.¹⁰⁹ The new Omicron variants had a substantial capability for immune evasion against bivalent booster vaccines.¹⁰⁶ These events have contributed to ongoing outbreaks and the pandemic transitioning to an endemic state.^{66,85,110,111}

2.6 Insufficient futuristic thinking and planning

The progress of medicine over the last three decades has been remarkable, offering a wide range of choices, advanced diagnostic methods, and innovative interventions;¹¹² most

recently, the use of artificial intelligence. However, information technologies have outpaced practical applications in the medical field. Despite the potential of machine learning and artificial intelligence,^{79,113} their integration into medical decisions and policy-making is still in its early stages.^{79,113} This lag in incorporating these technologies has resulted in shortcomings in drug and device safety assessments, interactions, and efficacy evaluation.^{114,115} Consequently, there have been delays in approving drugs, including cost-effective generic agents,^{79,113} and making timely healthcare policy decisions, presenting a significant obstacle to progress.^{79,114}

For instance, since mid-2020, compelling data from multi-disciplinary studies, RCTs,¹¹⁶⁻¹¹⁸ meta-analyses, and extensive database studies have consistently shown the effectiveness of widely available, unpatented agents and micronutrients in enhancing the immune system capabilities.^{5,119,120} These interventions have demonstrated the potential to prevent complications and reduce mortality associated with coronaviruses like SARS-CoV-2 (<https://c19early.org/dmeta>).⁹⁷ However, regulatory bodies and prominent health authorities have overlooked this evidence.

2.7 Inherent conflicts of interests

Two examples of conflicts of interest are repurposed agents vitamin D,¹²⁰⁻¹²² and ivermectin.^{123,124} In late 2020, each agent had over forty reasonably sized peer-reviewed published clinical studies, including RCTs (<https://c19early.org>).⁹⁷ By the end of 2022, there were over 140. By mid-2023, there were over 250 peer-reviewed publications for each agent.⁹⁷ In-depth details of all these publications are available free on one website—<https://c19vitaminmind.com>.⁹⁷ Despite this large amount of clinical trial data, regulators and healthcare agencies were unwilling to investigate these agents for their efficacy and safety.¹¹³ Furthermore, the pharmaceutical industry began designing clinical studies and funding for academia to carry out and poorly designed and biased studies and publish them via academic institutes. Some trials were designed to fail, and others intended to discredit generic agents.^{102,125}

The relevant data supporting the efficacy of early interventions had already been published, and outcomes were well-known. Therefore, authors should have known a high probability of failure of outcomes and, thus, harm to subjects when designing those failed clinical studies.^{126,127} Therefore, ethics committees (IRBs) should not have approved such clinical studies.

Most RCTs mentioned above utilized inappropriate and/or insufficient doses of economic agents like vitamin D and ivermectin.^{128,129,130,131} In some studies, they were administered at the wrong frequency or only as a bolus dose,^{128,129} in severely ill individuals (e.g., in ICUs) or late in the course of COVID-19.^{129,132,133} Others had defective study designs.^{130,131} It is noteworthy that most of these generic therapies are most effective when administered early (as possible) as a preventative measure prior to developing complications. Such an approach is intended to provide either prophylactic treatment to prevent symptomatic disease or on admission before complications develop.¹³⁴

These repurposed agents were approved for other conditions, are widely available worldwide without a prescription, and are economical. In addition, they have impressive safety profiles (refer to section 4.8).^{97,135,136} As discussed above, the main reason for not approving these generic agents was the automatic revocation of vaccines, antivirals, and

monoclonal antibodies temporarily approved under EUA (i.e., to protect their status) if these agents were approved—to protect the approval of the latter for clinical use.⁷⁹

2.8 The failure to approve repurposed early therapies

Despite four years into the pandemic, regulators have yet to approve any non-patented cost-effective therapies for preventing and treating SARS-CoV-2. This narrow perspective has resulted in missed opportunities to establish logic-based COVID-19 adjunctive algorithms to manage it better using algorithms with complementary therapies, even though it could have pejorative connotations in mainstream medicine.^{79,113} Such could have saved lives instead of relying on most countries' recommended supportive and/or wait-and-see strategy in 2020.¹³⁴ By mid-2020, substantial published data encompassing over 390,000 individuals (i.e., big data) and information on vulnerable patient groups were available.⁷⁹ Utilization of these would have offered insights into early diagnosis, age-sex-specific risks, and cost-effective early use of generic agents.^{113,116} No steps have been taken to integrate this information into clinical practice.

The evidence-based medicine paradigm established by big pharma for drug approvals has led regulators to focus solely on pharmaceutical industry-conducted, large, multi-center RCTs as the primary (and the only) evidence for testing medical hypotheses for drug approvals, which was extended to nutrients like vitamin D.^{5,137} This preferential approach negatively affects the approval process for unpatented and widely available inexpensive agents, such as vitamin D and ivermectin for SARS-CoV-2.⁹⁷ Due to this misconception, well-conducted extensive observational and ecological studies, along with smaller but well-designed and statistically powered RCTs conducted by sources outside the pharma sector, such as academia, were disregarded by regulators in the context of drug approvals. Contrary to this bias, these clinical studies are equally valid for understanding causative factors and assessing effectiveness, even though they may involve generics and nutrient interventions.⁵

Inference: Authorities neglected to employ innovative logical methods, extensive data analyses, big-data analysis, and novel machine-learning paradigms for decision-making despite the availability of a large set of data by the end of 2020^{79,113}. Utilizing these approaches could have provided evidence of the efficacies of vitamin D and ivermectin in mid-2020, confirming their clinical relevance, consistency, robust dose responses, and inverse correlations between vitamin D status and COVID-19-related outcomes,⁷³ similar to ivermectin.⁹⁷ Integrating new information and engineering technologies into large clinical study datasets could have minimized biases and guesswork used by health authorities over the past three years. The failure to do so resulted in fruitless policies like lockdowns and weak, ambiguous clinical recommendations, contributing to increased economic disruptions, healthcare costs, hospitalizations, intensive care usage, and deaths, especially in high-risk groups.^{79,113}

2.9 Misinformation and data suppression

Throughout the COVID-19 pandemic, there has been a proliferation of misinformation by individuals on social media regarding disease prevention and treatment aspects (57). Numerous challenges in accessing authentic information include preventing the dissemination of the beneficial effects of vitamin D and ivermectin via scientific articles and reviews.

Pharmaceutical companies and governments supported the process sponsored *fact-checking* organizations.¹³⁸⁻¹⁴⁰ Besides, a coordinated effort by mainstream media and major social media platforms was used to undermine the use of generics and repurposed medications for COVID-19.¹⁴¹⁻¹⁴³ Additionally, there had been attempts to divert regulatory attention from considering repurposed agents for COVID-19 by presenting sophisticated but irrelevant statistics instead of focusing on straightforward, traditional, and cost-effective analyses and assessments.¹⁴⁴

These repurposed agents' effectiveness has been downplayed, and adverse effects have been exaggerated,¹⁴⁵—designed to undermine the public trust. Such disinformation is defined as knowingly sharing false information to cause harm.¹⁴⁶ Even some scientific articles initially published were later found to contain unintentional misinformation.^{138,139,141,147} Medical boards and councils have inappropriately taken politically-driven disciplinary actions against physicians who prescribed these cost-effective medications with good intentions and the best judgment to benefit patients.^{140,143} The usefulness of generic agents like vitamin D, however, were included in a few, as in Front Line COVID-19 Critical Care Alliance (FLCCC) protocols¹⁴⁸ and World Council of Health treatment guidelines.¹⁴⁹ Repurposed agents like ivermectin (war on ivermectin)^{141,142,150} have been addressed in detail elsewhere,^{150,151,119} and can be explored further on the website <https://c19early.org/>, which provides published clinical research data of all medications tested against SARS-CoV-2 and real-time meta-analyses.⁹⁷

Inference: Special interest groups and patient advocacy groups have propagated contradictory recommendations to favor their position, worsening the confusion and hindering the use of repurposed agents and vitamin D.¹⁵² Consequently, big pharma and governments exploit the mass confusion and the lack of unity among scientists to their advantage for marketing patented medications, like and vaccines and anti-viral agents.⁷⁹ Big Pharma has employed similar approaches, utilizing their misinformation playbooks.¹⁰³

3.0 Controlling the pandemic with vaccines

As previously illustrated, vaccines developed against the original SARS-CoV-2 had positive effects in 2021, flattening infectious peaks/outbreaks and distributing admissions to hospitals and intensive care facilities (ICUs) that reduced the burden.^{33,153} This reduced the overloading of health personnel and facilities, but this advantage diminished toward the end of 2021. This was due to the development of immune evasion by dominant SARS-CoV-2 mutants and the failure of bivalent vaccines.⁵¹ Due to the converging viral mutations occurring within the RBDs of Spike proteins and the consequent reduced immune recognition, the effectiveness of bivalent booster vaccines was reduced to below 40%.⁸³

3.1 Benefits derived from COVID-19 vaccines

Despite the failure to prevent the transmission of SARS-CoV-2,^{2,80,81} published governmental data during 2021 indicated that COVID-19 vaccines had an impact on reducing hospitalizations.¹⁵³ However, by early-2022, the hospitalization rates from Omicron variants of SARS-CoV-2 were increasing. There were few differences between fully vaccinated and boosted individuals and unvaccinated individuals.⁷⁹ Notably, the definition of fully vaccinated

varies from country to country. In the USA, one dose of Janssen (J&J) viral vector vaccine or two doses of mRNA COVID-19 vaccine constituted complete vaccination.

Several studies have reported the limited capacity and short duration of protective immunity^{154,155} derived from mRNA COVID-19 vaccines.^{153,156-158} As a result, the interval between vaccine booster doses was reduced from the intended yearly to six months to maintain circulating antibody levels.¹⁵⁸ However, the ratio of neutralizing antibodies to other antibodies declined, making the response to infection less effective. This decision, however, ignored the importance of T-cell priming and memory cell functions. Besides the short-lived immunity,¹⁵⁷ the efficacy of vaccines, including the Omicron-specific bivalent vaccines, is significantly less than it was against the original SARS-CoV-2 virus.¹⁵⁸

3.2 Fundamentals of controlling a pandemic and expectations

During an epidemic or pandemic, an essential public health concept is to prioritize the reduction of the incidence (peaks) of infections, i.e., by reducing the transmissibility (R_0) and curtailing viral spread from person to person.²⁶ This is vivid with traditional vaccines. An effective vaccine is expected to curtail the transmission of viruses or bacteria. SARS-CoV-2 transmission is primarily airborne, occurring via droplet spread.

Wearing facemasks, observing social distancing, avoiding crowded gatherings, and maintaining good personal hygiene are typical public health disease-preventive measures used against respiratory viral diseases/epidemics. These measures are designed to minimize exposure to high viral loads and curb the spread of the disease, thereby limiting infections and ultimately controlling viral transmission, as in the case of SARS-CoV-2. They aim to flatten infectious peaks, alleviate healthcare burdens, and eventually facilitate herd immunity.^{104,159} However, despite widespread vaccination efforts, these intended protective characteristics have not been fully realized.

Inference: As with all previous immunizations, the logical process and expected outcome were that vaccinated individuals would cease becoming infected by others and would not transmit the microbes to unvaccinated persons, such as patients. However, neither of these promises materialized with COVID-19 vaccines. Despite the false claims by Big Pharma that mRNA vaccines prevent the transmission of SARS-CoV-2, institutions like the Centers for Disease Control (CDC) in the USA, the World Health Organization (WHO), and prominent governments propagated this pharma-derived falsehood to boost vaccine uptake and sales. Not until late 2022 did the manufacturers of mRNA COVID vaccines, Pfizer and Moderna, acknowledge that their vaccines do not prevent the spread of the disease or reinfection.²

3.3 Original COVID-19 vaccines reduced hospitalization but failed to stop SARS-CoV-2 transmission

The COVID-19 outbreaks in 2020 and early 2021 primarily impacted specific vulnerable groups, especially older individuals with co-morbidities, immune-compromised or suppressed individuals, institutionalized persons, and ethnic minorities with darker skin living in temperate countries.¹⁶⁰ A common denominator among these susceptible groups was having a high prevalence of vitamin D deficiency.^{120,161,162} Given that approximately 75% of immune system activities rely on sufficient vitamin D,⁷³ COVID-19 became a pandemic among

those with severe vitamin D deficiency.⁷⁷ Despite the availability of this critical data, major healthcare agencies overlooked it⁷⁹ and took no action to advise the population on strengthening their immune systems to combat the SARS-CoV-2 virus.⁷⁹ Consequently, severe hypovitaminosis D persisted and worsened due to prolonged curfews and lockdowns in many countries.^{77,78} This contributed to millions developing severe complications requiring hospitalization and thousands succumbing to SARS-CoV-2.¹²⁰

Mass-scale COVID-19 vaccination commenced in late 2020, driven by WHO-enforced strategies adopted by most governments. The USA initially initiated vaccine mandates, which Europe and other nations subsequently replicated without thorough scientific scrutiny. They failed to conduct comprehensive independent investigations and adherence to published scientific data. Following governmental mandates, industrialized countries achieved successful vaccination coverage ranging from 60 to 80% of their adult populations.

Despite the mass-scale vaccination efforts, since early 2022, SARS-CoV-2 outbreaks and local epidemics have been resurgent, occurring irrespective of vaccination status. These instances have been observed in communities with fully vaccinated, boosted, and unvaccinated individuals. Despite widespread media promotion of vaccination, these local epidemics have shown similar spread patterns among vaccinated and non-vaccinated individuals, irrespective of factors such as access to healthcare, availability, or per capita spending.^{2,3,80,81}

3.4 Original SARS-CoV-2 vs. Omicron variants—the status and outbreaks

The Omicron variant of SARS-CoV-2 exhibited a rapid spread, surpassing the infectivity of the Delta variant with a doubling of infections every three days. Despite its swifter transmission, Omicron presented as milder, parallel to flu or influenza, and displayed different symptomatology than the original SARS-CoV-2. Irrespective of the vaccination status, this milder nature resulted in healthy individuals typically not requiring hospitalization. However, individuals with uncontrolled coexisting co-morbidities and those with severe vitamin D deficiency [serum 25(OH)D concentration less than 12 ng/mL] were still at an elevated risk of developing complications and death.¹⁶⁰ Unlike earlier SARS-CoV-2 variants like Alpha and Delta, the two preceding lineages of Omicron (BQ and XBB) that dominated outbreaks in 2022/2023 exhibited increased resistance to mRNA-derived neutralizing antibodies. This resistance (immune evasion) was more noticeable with bivalent booster-generated antibodies.^{82,85}

New Omicron variants displayed a notable shift in affecting younger age groups compared to the Alpha and Delta variants. This change was attributed to the acquired immune resistance of Omicron variants, driven by modified Spike proteins, leading to significant deviations in their immune characteristics from the original SARS-CoV-2. This altered immune profile contributed to immune evasion against mRNA and adenovirus vector-based COVID-19 vaccines^{163,164} and antiviral agents.⁸⁷ Consequently, vaccine-derived immunity proved less effective in preventing infections caused by new mutants, even with the administration of booster doses.¹⁶⁵⁻¹⁶⁷ Immunoparesis resulting from repeated booster doses contributed to reinfections in individuals previously exposed to SARS-CoV-2,^{2,153} particularly impacting older adults.^{66,158}

Figure 2 illustrates the antigenic drift from the original SARS-CoV-2 and the Omicron variant to XBB and XBB.1. This suggests that these subvariants behave antigenically

distinct or, perhaps, as a new virus compared to SARS-CoV-2.⁸² This observation may elucidate why the bivalent vaccine and monoclonal antibodies fail to recognize new subvariants, contributing to the reduced effectiveness of current COVID-19 vaccines and an increased occurrence of breakthrough infections and reinfections.^{168,169}

< Figure 2 >




Figure 2: Compared to pseudo-typed D614G and Omicron subvariants BAs, neutralization of Omicron subvariants BQ.1, BQ.1.1, XBB, and XBB.1c, by sera from five different cohorts—the limit of detection of one hundred (indicated in a dotted line). Values above the bold symbols represent geometric mean, ID₅₀ values (details of the experiment are in the original manuscript) (from Wang et al., Lancet, 2023).⁸²

Inference: Repetitive administration of COVID-19 booster doses further weakened the immune system, resulting in diminished responses against pathogenic microbes. This immune suppression, akin to vaccine-acquired immune deficiency syndrome,³⁰ is also called immune paresis, a concern that health authorities should have addressed early.

3.5 Omicron variants and post-COVID syndrome

Mutations represent natural biological phenomena that heighten environmental pressures for viral survival. Increased utilization of COVID-19 vaccines, especially among immunocompromised individuals and those with vitamin D deficiency, along with administering bivalent mRNA vaccine booster doses, offer limited immunity. This may have contributed to the rate of developing SARS-CoV-2 viral mutations.⁸⁴ The dominant variants, such as Delta and Omicron, have higher infectivity than the original SARS-CoV-2 as primarily derived from the mutations in the RBD region, which increased the affinity to the membrane-bound ACE-2 receptors on epithelial cells. However, the resistance to COVID-19 vaccines may also

have contributed to it.^{156,170} Additionally, convergent evolution has led to diverse combinations of mutations in the virus's RBD.⁸³

Although the dominant mutants of SARS-CoV-2 exhibit a higher rate of spread (R_0) in the community than the original SARS-CoV-2 virus,⁸⁴ the mortality remained low. Nevertheless, the occurrence of the post-COVID syndrome (long-COVID) is similar,¹⁵³ or even higher than those associated with the original virus.¹⁷¹ Due to the uprising and fatigue of the public due to enforced restrictions, most countries relaxed public health measures, such as wearing facemasks and restrictions on crowded gatherings, way before the WHO's declaration that the pandemic was over. However, some governments continued to propagate expensive COVID-19 vaccines. The Omicron variant outbreaks and endemic spread were fueled by increased local transmission and global travel, dominated by variants like B.1 and XBB mutants, which significantly evade immune recognition by vaccine-derived antibodies.^{154,163}

Inference: Globally, few or no new SARS-CoV-2 mutants originate from countries or communities with minimal vaccine uptake.⁷⁰ Most mutants occur in immune-deficient or fully vaccinated individuals.⁸² Moreover, more boosters correlate with an increased likelihood of mutations and reinfection.^{172,173} Conversely, repeated vaccine boosters weaken the immune system, making individuals more vulnerable to SARS-CoV-2 mutants,⁶⁷ reinfections.^{172,173} It also increased the risks for re-infections with mutant viruses and other dormant and commensal organisms, such as *Mycobacterium tuberculosis* and herpes zoster.¹⁷⁴

3.6 Parallels with Influenza vaccines

Each year, the characteristics of influenza viruses change. Consequently, vaccines are developed based on predictions of the predominant species in the upcoming winter season. However, these predictions are often inaccurate, resulting in flu vaccines being 'generic' and not specific to the correct species dominating during the following year. As a result, the efficacy of flu vaccines in preventing symptomatic disease, hospitalization, and deaths remains low.¹² This low efficacy has led to distrust in the product, and more adults are now reluctant to take flu vaccines.⁷⁰ A similar trend began to emerge in 2022 with SARS-CoV-2.¹⁷⁵ Initially, the uptake of bivalent vaccines was high, but as emerging data became available, their population uptake rapidly decreased despite offering incentives.^{110,158,175-177}

Recent scientific reports indicate that booster vaccines are relatively ineffective and may not be a solution to control the pandemic (Figure 2).^{110,158,175-177} Safety concerns regarding serious adverse effects related to COVID-19 vaccines are also growing.^{21,104,106,111,178} The mishandling of COVID-19 by public health authorities has further contributed to the loss of public trust. With increased infectivity from new variants and the diminishing efficacy of bivalent vaccines coupled with immune evasion by mutant SARS-CoV-2 viruses, controlling outbreaks has become challenging, resulting in a transition from a pandemic to an endemic. Figure 3 illustrates the effectiveness of mono and bivalent COVID-19 vaccines and their susceptibility to immune escape.

< Figure 3 >




Figure 3: Absolute effectiveness of vaccines [95% Confidence Intervals (CI) presented in parenthesis, in Y-axis] against symptomatic SARS-CoV-2 infection for a single bivalent mRNA COVID-19 booster dose given after 2, 3, or 4 doses of monovalent vaccine (no data presented for single dose recipients) compared with no booster dose. Data presented by age group (X-axis) vs. number of monovalent COVID-19 vaccine doses received. The average effectiveness of 24% is presented as a broken horizontal line. The figure was drawn using the data from the CDC's increasing community access to the testing program, USA (Sept.–Nov. 2022, table 3).¹⁷⁹

Inference: Research, including data from the CDC (Figure 3), demonstrates a decline in vaccine effectiveness against severe COVID-19, particularly after the second booster.^{110,179} As the efficacy of COVID-19 vaccines approaches levels comparable to placebos,¹⁷⁹ health authorities should have explored complementary options to prevent hospitalizations from Omicron variants using other means.

4.0 Development of Mutants and Rationale for Reduced Effectiveness of COVID-19 Booster Vaccines

4.1 Widespread vaccination is associated with increased SARS-CoV mutants

Variants such as Alpha, Beta, Delta, and Omicron exhibit mutations, deletions, and alterations in the sequence of the NTD antigenic supersite, a larger glycan-free region in the spike protein within the viral membrane.¹⁵⁶ Deletions in variants B.1.1.523 were identified in early 2021, while B.1.617.2 includes the E484K mutation in most variants.^{82,180,181} The E484K mutation is situated within the membrane RBD region, enhancing affinity to ACE-2 receptors and modifying immune detection.

The E484K mutation was also found in B.1.351 and P.1 variants, reducing vaccine efficacy by elevating vaccine resistance. This mutation also diminishes the effectiveness of monoclonal antibodies and convalescent plasma therapy, making them ineffective.¹⁸¹

Although these new variants evade immunity without increasing virulence, emergent variants like BQ.1.1 and XBB1.5 and future variants might lead to increased virulence.^{87,164,182} Consequently, the potential impact of future mutations should be a significant concern.⁸⁵

4.2 Viral infections and vaccinations consume vitamin D

The escalation of infection-induced vitamin D deficiency hinders the entry of D₃ and 25(OH)D into target cells, preventing the intracellular production of 1,25(OH)₂D in immune cells.^{74,127} This, in turn, disrupts the intracellular autocrine and paracrine signaling functions of calcitriol and activation of VDR (CTR) molecules.^{73,77} Maintaining a consistent D₃ and 25(OH)D supply can overcome this challenge and support a robust immune system. Studies have indicated that vaccinating individuals previously infected and recovered does not confer additional benefits against SARS-CoV-2 or its variants.¹⁸³ Furthermore, there is no discernible advantage to administering more than one booster shot to individuals fully vaccinated or recovered from SARS-CoV-2 infection.¹⁶⁷

The robust immune responses triggered by SARS-CoV-2 infections and following vaccinations increased vitamin D/25(OH)D utilization. Without continuous vitamin D supplementation during an infection, there is a risk of developing or existing vitamin D deficiency. The administration of more vaccines, particularly booster doses, correlates with a higher rate of depletion of vitamin D stores in the body and an accelerated reduction in serum 25(OH)D concentrations.^{73,184}

Additionally, zinc and magnesium, which are crucial cofactors for vitamin D's effects on VDR/CTR and other immune functions, are also depleted during this process.¹⁸⁵ The increased consumption of vitamin D exacerbates insufficiencies in zinc and magnesium.⁷⁴ Deficiencies in these components result in a dysfunctional innate immune system,⁵ and immune paresis,³⁰ contributing to delay in recovery, the spread of microbes, and increased complications and deaths. Several datasets indicate a positive association between the number of COVID-19 booster doses administered in the population or individuals and increased infection rates (Figure 4).^{163,166,167,170,186}

< **Figure 4** >




Figure 4: A Simon-Makuch plot illustrates data from healthcare workers at Cleveland Clinic (n=51,011) USA—comparisons of the risk of acquiring COVID-19, with and without COVID-19 vaccines and booster doses. Data were analyzed using Cox proportional hazards regression, point estimates, and 95% confidence intervals. The risk of COVID-19 infection linearly increases with the number of COVID-19 booster doses (adapted from Shrestha, NK et al. with permission; pending peer review).⁶⁷

The gradual decline in the efficacy of COVID-19 vaccines and bivalent boosters is attributed to many factors. These include mutations that lead to immune evasion by viruses, repeated booster doses causing immune exhaustion and depletion of immune cells, and the increasing deficiency of vitamin D and cofactors, resulting in immune paresis, etc., analog to vaccine-acquired immune deficiency syndrome.³⁰ The weakening of the immune system is particularly evident, as depicted in Figure 4, after administering the second booster dose of COVID-19 vaccines.

Inference: Deficiencies in vitamin D and cofactors substantially compromise innate and post-immunization immune responses,¹⁶⁷ rendering individuals more susceptible to SARS-CoV-2 reinfections,^{172,173} infections from mutant viruses,⁶⁷ and other commensal organisms.¹⁷⁴ Collectively, these factors contribute to heightened morbidity and mortality.

4.3 Development of immune evasions by Omicron variants

Several recent studies have documented the diminishing effectiveness of COVID-19 vaccines.^{2,3,67,172,173} An RCT conducted at Cleveland Clinic in the USA involved 51,011 healthcare workers who received bivalent COVID-19 vaccines.⁶⁷ 13 weeks follow-up data revealed an increased risk of contracting Omicron variants with the number of vaccine booster doses received.^{67,167} The data suggests that the higher the vaccine booster doses, the higher the risk of contracting Omicron disease, as illustrated in Figure 4.

Additional booster doses decrease immune capabilities and heighten vulnerability to SARS-CoV-2,⁶⁷ reinfections,^{91,172,173} and other infections,¹⁷⁴ partially attributed to immune paresis.¹⁶⁵ The effectiveness of current bivalent COVID vaccines in shielding against infection in the BA.4/BA.5 lineages of the Omicron variant is reported to be approximately 30%, akin to a placebo.^{67,167} Strikingly, each successive booster dose correlates with a step-wise escalation in susceptibility to SARS-CoV-2 infection, as depicted in Figure 4.

4.4 Actions that may help to reduce poor outcomes in future pandemics

Taking preventive measures such as micronutrient sufficiency could naturally strengthen the immune system against microbes and prevent vaccine-associated immunoparesis and the incidence of post-COVID syndrome among both vaccinated and unvaccinated individuals.¹⁶⁵⁻¹⁶⁷ Given the ongoing viral mutations, exploring alternative strategies beyond vaccines alone is imperative. Early therapies incorporating unpatented, repurposed, and widely available agents like vitamin D, ivermectin, and melatonin should be considered.⁹⁷

The <https://c19early.org/> website provides information on over forty generic agents investigated and reported in peer-reviewed literature.⁹⁷ The first scientific publication recommending an effective remedy against SARS-CoV-2—vitamin D—was released in February 2020.¹⁸⁷ The failure to diversify the therapeutic options, influenced by conflicts of interest, pharmaceutical bias, and administrative pressures to maintain the Emergency Use Authorization (EUA) status of patented vaccines and antiviral drugs, resulted in chaos. Replicating such a flawed approach could have disastrous consequences in future pandemics and should be avoided.

Molecular biological techniques offer a practical approach to combating viruses with rapid mutation, aiming to understand and detect them at an early stage. However, it is crucial to identify the complete viral antigenic profile and employ a broader antigenic exposure strategy to induce immunity and antibodies against a broader antigenic spectrum, sensitizing memory cells. The production of such therapies must rapidly address viruses that mutate within a year. Despite the advantages of mRNA technology deployment, the serious adverse effects of Spike protein toxicity and the rapid loss of efficacy raise concerns, making it questionable to use similar choices and technology in the future. As the current mRNA technology is still evolving, introducing insufficiently tested products directly into humans via vaccine injections¹⁸⁸ or orally¹⁸⁹ without establishing proper safety profiles was deemed premature and likely to have serious negative consequences.

4.5 Reported serious adverse events with mRNA vaccines

In the Moderna randomized controlled trial (RCT), serious adverse effects (SAEs) occurred at a rate of 15.1 per 10,000 participants, compared to 6.4 per 10,000 participants in the placebo group—a 2.4-fold higher risk of SAEs than the reduction observed in COVID-19 hospitalization.¹⁹⁰ Similarly, in the Pfizer RCT, the excess risk of SAEs was 10.1 per 10,000 participants, relative to 2.3 per 10,000 in the placebo group¹⁰—a 4.4-fold higher incidence of SAEs compared to the risk reduction for COVID-19 hospitalization.^{10,191}

Combined mRNA vaccine data was associated with an excess risk of SAEs of 12.5 per 10,000 vaccinated (95 % CI 2.1 to 22.9)—a risk ratio of 1.43 (95 % CI 1.07 to 1.92). The Pfizer RCT had a 36 % higher risk of SAEs in the vaccine group with a risk difference of 18.0

per 10,000 vaccinated (95 % CI 1.2 to 34.9)—a risk ratio 1.36 (95 % CI 1.02 to 1.83), a similar trend.¹⁰ The SAEs and risk ratios in these RCTs are significantly higher in the mRNA-vaccinated groups than in the placebo groups,^{190,191} but the regulators overlooked these data. Independent scientists' lack of access to primary data further hindered impartial analysis.

4.6 Booster doses: SARS-CoV-2 spread, duration of protection, and impact on mortality

Data confirmed that vaccinated and unvaccinated could harbor, shed the virus, and infect others equally.^{2,80,81,163} The primary series of COVID-19 vaccines partially protects from complications, thus reducing hospitalizations, but only for a few months.^{33,155,192} Neither antiviral agents nor monoclonal antibody therapies activate the immune system or provide longer-term benefits. These treatments typically cost \$500 to \$2,000 per patient. Nevertheless, they exhibit lower effectiveness than generic agents, such as vitamin D and ivermectin, which treatment costs around two dollars per person.¹² Calcifediol is more effective than vitamin D, which costs about ten dollars per patient. Many published clinical studies in combating SARS-CoV-2 infection. Clinical studies using generic agents against SARS-CoV-2 infection real-time meta-analyses and summarized data available on c19early.org/dmeta.html support these concepts.⁹⁷

Given the failure of vaccines to prevent the transmission of SARS-CoV-2,^{91,92 48,49,59 2,3,67,172,173} relying on vaccination and mandating status (proof of vaccination) cards or digital identities and segregating people lacks scientific merit. Even in countries like the USA, the UK, Israel, etc., where over 75% of adults and adolescents are vaccinated, achieving herd immunity or halting viral spread failed.^{159,193} The persistence of outbreaks was attributed, in part, to transmission within communities and introduction from outside.

The challenges are compounded by the evasion of vaccine-induced immunity by dominant new variants like the XBB series, along with a reduction in the ability of antibodies to neutralize viruses to under 40%.^{154,163} With emerging new variants developing resistance to vaccines and antiviral agents,⁸⁷ the reliance on vaccinations alone would fail to prevent COVID-19 outbreaks, establish herd immunity, or eliminate SARS-CoV-2 and its mutant viruses.

In certain Western countries, the proportion of hospitalization attributed to Omicron remains lower among the vaccinated. However, after the introduction of bivalent vaccines, this ratio changed; the reasons for this were discussed above.¹⁹⁴ to the high vaccination rates in industrialized countries, the absolute numbers of infections and re-infections, as well as hospitalization, became higher among the fully vaccinated than the unvaccinated.¹⁹⁵ Making firm conclusions requires appropriately adjusting for confounding factors and denominators. For instance, a higher proportion of the vaccinated population comprises older people with co-morbidities. Cautionary measures are warranted unless the observed trend is consistent among younger individuals. Figure 5 illustrates broader effects using an acute increase in reported sudden deaths in Germany in 2021 and provides comparative data on vaccine uptakes and deaths from 2021-22 globally, in Sri Lanka and Israel.

< **Figure 5** >




Figure 5: A) illustrates the change in reported deaths in Germany between 2021 and 2022. Figures C through D depict the parallel relationship between the daily administration of COVID-19 booster vaccine doses and reported deaths. Figure B presents global data, showcasing the daily number of vaccine booster doses administered per 100 people and deaths from COVID-19 (sourced from OneWorld in Data, Human Mortality Database, and World Mortality Dataset, 2023). Figures C and D also provide specific data for Sri Lanka and Israel, respectively. In Figure C, daily vaccination statistics are compared to daily reported deaths from COVID-19 in Sri Lanka from January to September 2021 (adapted from E. Dias). In Figure D, Israel's data illustrates the number of persons fully vaccinated versus daily deaths from COVID-19 from December 2020 to September 2021 (source: <https://our-worldindata.org/>), highlighting the observed parallelism.

As highlighted earlier, increased complications and hospitalizations can be attributed partly to the immune evasion exhibited by dominant mutant viruses and the immunoparesis of hosts resulting from booster doses or repeated SARS-CoV-2 infections.^{163,166,167,170,186} This pattern has been observed previously with other viruses, especially in individuals who are vitamin D deficient or immune-compromised.¹⁹⁶

Inference: Consequently, advocating for multiple booster doses and vaccinating younger children lacks rationale and merit. Instead, the primary focus of vaccination should have been on vulnerable populations to enhance their adaptive immune system, activate memory cells, and prioritize the maintenance of a robust immune system for effective defense

against the virus. This becomes particularly evident when considering data demonstrating declining mortality rates during ivermectin treatment campaigns.¹⁹⁷

4.7 Countries with fewer than 15% vaccinated had no new COVID outbreaks

Approximately 2.8 billion people within 1400 km of the equator experience sufficient sunlight exposure throughout the year. Most individuals in this population maintain their serum 25(OH)D concentrations above 30 ng/mL,^{74,126} the minimum necessary for certain biological activities, except those deliberately avoiding sun exposure. However, the average serum population serum 25(OH)D concentration in most countries is about 20 ng/mL, including in the USA,¹⁹⁸ and most developing countries and emerging economies.

The countries mentioned above had insufficient dollars to purchase the required COVID-19 vaccines. With this limited access, their COVID-19 vaccine uptake was between 5 and 15%; the majority only received the first dose. In contrast, over 70% of adults in industrialized nations were fully vaccinated, with a similar percentage having received booster doses. Despite these disparities and healthcare inequalities, the tropical countries reported fewer or no COVID-19 outbreaks, hospitalizations, and fewer deaths from 2022 onwards.

Furthermore, adults in affluent countries received multiple booster vaccine doses, around three boosters per person, yet exhibited lesser protection against SARS-CoV-2, evidenced by re-infections.¹⁹⁹ This significant disconnect is highlighted in Figure 6, which illustrates the polarized SARS-CoV-2-related deaths expressed per million population in industrialized countries (characterized by higher GDP and per capita health expenditure) with rates of high vaccination (Cluster 1) compared to developing countries with low per capita GDP and low vaccination rates at the time (Cluster 2). As illustrated in Figure 6, affluent countries (Cluster 1) experienced a four-hundred-fold higher death rate from COVID-19 than developing countries.

< Figure 6 >




Figure 6: The data on deaths from COVID-19, spanning from November 2021 to February 2023, presented as the rolling seven-day average deaths per million population, clearly illustrates a stark 400-fold difference in death rates between the two clusters. Cluster 1 comprises industrialized countries with over 70% of the population fully vaccinated, while Cluster 2 represents African countries with less than 15% vaccination coverage. Notably, India and Sri Lanka achieved higher vaccination percentages in late 2022, and their death rates were already low before the initiation of vaccination campaigns. The data is expressed as deaths per million population and adapted from Our World COVID data based on the Johns Hopkins University database.

4.8 The cost-effectiveness (NNT) of various agents examined against SARS-CoV-2

The website <https://c19early.org> is a reliable and comprehensive resource featuring over a hundred peer-reviewed clinical research articles focused on three prominent treatments for SARS-CoV-2: vitamin D and ivermectin. {Group-C19.com, 2022 #17141} This unique site independently compiled data from diverse clinical research teams worldwide, providing individual research study data and collective analyses over forty-five intervention agents published against SARS-CoV-2. This site analyzed individual published clinical studies—305 with vitamin D and 99 with ivermectin, consisting of several thousand study participants (i.e., reliable big data sets consisting of SARS-CoV-related 4,423 clinical studies). {Group-C19.com, 2022 #17141} These two agents emerged as highly cost-effective options. The estimated costs (also imply the number needed to treat; NNT) are reported as \$11 for vitamin D and \$25 for ivermectin (Table 1).

High Cost-Effectiveness				Low Cost-Effectiveness			
Agent	Cost	Agent	Cost	Agent	Cost	Agent	Cost
Melatonin	\$8	Aspirin	\$33	Fluvoxamine	\$411	Casirivimab	\$203958
Vitamin D	\$11	Curcumin	\$59	Budesonide	\$574	Paxlovid	\$206705
Alkalization	\$11	Famotidine	\$94	Nitazoxanide	\$680	Bamlaniv	\$301549
Zinc	\$15	Probiotics	\$99	Azvodine	\$1248	Sotrovimab	\$325800
Ivermectin	\$25	Quercetin	\$127	Favipiravir	\$1934	Bebtelovimab	\$737601
HCQ	\$26	Metformin	\$133	Tixagev	\$74506	Remdesivir	\$1558440
Colchicine	\$26	Nigella Sativa	\$187	Regdanvimab	\$139860	Molnupiravir	\$2400867

Table 1: The cost for a life saved (i.e., cost-effectiveness) of several tested agents in SARS-CoV-2*

*The cost of an agent needs to save one life) for SARS-CoV-2 infection. Low-cost but low-efficacious agents were excluded (<https://c19early.org/>). {Group-C19.com, 2022 #17141}

In addition to the two agents mentioned above (vitamin D and ivermectin), there are other generic agents with reported clinical significance: Fluvoxamine (an antidepressant): 20 clinical studies reported a 40 to 80% reduction in hospitalization;^{200,201} hydroxychloroquine, 27 clinical studies reporting 80% reduction in hospitalization; Elderberry (from European black elder - *Sambucus nigra*): Demonstrated a 50% improvement;²⁰² Curcumin (oral nano-curcumin): Collective efficacy of 26 clinical studies with a significant p-value of 0.0000093;²⁰³ Nigella (*Nigella sativa*, black cumin seed): 12 clinical studies showing over 50% reduction in hospitalization;²⁰⁴ Melatonin 11 clinical studies with a 50% reduction in mortality;²⁰⁵ Quercetin: 11 clinical studies showing over 50% reduction in hospitalization.²⁰⁶ All agents were reported to have clinically significant anti-viral effects.

In contrast, the anti-viral agent favipiravir, with 67 studies, reported a pooled effect of ~20% reduction of risks,²⁰⁵ Molnupiravir with ~20% lower risks²⁰⁷ (8% reduction of post-COVID syndrome),²⁰⁸ Paxlovid (*nirmatrelvir-ritonavir*) ~50% reduction of risks²⁰⁹ (1% improvement of post-COVID syndrome/complications,²¹⁰ and Remdesivir, 62 clinical studies, 0-10% Improvement²¹¹ and higher mortality.²¹² The pooled effects and mean outcomes were obtained from <https://c19early.org/ader.html>.⁹⁷

4.9 Ivermectin as a cost-effective alternative therapy

Ivermectin is a broad-spectrum antiparasitic drug with additional antiviral activities.²¹³⁻²¹⁵ While several negative studies on ivermectin were published, many were funded by pharmaceutical companies and biased. Some were designed to fail, indicating conflicts of interest. Over a hundred clinical studies and 50+ RCTs have investigated ivermectin—316 clinical trials in 35 countries involving 446,237 subjects with COVID-19. Besides, some RCTs reported a strong association and a significant reduction of mortality with ivermectin use (RR, 0.25; 95% CI, 0.09-0.70; p = p.008);²¹⁴ other analyses support this.²¹⁶

4.9.1 Ivermectin positive studies

Recently published positive clinical studies include seven meta-analyses and a systematic review of trials conducted in 2020 (n=382); Ragó et al. reported significant benefits associated with ivermectin.²¹⁷ These include faster viral clearance, an 81% reduction in symptomatic disease, 38% fewer hospitalizations, and a 49% decrease in mortality²¹⁷. Another meta-analysis, encompassing 15 clinical trials (n = 2438), indicated that prophylactic use of ivermectin resulted in an 86% reduction in COVID-19 infections (95% confidence interval 79%–91%) and a 62% decrease in mortality risk compared to placebo confirmed by Trail Sequential analysis (95% CI 0.19–0.73).²¹⁸

Several studies have also affirmed the safety and cost-effectiveness of ivermectin when used at doses ranging from 0.2 to 0.4 mg/kg^{219,220}. Additionally, RCT by Varnaseri et al. (n=110) showed ivermectin significantly reduced ICU admissions (32.7% vs. 5.5%; p<0.001), hospitalization duration (6 vs. 4 days; p<0.001), and median time to symptom resolution (p<0.05) compared to the placebo group, with no serious adverse effects reported²²¹.

Recent studies on ivermectin were controversial, but well-designed RCTs provided compelling data. The PRINCIPLE study by Hayward et al. (n=3,963) revealed a 36% reduction in COVID-19-specific symptoms (p<0.0001), with ivermectin demonstrating superiority in primary recovery outcomes with a probability of superiority >0.999.²²² Early administration

of ivermectin resulted in faster recovery, complete symptom alleviation, and sustained relief ($p < 0.0001$), all at a significantly lower cost than other treatments, such as Molnupiravir, which can cost over \$700 per patient.

Ivermectin has been the subject of 103 peer-reviewed clinical studies, including 50 RCTs; 142,417 subjects participated and were contributed by 1,193 scientists. Real-time meta-analyses of these studies reported significant efficacy ($p < 0.00000001$) in reducing the risk of infection, hospitalization, and mortality and increased viral clearance. A random pooled analysis of 50 RCTs reported that ivermectin demonstrated 85% effectiveness in reducing the risk of COVID-19 and 49% efficacious in reducing mortality in early treatment studies.

4.9.2 *Negative studies using ivermectin*

Nevertheless, there were several negative studies using ivermectin. The TOGETHER study by Reis et al. ($n=679$) reported no significant difference between ivermectin and placebo groups in the risk of hospitalization or emergency department visits for COVID-19.²²³ ACTIV-6 study by Naggie et al. used (400 ug/kg for three days) found no significant benefit with ivermectin compared to placebo on time to sustained recovery (defined as three consecutive days without symptoms) (hazard ratio 1.07, 95%; CI 0.96–1.17).

Nevertheless, TOGETHER²²³ and ACTIV-6²²⁴ RCTs failed to demonstrate significant benefits using doses of ivermectin. These studies had major design errors and potential bias, raising doubts about the validity of conclusions (<https://c19ivm.org/>). Similarly, the COVID-OUT study by Bramante et al. ($n=1323$) initially reported no effect from ivermectin, metformin, or fluvoxamine.²²⁵ However, later analyses suggested a notable reduction in hospitalizations exceeding 50%.²²⁶

Nevertheless, some evidence on ivermectin in preventing SARS-CoV-2 infection is contradictory.^{213,214,226-229} In contrast, well-designed trials with no conflicts of interest showed substantial benefits.^{11-13,214,216,230} These conflicting findings underscore the importance of careful interpretation and further investigation. Based on local research, ivermectin has been approved and used widely in over forty countries to prevent and treat SARS-CoV-2.²²⁸

Some of these negative studies were placebo-controlled RCTs. These have been used in a few systematic reviews and meta-analyses that reported equivocal outcomes,^{213,215,227,228,231,232} and from large, placebo-controlled outpatient trials conducted in the USA (COVID-OUT and ACTIV-6 (two trials, both evaluating a similar dose to this study of 400 $\mu\text{g}/\text{kg}$ for three days,^{225,226} and ACTIV-6 additionally evaluating 600 $\mu\text{g}/\text{kg}$ for six days,²²⁹ and TOGETHER study, 400 $\mu\text{g}/\text{kg}$ for three days.²²³ However, subjects in these studies were heterogeneous, and critical study group characteristics like vaccination status and comorbidities (e.g., high BMI) were not comparable and higher in the interventional groups²²⁵. None of the RCTs mentioned above reported any safety concerns regarding ivermectin.

The Cochrane Reviews encompass all negative studies, including those with significant study design flaws, which dilute potential benefits from investigational agents, such as ivermectin. This phenomenon is not unique to ivermectin but has also been observed in previous Cochrane reviews evaluating other agents. It underscores an inherent limitation of Cochrane reviews.

Inference: Despite a few negative studies, primarily employing improper doses in late-stage disease, the overall data strongly support positive clinical outcomes when ivermectin is used in the early stages of COVID-19 (early therapy) with appropriate doses. Both vitamin D and ivermectin are low-cost, readily available generic agents that have been shown to have no adverse effects at recommended doses. Therefore, these treatments should be accessible for prophylactic use and as adjunct therapies against SARS-CoV-2.

4.10 Consensus on vitamin D and ivermectin—implications for mRNA vaccines

A summary of expert consensus from c19early.org highlights the website as a comprehensive resource providing data from all published clinical studies (positive and negative) on various therapies combating SARS-CoV-2.⁹⁷ The site distinguishes itself as a reliable, dynamic real-time meta-analysis, offering an evolving overview of collective findings. Additionally, it presents calculated evidence on the cost-effectiveness of multiple treatments, incorporating NNT metrics to save one life. The website elucidates the cost-effectiveness of each intervention in terms of US dollar expenditure per life saved, offering valuable insights across diverse therapeutic approaches (Table 1).

As detailed in previous references, the reported negative studies remain below 10%. These negative studies consistently revealed significant shortcomings in their study designs and flawed statistical methods used.⁷⁹ Metanalyses of these treatments demonstrated a high level of significance ($p < 0.0000000001$), with an overall true efficacy (not relative efficacies as reported with COVID-19 vaccines), approximately 50% in preventing hospitalizations and deaths for each agent.⁹⁷ Sufficient cost-efficacy and safety data supporting these treatments were available by late 2020, warranted regulatory approval for clinical use, but never materialized.⁷⁹

In contrast to the documented safety records of mRNA vaccines,^{51,233} vitamin D and ivermectin, mentioned above, exhibited decades of safety and efficacy in managing infections during the pre-pandemic,²³⁴⁻²³⁶ and post-pandemic era.²³⁷⁻²³⁹ Despite their established efficacy and safety track record, western health authorities opted for experimental methods with little safety data, influenced by lobbying from the pharmaceutical industry. These contributed to suppressing evidence that supported early treatments that could have saved lives and reduced ICU occupancy and healthcare costs. NIH analyzed vitamin D for COVID-19 and concluded there is insufficient evidence to recommend it.²⁴⁰ This 2024 update from the NIH is despite 316 clinical trials published, including over 50 RCTs. However, it overlooked 86% of the published RCTs but included many failed studies, suggesting a significant selection bias.

Inference: Regulators did not consider alternative approaches to COVID-19 vaccines and antiviral agents for the above reasons. Relaxation of patents related to mRNA vaccine technology and allowing raw material availability during the early COVID-19 emergency would permit countries like India and China with well-established traditional vaccine technology and facilities to develop vaccines at a significantly lower cost.

Despite taking longer, adopting conventional vaccines would have significantly reduced production costs, allowing simultaneous global production at an affordable cost and increased availability of COVID-19 vaccines. It provided the opportunity to thoroughly investigate mRNA technology for its safety and efficacy independent of Pfizer and Moderna

companies and test for a broader range of viral antigens before generating and widespread deployment. Considering the serious adverse effects,^{49,50} and the inability of mRNA vaccines to prevent infections and the spread,^{91,92,48,49,59,2,3,67,172,173} funding and focusing only on mRNA vaccines may have been an error made in haste.

5.0 Conclusions

The mRNA and adenoviral vector vaccines were developed only against the Spike protein sequences of SARS-CoV-2.¹⁸ The bivalent vaccines combined sequences of the original viral RBD sequences and Omicron mutants.^{106,241} In either case, the immune recognition regions were limited and targeted only parts of the RBD portions of the Spike protein, omitting all of the core proteins. Furthermore, the resultant antibodies had no secretory mucosal immunity protection. Continuing mutations of the Spike protein led to new mutants escaping the host's immune recognition (by neutralizing antibodies), creating immune evasion.⁸³ In addition, booster doses caused immunoparesis (i.e., further weakening of the host's immune system),¹⁶⁵⁻¹⁶⁷ thereby increasing the vulnerability to SARS-CoV-2 infections, re-infections,¹⁷³ and re-activation of dormant viral and bacterial infections with a secondary flare-up of infections.

Consequently, despite mass vaccination programs covering approximately 5.6 billion people globally, predominantly from industrialized (affluent) countries, hospitalizations and deaths from SARS-CoV-2 and its mutants continue, though at a lower rate. None of the COVID-19 vaccines failed to prevent SARS-CoV-2 infections or dissemination of infection in vaccinated people. Due to the changing characteristics and behavior of mutant viruses, such as immune evasion, the efficacies of the original mRNA vaccine and the bivalent vaccines against the original SARS-CoV-2 virus and Omicron variants are less than 40%. Furthermore, COVID-19 vaccines, as well as boosters, were associated with severe adverse effects.

Therefore, widening the armamentaria for prevention and treatment became essential, especially before the next round of dominant SARS-CoV-2 mutants. Approximately four hundred clinical studies have reported that previously approved repurposed and globally available agents, such as vitamin D and ivermectin, available without a prescription, are cost-effective adjunct therapeutic options. However, regulators continued to be overlooked despite convincing clinical studies and RCTs demonstrating their effectiveness for vitamin D and ivermectin. Meanwhile, a few conflicted publications sponsored and funded by pharma attempted to discredit the efficacies of unpatented agents (e.g., Together, ACTIV-6, COVID-Out). These are illustrated at <https://c19ivm.org/meta.html> website.

An additional advantage of these generic agents is their broader efficacy against pathogenic microbes, including bacteria, viruses, and parasites, and their long history of safety. Unlike COVID-19 vaccines and patented antiviral agents, the effectiveness of this non-proprietary early therapy remains unaffected by viral mutations. Given the declining efficacy of all COVID-19 vaccines and the poor efficacy of antiviral agents as well as significant adverse effects, regulators and health agencies must promptly assess these non-patented, widely available agents. This evaluation should occur before the next round of dominant mutants emerge or the next pandemic. Additionally, it should be conducted in conjunction with appropriate

public healthcare measures, which may involve considering vaccines targeting the entire virus with new, traditional vaccines.

Conflicts of interest: The author declares no conflicts of interest. The author did not receive any funding or writing assistance.

Acknowledgments: The author appreciates the input from Drs. Athula Polonowita and Maneesha Weerasooriya.

Funding: None

Ethical approval: Not applicable

Ethics statement: This study does not have issues related to any ethical aspect.

Availability of data and materials: Data is included in the article/supp. material is referenced in the article.

Funding: None

Copyright: Not applicable (covered under CCBY license)

References

1. Springer DN, Bauer M, Medits I, et al. Bivalent COVID-19 mRNA booster vaccination (BA.1 or BA.4/BA.5) increases neutralization of matched Omicron variants. *NPJ Vaccines*. Aug 4 2023;8(1):110. doi:10.1038/s41541-023-00708-9
2. Singanayagam A, Hakki S, Dunning J, et al. Community transmission and viral load kinetics of the SARS-CoV-2 delta (B.1.617.2) variant in vaccinated and unvaccinated individuals in the UK: a prospective, longitudinal, cohort study. *Lancet Infect Dis*. Feb 2022;22(2):183-195. doi:10.1016/S1473-3099(21)00648-4
3. Bleier BS, Ramanathan M, Jr., Lane AP. COVID-19 Vaccines May Not Prevent Nasal SARS-CoV-2 Infection and Asymptomatic Transmission. *Otolaryngology--head and neck surgery : official journal of American Academy of Otolaryngology-Head and Neck Surgery*. Feb 2021;164(2):305-307. doi:10.1177/0194599820982633
4. Guo Y, Han J, Zhang Y, et al. SARS-CoV-2 Omicron variant: epidemiological features, biological characteristics, and clinical significance. *Front Immunol*. 2022;13:877101. doi:10.3389/fimmu.2022.877101
5. Charoenngam N, Holick MF. Immunologic effects of vitamin D on human health and disease. *Nutrients*. Jul 15 2020;12(7)doi:10.3390/nu12072097
6. Ahmadi S, Bazargan M, Elahi R, Esmailzadeh A. Immune evasion of severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2); molecular approaches. *Mol Immunol*. Apr 2023;156:10-19. doi:10.1016/j.molimm.2022.11.020
7. Kindler E, Thiel V. To sense or not to sense viral RNA--essentials of coronavirus innate immune evasion. *Curr Opin Microbiol*. Aug 2014;20:69-75. doi:10.1016/j.mib.2014.05.005
8. Yuan W, Pan W, Kong J, et al. 1,25-dihydroxyvitamin D3 suppresses renin gene transcription by blocking the activity of the cyclic AMP response element in the renin gene promoter. *J Biol Chem*. Oct 12 2007;282(41):29821-30. doi:10.1074/jbc.M705495200
9. FDA. Emergency use authorization for vaccines explained. FDA.gov. Accessed May 22nd, 2022. <https://www.fda.gov/vaccines-blood-biologics/vaccines/emergency-use-authorization-vaccines-explained>
10. Fraiman J, Erviti J, Jones M, et al. Serious adverse events of special interest following mRNA COVID-19 vaccination in randomized trials in adults. *Vaccine*. Sep 22 2022;40(40):5798-5805. doi:10.1016/j.vaccine.2022.08.036
11. Okumus N, Demirturk N, Cetinkaya RA, et al. Evaluation of the effectiveness and safety of adding ivermectin to treatment in severe COVID-19 patients. *BMC Infect Dis*. May 4 2021;21(1):411. doi:10.1186/s12879-021-06104-9
12. Wimalawansa SJ. Decoding the paradox: Understanding elevated hospitalization and reduced mortality in SARS-CoV-2 variants. *Int J Frontiers in Sci and Technol Res*. 2024;6(2):1-20. doi:10.53294/ijfstr.2024.6.2.0031
13. Shahbaznejad L, Davoudi A, Eslami G, et al. Effects of ivermectin in patients With COVID-19: A multicenter, double-blind, randomized, controlled clinical trial. *Clin Ther*. Jun 2021;43(6):1007-1019. doi:10.1016/j.clinthera.2021.04.007
14. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. Mar 29 2021;372:n71. doi:10.1136/bmj.n71
15. Heinz FX, Stiasny K. Distinguishing features of current COVID-19 vaccines: knowns and unknowns of antigen presentation and modes of action. *NPJ Vaccines*. Aug 16 2021;6(1):104. doi:10.1038/s41541-021-00369-6
16. Jain S, Venkataraman A, Wechsler ME, Peppas NA. Messenger RNA-based vaccines: Past, present, and future directions in the context of the COVID-19 pandemic. *Adv Drug Deliv Rev*. Dec 2021;179:114000. doi:10.1016/j.addr.2021.114000
17. Arashkia A, Jalilvand S, Mohajel N, et al. Severe acute respiratory syndrome-coronavirus-2 spike (S) protein based vaccine candidates: State of the art and future prospects. *Rev Med Virol*. May 2021;31(3):e2183. doi:10.1002/rmv.2183
18. Park JW, Lagniton PNP, Liu Y, Xu RH. mRNA vaccines for COVID-19: what, why and how. *International journal of biological sciences*. 2021;17(6):1446-1460. doi:10.7150/ijbs.59233
19. Yang H, Rao Z. Structural biology of SARS-CoV-2 and implications for therapeutic development. *Nat Rev Microbiol*. Nov 2021;19(11):685-700. doi:10.1038/s41579-021-00630-8
20. Yang L, Tang L, Zhang M, Liu C. Recent Advances in the Molecular Design and Delivery Technology of mRNA for Vaccination Against Infectious Diseases. *Front Immunol*. 2022;13:896958. doi:10.3389/fimmu.2022.896958
21. Stati G, Amerio P, Nubile M, Sancilio S, Rossi F, Di Pietro R. Concern about the effectiveness of mRNA vaccination technology and Its long-term safety: potential interference on miRNA machinery. *Int J Mol Sci*. Jan 11 2023;24(2)doi:10.3390/ijms24021404

22. Bouwman W, Verhaegh W, Holtzer L, van de Stolpe A. Measurement of Cellular Immune Response to Viral Infection and Vaccination. *Front Immunol.* 2020;11:575074. doi:10.3389/fimmu.2020.575074
23. Auladell M, Jia X, Hensen L, et al. Recalling the Future: Immunological Memory Toward Unpredictable Influenza Viruses. *Front Immunol.* 2019;10:1400. doi:10.3389/fimmu.2019.01400
24. Trougakos IP, Terpos E, Alexopoulos H, et al. Adverse effects of COVID-19 mRNA vaccines: the spike hypothesis. *Trends Mol Med.* Jul 2022;28(7):542-554. doi:10.1016/j.molmed.2022.04.007
25. Fang E, Liu X, Li M, et al. Advances in COVID-19 mRNA vaccine development. *Signal Transduct Target Ther.* Mar 23 2022;7(1):94. doi:10.1038/s41392-022-00950-y
26. Wimalawansa SJ. What modelling and reproduction numbers are useful in predicting COVID-19 spread? *Can J Biomed Res & Tech.* 2020;3(3):1-4.
27. Havasi A, Visan S, Cainap C, Cainap SS, Mihaila AA, Pop LA. Influenza A, Influenza B, and SARS-CoV-2 similarities and differences - A focus on diagnosis. *Front Microbiol.* 2022;13:908525. doi:10.3389/fmicb.2022.908525
28. Schlake T, Thess A, Fotin-Mleczek M, Kallen KJ. Developing mRNA-vaccine technologies. *RNA Biol.* Nov 2012;9(11):1319-30. doi:10.4161/rna.22269
29. Bellavite P, Ferraresi A, Isidoro C. Immune response and molecular mechanisms of cardiovascular adverse effects of Spike Proteins from SARS-CoV-2 and mRNA vaccines. *Biomedicines.* Feb 3 2023;11(2)doi:10.3390/biomedicines11020451
30. Seneff S, Nigh G, Kyriakopoulos AM, McCullough PA. Innate immune suppression by SARS-CoV-2 mRNA vaccinations: The role of G-quadruplexes, exosomes, and MicroRNAs. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association.* Jun 2022;164:113008. doi:10.1016/j.fct.2022.113008
31. Shahcheraghi SH, Ayatollahi J, Aljabali AA, et al. An overview of vaccine development for COVID-19. *Ther Deliv.* Mar 2021;12(3):235-244. doi:10.4155/tde-2020-0129
32. Cagigi A, Lore K. Immune responses induced by mRNA vaccination in mice, monkeys and humans. *Vaccines (Basel).* Jan 18 2021;9(1)doi:10.3390/vaccines9010061
33. Thompson MG, Stenehjem E, Grannis S, et al. Effectiveness of Covid-19 Vaccines in Ambulatory and Inpatient Care Settings. *N Engl J Med.* Oct 7 2021;385(15):1355-1371. doi:10.1056/NEJMoa2110362
34. Office-GAO-21-319 GA. Operation Warp Speed: Accelerated COVID-19 vaccine development status and efforts to address manufacturing challenges. GAO.
35. Zhang S, Yang Z, Chen ZL, Yue SJ, Zhang S, Tang YP. Why does COVID-19 continue to spread despite mass vaccination? *Front Public Health.* 2022;10:938108. doi:10.3389/fpubh.2022.938108
36. Willis GA, Bloomfield L, Berry M, et al. The impact of a vaccine mandate and the COVID-19 pandemic on influenza vaccination uptake in Western Australian health care students. *Vaccine.* Sep 16 2022;40(39):5651-5656. doi:10.1016/j.vaccine.2022.08.040
37. Adhikari B, Yeong Cheah P, von Seidlein L. Trust is the common denominator for COVID-19 vaccine acceptance: A literature review. *Vaccine X.* Dec 2022;12:100213. doi:10.1016/j.jvacx.2022.100213
38. Younus MM, Al-Jumaili AA. An overview of COVID-19 vaccine safety and post-marketing surveillance systems. *Innov Pharm.* 2021;12(4)doi:10.24926/iip.v12i4.4294
39. van der Pol KH, Aljofan M, Blin O, et al. Drug repurposing of generic drugs: Challenges and the potential role for government. *Applied health economics and health policy.* Nov 2023;21(6):831-840. doi:10.1007/s40258-023-00816-6
40. Kumari M, Lu RM, Li MC, et al. A critical overview of current progress for COVID-19: development of vaccines, antiviral drugs, and therapeutic antibodies. *J Biomed Sci.* Sep 12 2022;29(1):68. doi:10.1186/s12929-022-00852-9
41. Union E. the European Parliament and of the Council on the Community code relating to medicinal products for human use: Commission Directive 2003/63/EC. European Union. Accessed Nov 12, 2022. <https://eur-lex.europa.eu/legal-content/en/ALL/?uri=CELEX%3A32003L0063>
42. Pioli PD. Plasma Cells, the Next Generation: Beyond Antibody Secretion. *Front Immunol.* 2019;10:2768. doi:10.3389/fimmu.2019.02768
43. Duan Y, Luo J, Zhao L, et al. Reporting and data sharing level for COVID-19 vaccine trials: A cross-sectional study. *EBioMedicine.* Apr 2022;78:103962. doi:10.1016/j.ebiom.2022.103962
44. Puspitarani F, Sitaresmi MN, Ahmad RA. Adverse events following immunization of COVID-19 vaccine among children aged 6-11 years. *Front Public Health.* 2022;10:999354. doi:10.3389/fpubh.2022.999354
45. Chi WY, Li YD, Huang HC, et al. COVID-19 vaccine update: vaccine effectiveness, SARS-CoV-2 variants, boosters, adverse effects, and immune correlates of protection. *J Biomed Sci.* Oct 15 2022;29(1):82. doi:10.1186/s12929-022-00853-8

46. Stowe J, Miller E, Andrews N, Whitaker HJ. Risk of myocarditis and pericarditis after a COVID-19 mRNA vaccine booster and after COVID-19 in those with and without prior SARS-CoV-2 infection: A self-controlled case series analysis in England. *PLoS Med.* Jun 2023;20(6):e1004245. doi:10.1371/journal.pmed.1004245
47. Matsumoto N, Shimizu J, Yokoyama Y, Tsukahara H, Yorifuji T. Adverse reactions and attitudes toward the BNT162b2 COVID-19 vaccine in children 5 to 11 years of age in Japan. *J Epidemiol.* Feb 5 2023;33(2):110-111. doi:10.2188/jea.JE20220265
48. Chowdhury P, Bhattacharya S, Gogoi B, Veeranna RP, Kumar S. An update on complications associated with SARS-CoV-2 infection and COVID-19 vaccination. *Vaccines (Basel).* Sep 29 2022;10(10)doi:10.3390/vaccines10101639
49. Almadani OA, Alshammari TM. Vaccine adverse event reporting system (VAERS): Evaluation of 31 years of reports and pandemics' impact. *Saudi Pharm J.* Dec 2022;30(12):1725-1735. doi:10.1016/j.jsps.2022.10.001
50. Santi Laurini G, Montanaro N, Broccoli M, Bonaldo G, Motola D. Real-life safety profile of mRNA vaccines for COVID-19: An analysis of VAERS database. *Vaccine.* May 2 2023;41(18):2879-2886. doi:10.1016/j.vaccine.2023.03.054
51. Xu W, Ren W, Wu T, et al. Real-world safety of COVID-19 mRNA vaccines: A systematic review and meta-analysis. *Vaccines (Basel).* Jun 19 2023;11(6)doi:10.3390/vaccines11061118
52. Jiao T, Huang Y, Sun H, Yang L. Research progress of post-acute sequelae after SARS-CoV-2 infection. *Cell Death Dis.* Apr 11 2024;15(4):257. doi:10.1038/s41419-024-06642-5
53. Lam ICH, Zhang R, Man KKC, et al. Persistence in risk and effect of COVID-19 vaccination on long-term health consequences after SARS-CoV-2 infection. *Nat Commun.* Feb 26 2024;15(1):1716. doi:10.1038/s41467-024-45953-1
54. Ittiwut C, Mahasirimongkol S, Srisont S, et al. Genetic basis of sudden death after COVID-19 vaccination in Thailand. *Heart Rhythm.* Nov 2022;19(11):1874-1879. doi:10.1016/j.hrthm.2022.07.019
55. Salomao MLM, Queiroz F, Mendes LM, et al. COVID-19-related multisystem inflammatory syndrome in adult: the first death in Brazil. *Rev Inst Med Trop Sao Paulo.* 2023;65:e50. doi:10.1590/S1678-9946202365050
56. Patel T, Kelleman M, West Z, et al. Comparison of multisystem inflammatory syndrome in children-related myocarditis, classic viral myocarditis, and COVID-19 vaccine-related myocarditis in children. *J Am Heart Assoc.* May 3 2022;11(9):e024393. doi:10.1161/JAHA.121.024393
57. Komine-Aizawa S, Haruyama Y, Deguchi M, et al. The vaccination status and adverse effects of COVID-19 vaccine among pregnant women in Japan in 2021. *J Obstet Gynaecol Res.* Jul 2022;48(7):1561-1569. doi:10.1111/jog.15285
58. Yu CK, Tsao S, Ng CW, et al. Cardiovascular assessment up to one year after COVID-19 vaccine-associated myocarditis. *Circulation.* Aug 2023;148(5):436-439. doi:10.1161/CIRCULATIONAHA.123.064772
59. Boschi C, Scheim DE, Bancod A, et al. SARS-CoV-2 Spike protein induces hemagglutination: implications for COVID-19 morbidities and therapeutics and for vaccine adverse effects. *Int J Mol Sci.* Dec 7 2022;23(24)doi:10.3390/ijms232415480
60. Pulvirenti F, Mortari EP, Putotto C, et al. COVID-19 severity, cardiological outcome, and Immunogenicity of mRNA vaccine on adult patients with 22q11.2 DS. *J Allergy Clin Immunol Pract.* Jan 2023;11(1):292-305 e2. doi:10.1016/j.jaip.2022.10.010
61. Shin L, Shahsavari S, Lee C, Laborada J, Egeberg A, Wu JJ. Adverse Effects of the COVID-19 Vaccine in Patients With Psoriasis. *Cutis.* Feb 2023;111(2):80-81. doi:10.12788/cutis.0700
62. Pour Mohammad A, Mashayekhi F, Seirafianpour F, Gholizadeh Mesgarha M, Goodarzi A. COVID-19 and COVID-19 vaccine-related dermatological reactions: An interesting case series with a narrative review of the potential critical and non-critical mucocutaneous adverse effects related to virus, therapy, and the vaccination. *Clin Case Rep.* Apr 2022;10(4):e05775. doi:10.1002/ccr3.5775
63. Ichhpujani P, Parmar UPS, Duggal S, Kumar S. COVID-19 vaccine-associated ocular adverse effects: An overview. *Vaccines (Basel).* Nov 7 2022;10(11)doi:10.3390/vaccines10111879
64. Finsterer J. Vaccine adverse Event reporting system could miss or misinterpret neurological side effects of COVID-19 vaccinations. *Ann Neurol.* Jul 2022;92(1):157-158. doi:10.1002/ana.26369
65. Desai AD, Lavelle M, Boursiquot BC, Wan EY. Long-term complications of COVID-19. *Am J Physiol Cell Physiol.* Jan 1 2022;322(1):C1-C11. doi:10.1152/ajpcell.00375.2021
66. FDA. COVID-19 bivalent vaccine boosters. FDA. Accessed Dec 15, 2022. <https://www.fda.gov/emergency-preparedness-and-response/coronavirus-disease-2019-covid-19/covid-19-bivalent-vaccine-boosters>

67. Shrestha NK, Burke PC, Nowacki AS, Simon JF, Hagen A, Gordon SM. Effectiveness of the coronavirus disease 2019 (COVID-19) bivalent vaccine. *Preprint: MedRxiv*. 2023;doi:10.1101/2022.12.17.22283625
68. Nuwarda RF, Ramzan I, Weekes L, Kayser V. Vaccine hesitancy: Contemporary issues and historical background. *Vaccines (Basel)*. Sep 22 2022;10(10)doi:10.3390/vaccines10101595
69. Lau JJ, Cheng SMS, Leung K, et al. Author Correction: Real-world COVID-19 vaccine effectiveness against the Omicron BA.2 variant in a SARS-CoV-2 infection-naive population. *Nat Med*. Oct 24 2023;doi:10.1038/s41591-023-02648-2
70. Romer D, Jamieson KH. Conspiratorial thinking as a precursor to opposition to COVID-19 vaccination in the US: a multi-year study from 2018 to 2021. *Sci Rep*. Nov 3 2022;12(1):18632. doi:10.1038/s41598-022-22014-5
71. Glaser N, Diexer S, Klee B, et al. The contribution of SARS-CoV-2 to the burden of acute respiratory infections in winter season 2022/2023: Results from the DigiHero study. *Int J Infect Dis*. Apr 15 2024;107057. doi:10.1016/j.ijid.2024.107057
72. Maeda SS, Saraiva GL, Hayashi LF, et al. Seasonal variation in the serum 25-hydroxyvitamin D levels of young and elderly active and inactive adults in Sao Paulo, Brazil: The Sao Paulo Vitamin D Evaluation Study (SPADES). *Dermatoendocrinol*. Jan 1 2013;5(1):211-7. doi:10.4161/derm.24476
73. Wimalawansa SJ. Rapidly increasing serum 25(OH)D boosts the immune system, against infections-sepsis and COVID-19. *Nutrients*. Jul 21 2022;14(14)doi:10.3390/nu14142997
74. Wimalawansa SJ. Physiological basis for using vitamin D to improve health. *Biomedicines*. May 26 2023;11(6)doi:10.3390/biomedicines11061542
75. Chung KH, Cho SJ. General guideline in the endoscopy room to avoid air-borne infection during the COVID-19 pandemic. *Clin Endosc*. Sep 2022;55(5):688-690. doi:10.5946/ce.2022.146
76. Wimalawansa SJ. Modes of preventing secondary community spread and future epidemics of COVID-19. *J Immuno Biol*. 2020;5(1)(10.37421/JIB.2020.5.147):1-5. doi:10.37421/JIB.2020.5.147
77. Wimalawansa S. Overcoming infections including COVID-19, by maintaining circulating 25(OH)D concentrations above 50 ng/mL. *Pathology & Lab Medicine Int*. 2022;14:37–60.
78. Mosca C, Colucci A, Savoia F, et al. Vitamin D levels in the pre- and post-COVID-19 pandemic periods and related confinement at pediatric age. *Nutrients*. Apr 26 2023;15(9)doi:10.3390/nu15092089
79. Polonowita A, Wimalawansa SJ. Molecular quantum and logic process of consciousness—Vitamin D big-data in COVID-19—A case for incorporating machine learning in medicine. *Euro J Biomedical Pharma Sci*. 2023;10(12):1-17.
80. Acharya CB, Schrom J, Mitchell AM, et al. Viral Load Among Vaccinated and Unvaccinated, Asymptomatic and Symptomatic Persons Infected With the SARS-CoV-2 Delta Variant. *Open Forum Infect Dis*. May 2022;9(5):ofac135. doi:10.1093/ofid/ofac135
81. Hagan L, McCormick, DW, Lee, C, et al. Outbreak of SARS-CoV-2 B.1.617.2 (delta) variant infections among incarcerated persons in a federal prison—Texas, July–August 2021. *The Lancet-Infectious diseases*. 2021;70:1349-1354.
82. Wang Q, Iketani S, Li Z, et al. Alarming antibody evasion properties of rising SARS-CoV-2 BQ and XBB subvariants. *Cell*. Jan 19 2023;186(2):279-286 e8. doi:10.1016/j.cell.2022.12.018
83. Focosi D, Maggi F. Neutralising antibody escape of SARS-CoV-2 spike protein: Risk assessment for antibody-based Covid-19 therapeutics and vaccines. *Rev Med Virol*. Nov 2021;31(6):e2231. doi:10.1002/rmv.2231
84. Liu Y, Rocklov J. The reproductive number of the Delta variant of SARS-CoV-2 is far higher compared to the ancestral SARS-CoV-2 virus. *J Travel Med*. Oct 11 2021;28(7)doi:10.1093/jtm/taab124
85. Konishi T. Mutations in SARS-CoV-2 are on the increase against the acquired immunity. *PLoS One*. 2022;17(7):e0271305. doi:10.1371/journal.pone.0271305
86. Wu Y, Shi J, He X, et al. Protection of the receptor binding domain (RBD) dimer against SARS-CoV-2 and its variants. *J Virol*. Oct 16 2023:e0127923. doi:10.1128/jvi.01279-23
87. Takashita E, Yamayoshi S, Simon V, et al. Efficacy of Antibodies and Antiviral Drugs against Omicron BA.2.12.1, BA.4, and BA.5 Subvariants. *N Engl J Med*. Aug 4 2022;387(5):468-470. doi:10.1056/NEJMc2207519
88. Shayak B, Sharma MM, Gaur M, Mishra AK. Impact of reproduction number on the multiwave spreading dynamics of COVID-19 with temporary immunity: A mathematical model. *Int J Infect Dis*. Mar 2021;104:649-654. doi:10.1016/j.ijid.2021.01.018
89. Puhach O, Adea K, Hulo N, et al. Infectious viral load in unvaccinated and vaccinated individuals infected with ancestral, Delta or Omicron SARS-CoV-2. *Nat Med*. Jul 2022;28(7):1491-1500. doi:10.1038/s41591-022-01816-0

90. Canning AG, Watson KE, McCreedy KE, Olawepo JO. Ethics and effectiveness of US COVID-19 vaccine mandates and vaccination passports: A review. *J Res Health Sci*. May 11 2022;22(2):e00546. doi:10.34172/jrhs.2022.81
91. Y T. Stop calling it a pandemic of the unvaccinated. <https://www.theatlantic.com/ideas/archive/2021/09/persuade-unvaccinated-protect-unvaccinated/620091/>. Accessed Feb 4th 2023, 2023. <https://www.theatlantic.com/ideas/archive/2021/09/persuade-unvaccinated-protect-unvaccinated/620091/>
92. Bergwerk M, Gonen T, Lustig Y, et al. Covid-19 Breakthrough Infections in Vaccinated Health Care Workers. *N Engl J Med*. Oct 14 2021;385(16):1474-1484. doi:10.1056/NEJMoa2109072
93. Duroseau B, Kipshidze N, Limaye RJ. The impact of delayed access to COVID-19 vaccines in low- and lower-middle-income countries. *Front Public Health*. 2022;10:1087138. doi:10.3389/fpubh.2022.1087138
94. Daems R, Maes E. The race for COVID-19 vaccines: accelerating innovation, fair allocation and distribution. *Vaccines (Basel)*. Sep 2 2022;10(9)doi:10.3390/vaccines10091450
95. Forman R, Shah S, Jeurissen P, Jit M, Mossialos E. COVID-19 vaccine challenges: What have we learned so far and what remains to be done? *Health Policy*. May 2021;125(5):553-567. doi:10.1016/j.healthpol.2021.03.013
96. Yarlagaadda H, Patel MA, Gupta V, et al. COVID-19 vaccine challenges in developing and developed countries. *Cureus*. Apr 2022;14(4):e23951. doi:10.7759/cureus.23951
97. Group-C19.com. Vitamin D for COVID-19: real-time analysis of all 300 studies: <https://c19early.org/d>. Accessed January 25, 2023, <https://c19vitamind.com/>
98. Anonymos. Vaccine monopolies make cost of vaccinating the world against COVID at least 5 times more expensive than it could be. Oxfam. 2023 March 10. <https://www.oxfam.org/en/press-releases/vaccine-monopolies-make-cost-vaccinating-world-against-covid-least-5-times-more>
99. E E-P. It will be hard to find a farmer left': Sri Lanka reels from rash fertiliser ban. *Guardian*. 2023, May 22. <https://www.theguardian.com/world/2022/apr/20/sri-lanka-fertiliser-ban-president-rajapaksa-farmers-harvests-collapse>
100. Athas I LC, Mogul R, Gonzalez-Roman D. Sri Lanka is 'bankrupt,' Prime Minister says. *CNN*, Sri Lanka. 2023, May 23. <https://www.cnn.com/2022/07/05/asia/sri-lanka-bankrupt-fuel-crisis-intl-hnk/index.html>
101. Shrestha LB, Foster C, Rawlinson W, Tedla N, Bull RA. Evolution of the SARS-CoV-2 omicron variants BA.1 to BA.5: Implications for immune escape and transmission. *Rev Med Virol*. Sep 2022;32(5):e2381. doi:10.1002/rmv.2381
102. Reed G, Hendlin Y, Desikan A, MacKinney T, Berman E, Goldman GT. The disinformation playbook: how industry manipulates the science-policy process-and how to restore scientific integrity. *J Public Health Policy*. Dec 2021;42(4):622-634. doi:10.1057/s41271-021-00318-6
103. Grant WB. Vitamin D acceptance delayed by big pharma following the Disinformation Playbook. Accessed March 22nd, 2022, <http://orthomolecular.org/resources/omns/v14n22.shtml>
104. Hess S, Lancsar E, Mariel P, et al. The path towards herd immunity: Predicting COVID-19 vaccination uptake through results from a stated choice study across six continents. *Soc Sci Med*. Apr 2022;298:114800. doi:10.1016/j.socscimed.2022.114800
105. Spencer S. Widespread claims misrepresent effectiveness of COVID-19 vaccines. *FactCheck.org*. Accessed Nov 10, 2022.
106. FDA. Coronavirus (COVID-19) update: FDA authorizes Moderna, Pfizer-BioNTech bivalent COVID-19 vaccines for use as a booster dose. FDA. January 5th, 2023. <https://www.fda.gov/news-events/press-announcements/coronavirus-covid-19-update-fda-authorizes-moderna-pfizer-biontech-bivalent-covid-19-vaccines-use>
107. Sharma K, Li-Kim-Moy J. COVID-19 vaccines in 2023. *Aust Prescr*. Oct 2023;46(3):60-63. doi:10.18773/austprescr.2023.020
108. Chalkias S, Harper C, Vrbecky K, et al. A bivalent Omicron-containing booster vaccine against Covid-19. *N Engl J Med*. Oct 6 2022;387(14):1279-1291. doi:10.1056/NEJMoa2208343
109. Chatzilena A, Hyams C, Challen R, et al. Relative vaccine effectiveness of mRNA COVID-19 boosters in people aged at least 75 years during the spring-summer (monovalent vaccine) and autumn-winter (bivalent vaccine) booster campaigns: a prospective test negative case-control study, United Kingdom, 2022. *Euro surveillance : bulletin Europeen sur les maladies transmissibles = European communicable disease bulletin*. Nov 2023;28(48)doi:10.2807/1560-7917.ES.2023.28.48.2300173
110. Chenchula S, Karunakaran P, Sharma S, Chavan M. Current evidence on efficacy of COVID-19 booster dose vaccination against the Omicron variant: A systematic review. *J Med Virol*. Jul 2022;94(7):2969-2976. doi:10.1002/jmv.27697

111. Lee J, Park S, Kim JY, et al. No correlation of neutralizing antibody titers against the Omicron variant after a booster dose of COVID-19 vaccines with subsequent breakthrough Omicron infections among healthcare workers. *J Infect.* Dec 2022;85(6):e177-e180. doi:10.1016/j.jinf.2022.10.007
112. Thimbleby H. Technology and the future of healthcare. *J Public Health Res.* Dec 1 2013;2(3):e28. doi:10.4081/jphr.2013.e28
113. Polonowita A, Wimalawansa, SJ. Impact of Withholding Cost-Effective Early Treatments, Such as Vitamin D, on COVID-19: An Analysis Using an Innovative Logical Paradigm. *Wold J Adanced Pharma Life Sci.* 2023;(in press)
114. Basu K, Sinha R, Ong A, Basu T. Artificial intelligence: How is it changing medical sciences and its future? *Indian J Dermatol.* Sep-Oct 2020;65(5):365-370. doi:10.4103/ijd.IJD_421_20
115. Junaid SB, Imam AA, Balogun AO, et al. Recent advancements in emerging technologies for healthcare management systems: A survey. *Healthcare (Basel).* Oct 3 2022;10(10)doi:10.3390/healthcare10101940
116. Kaufman HW, Niles JK, Kroll MH, Bi C, Holick MF. SARS-CoV-2 positivity rates associated with circulating 25-hydroxyvitamin D levels. *PLoS One.* 2020;15(9):e0239252. doi:10.1371/journal.pone.0239252
117. Raisi-Estabragh Z, McCracken C, Bethell MS, et al. Greater risk of severe COVID-19 in Black, Asian and Minority Ethnic populations is not explained by cardiometabolic, socioeconomic or behavioural factors, or by 25(OH)-vitamin D status: study of 1326 cases from the UK Biobank. *Journal of public health.* Aug 18 2020;42(3):451-460. doi:10.1093/pubmed/fdaa095
118. Hastie CE, Pell JP, Sattar N. Vitamin D and COVID-19 infection and mortality in UK Biobank. *Eur J Nutr.* Feb 2021;60(1):545-548. doi:10.1007/s00394-020-02372-4
119. Kumar R, Rathi H, Haq A, Wimalawansa SJ, Sharma A. Putative roles of vitamin D in modulating immune response and immunopathology associated with COVID-19. *Virus Res.* Jan 15 2021;292:198235. doi:10.1016/j.virusres.2020.198235
120. Wimalawansa SJ, Polonowita, A. Boosting immunity with vitamin D for preventing complications and deaths from COVID-19. National Science Foundation, Sri Lanka.; 2021:171-198.
121. Villasis-Keever MA, Lopez-Alarcon MG, Miranda-Novales G, et al. Efficacy and Safety of Vitamin D Supplementation to Prevent COVID-19 in Frontline Healthcare Workers. A Randomized Clinical Trial. *Archives of medical research.* Jun 2022;53(4):423-430. doi:10.1016/j.arcmed.2022.04.003
122. Entrenas Castillo M, Entrenas Costa LM, Vaquero Barrios JM, et al. "Effect of calcifediol treatment and best available therapy versus best available therapy on intensive care unit admission and mortality among patients hospitalized for COVID-19: A pilot randomized clinical study". *J Steroid Biochem Mol Biol.* Oct 2020;203:105751. doi:10.1016/j.jsbmb.2020.105751
123. Rajter JC, Sherman MS, Fatteh N, Vogel F, Sacks J, Rajter JJ. Use of Ivermectin Is Associated With Lower Mortality in Hospitalized Patients With Coronavirus Disease 2019: The Ivermectin in COVID Nineteen Study. *Chest.* Jan 2021;159(1):85-92. doi:10.1016/j.chest.2020.10.009
124. Padhy BM, Mohanty RR, Das S, Meher BR. Therapeutic potential of ivermectin as add on treatment in COVID 19: A systematic review and meta-analysis. *J Pharm Pharm Sci.* 2020;23:462-469. doi:10.18433/jpps31457
125. Grant WB. Vitamin D acceptance delayed by big pharma following the Disinformation Playbook. Accessed March 22nd, 2022, <http://orthomolecular.org/resources/omns/v14n22.shtml>
126. Wimalawansa SJ. Infections and autoimmunity-The immune system and vitamin D: A Systematic Review. *Nutrients.* Sep 2 2023;15(17)doi:10.3390/nu15173842
127. Wimalawansa SJ. Prophylactic use of vitamin D to maintain a robust immune system against infections like SARS-CoV-2. *Glob J Endocrinol Metab GJEM 000571.* 2023;3(5):1-9. doi:10.31031/GJEM.2023.03.000571
128. Cannata-Andia JB, Diaz-Sottolano A, Fernandez P, et al. A single-oral bolus of 100,000 IU of cholecalciferol at hospital admission did not improve outcomes in the COVID-19 disease: the COVID-VIT-D-a randomised multicentre international clinical trial. *BMC Med.* Feb 18 2022;20(1):83. doi:10.1186/s12916-022-02290-8
129. Murai IH, Fernandes AL, Sales LP, et al. Effect of a single high dose of vitamin D3 on hospital length of stay in patients Wpwith moderate to severe COVID-19: A randomized clinical trial. *JAMA.* Mar 16 2021;325(11):1053-1060. doi:10.1001/jama.2020.26848
130. Al-Jarallah M, Rajan R, Dashti R, et al. In-hospital mortality in SARS-CoV-2 stratified by serum 25-hydroxy-vitamin D levels: A retrospective study. *J Med Virol.* Oct 2021;93(10):5880-5885. doi:10.1002/jmv.27133

131. Pecina JL, Merry SP, Park JG, Thacher TD. Vitamin D Status and Severe COVID-19 Disease Outcomes in Hospitalized Patients. *J Prim Care Community Health*. Jan-Dec 2021;12:21501327211041206. doi:10.1177/21501327211041206
132. Cereda E, Bogliolo L, Lobascio F, et al. Vitamin D supplementation and outcomes in coronavirus disease 2019 (COVID-19) patients from the outbreak area of Lombardy, Italy. *Nutrition*. Feb 2021;82:111055. doi:10.1016/j.nut.2020.111055
133. Amin HA, Drenos F. No evidence that vitamin D is able to prevent or affect the severity of COVID-19 in individuals with European ancestry: a Mendelian randomisation study of open data. *BMJ Nutr Prev Health*. 2021;4(1):42-48. doi:10.1136/bmjnp-2020-000151
134. Ben-Zuk N, Dechtman ID, Henn I, et al. Potential prophylactic treatments for COVID-19. *Viruses*. Jul 2 2021;13(7)doi:10.3390/v13071292
135. Sharun K, Dhama K, Patel SK, et al. Ivermectin, a new candidate therapeutic against SARS-CoV-2/COVID-19. *Ann Clin Microbiol Antimicrob*. May 30 2020;19(1):23. doi:10.1186/s12941-020-00368-w
136. Wimalawansa SJ. Fighting against COVID-19: Boosting the immunity with micronutrients, stress reduction, physical activity, and vitamin D. *Nutrition and Food Science Journal (Sci Literature)*. 2020c;3(1)(126):1-4.
137. Pilz S, Zittermann A, Trummer C, et al. Vitamin D testing and treatment: a narrative review of current evidence. *Endocr Connect*. Feb 1 2019;8(2):R27-R43. doi:10.1530/EC-18-0432
138. Martinez MA. Lack of effectiveness of repurposed drugs for COVID-19 treatment. *Front Immunol*. 2021;12:635371. doi:10.3389/fimmu.2021.635371
139. MacFarlane D, Tay LQ, Hurlstone MJ, Ecker UKH. Refuting spurious COVID-19 treatment claims reduces demand and misinformation sharing. *J Appl Res Mem Cogn*. Jun 2021;10(2):248-258. doi:10.1016/j.jarmac.2020.12.005
140. Yang YT, Schaffer DeRoo S. Disciplining physicians who spread medical misinformation. *J Public Health Manag Pract*. Nov-Dec 01 2022;28(6):595-598. doi:10.1097/PHH.0000000000001616
141. Schellack N, Strydom M, Pepper MS, et al. Social media and COVID-19-perceptions and public deceptions of ivermectin, colchicine and hydroxychloroquine: Lessons for future pandemics. *Antibiotics (Basel)*. Mar 25 2022;11(4)doi:10.3390/antibiotics11040445
142. FDA. Why you should not use ivermectin to treat or prevent COVID-19. FDA. 2023 March 12. <https://www.fda.gov/consumers/consumer-updates/why-you-should-not-use-ivermectin-treat-or-prevent-covid-19>
143. Sun LH WL, GodfreyH. Doctors who put lives at risk with COVID misinformation rarely punished. *Washington Post*. 2023, Aug 10. <https://www.washingtonpost.com/health/2023/07/26/covid-misinformation-doctor-discipline/>
144. Maziarz M, Stencil A. The failure of drug repurposing for COVID-19 as an effect of excessive hypothesis testing and weak mechanistic evidence. *Hist Philos Life Sci*. Oct 18 2022;44(4):47. doi:10.1007/s40656-022-00532-9
145. Anderson AA SG. Risks elaborated vs. risks downplayed: The effect of risk comparisons in mainstream media during COVID-19 on risk perceptions and anxiety levels. *Frontiers: Sci Environ Commun*. 2021;6doi:10.3389/fcomm.2021.646001
146. Colomin C SMH, Youngs, R. The impact of disinformation on democratic processes and human rights in the world. *European Parliament*. 2023, May 7. [https://www.europarl.europa.eu/RegData/etudes/STUD/2021/653635/EXPO_STU\(2021\)653635_EN.pdf](https://www.europarl.europa.eu/RegData/etudes/STUD/2021/653635/EXPO_STU(2021)653635_EN.pdf)
147. Chavda VP, Sonak SS, Munshi NK, Dhamade PN. Pseudoscience and fraudulent products for COVID-19 management. *Environmental science and pollution research international*. Sep 2022;29(42):62887-62912. doi:10.1007/s11356-022-21967-4
148. anonymous. FLCCC treatment protocols. FLCCC Alliance. 2023, Nov. 22. <https://covid19criticalcare.com/treatment-protocols/>
149. Anonymus. At home treatment guide. World Council of Health. 2022, August 7. https://worldcouncilforhealth.org/wp-content/uploads/2021/09/WCH-At-Home-Treatment-Guide_30-Sept-2021.pdf
150. Kory P MJ. *The war on ivermectin: The medicine that saved millions and could have ended the pandemic* [<https://rumble.com/v41agxg-2023-12-14-mikki-willis-pierre-kory-the-war-on-ivermectin.html>]. vol 1. ICAN Press; 2023.
151. A. A. Why cheap, older drugs that might treat COVID never get out of the lab. <https://kffhealthnews.org/news/article/repurposing-cheap-older-drugs-as-covid-treatment-camostat/>
152. Henrina J, Lim MA, Pranata R. COVID-19 and misinformation: how an infodemic fuelled the prominence of vitamin D. *Br J Nutr*. Feb 14 2021;125(3):359-360. doi:10.1017/S0007114520002950

153. Chanda D, Hines JZ, Itoh M, et al. COVID-19 Vaccine Effectiveness Against Progression to In-Hospital Mortality in Zambia, 2021-2022. *Open Forum Infect Dis*. Sep 2022;9(9):ofac469. doi:10.1093/ofid/ofac469
154. Samanovic MI, Oom AL, Cornelius AR, et al. Vaccine-Acquired SARS-CoV-2 Immunity versus Infection-Acquired Immunity: A Comparison of Three COVID-19 Vaccines. *Vaccines (Basel)*. Dec 15 2022;10(12)doi:10.3390/vaccines10122152
155. Israel A, Shenhar Y, Green I, et al. Large-Scale Study of Antibody Titer Decay following BNT162b2 mRNA Vaccine or SARS-CoV-2 Infection. *Vaccines (Basel)*. Dec 31 2021;10(1)doi:10.3390/vaccines10010064
156. Guerin P, Yahi N, Azzaz F, Chahinian H, Sabatier JM, Fantini J. Structural Dynamics of the SARS-CoV-2 Spike Protein: A 2-Year Retrospective Analysis of SARS-CoV-2 Variants (from Alpha to Omicron) Reveals an Early Divergence between Conserved and Variable Epitopes. *Molecules*. Jun 15 2022;27(12)doi:10.3390/molecules27123851
157. Kontopoulou K, Nakas CT, Belai C, Papazisis G. Antibody titers after a third dose of the SARS-CoV-2 BNT162b2 vaccine in immunocompromised adults in Greece: Is a fourth dose necessary? *J Med Virol*. Oct 2022;94(10):5056-5060. doi:10.1002/jmv.27954
158. van Ewijk CE, Hazelhorst EI, Hahne SJM, Knol MJ. COVID-19 outbreak in an elderly care home: Very low vaccine effectiveness and late impact of booster vaccination campaign. *Vaccine*. Nov 2 2022;40(46):6664-6669. doi:10.1016/j.vaccine.2022.09.080
159. Suryawanshi YN, Biswas DA. Herd immunity to fight against COVID-19: A narrative review. *Cureus*. Jan 2023;15(1):e33575. doi:10.7759/cureus.33575
160. Biswas M, Rahaman S, Biswas TK, Haque Z, Ibrahim B. Association of sex, sge, and comorbidities with mortality in COVID-19 Patients: A systematic review and meta-analysis. *Intervirology*. Dec 9 2020;1-12. doi:10.1159/000512592
161. Wimalawansa SJ. Maintaining optimum health requires longer-term stable vitamin D concentrations. *Int J Regen Med*. 2020;3(3):1-5. doi:10.31487/j.RGM.2020.03.03
162. Wimalawansa SJ. Effective and practical ways to overcome vitamin D deficiency. *J Family Med Community Health*. 2021;8(1):1-8. doi:https://www.jscimedcentral.com/FamilyMedicine/familymedicine-7-1170.pdf
163. Lind ML, Copin R, McCarthy S, et al. Use of Whole-Genome Sequencing to Estimate the Contribution of Immune Evasion and Waning Immunity on Decreasing COVID-19 Vaccine Effectiveness. *J Infect Dis*. Mar 1 2023;227(5):663-674. doi:10.1093/infdis/jiac453
164. Zhang X, Chen LL, Ip JD, et al. Omicron sublineage recombinant XBB evades neutralising antibodies in recipients of BNT162b2 or CoronaVac vaccines. *Lancet Microbe*. Mar 2023;4(3):e131. doi:10.1016/S2666-5247(22)00335-4
165. Aslam S, Danziger-Isakov L, Mehra MR. COVID-19 vaccination immune paresis in heart and lung transplantation. *J Heart Lung Transplant*. Aug 2021;40(8):763-766. doi:10.1016/j.healun.2021.04.018
166. Shrestha NK, Burke PC, Nowacki AS, Simon JF, Hagen A, Gordon SM. Effectiveness of the coronavirus disease 2019 bivalent vaccine. *Open Forum Infect Dis*. Jun 2023;10(6):ofad209. doi:10.1093/ofid/ofad209
167. Shrestha NK, Shrestha P, Burke PC, Nowacki AS, Terpeluk P, Gordon SM. Coronavirus Disease 2019 Vaccine Boosting in Previously Infected or Vaccinated Individuals. *Clin Infect Dis*. Dec 19 2022;75(12):2169-2177. doi:10.1093/cid/ciac327
168. Cegolon L, Negro C, Mastrangelo G, Filon FL, group Ow. Primary SARS-CoV-2 Infections, Re-infections and Vaccine Effectiveness during the Omicron Transmission Period in Healthcare Workers of Trieste and Gorizia (Northeast Italy), 1 December 2021-31 May 2022. *Viruses*. Nov 30 2022;14(12)doi:10.3390/v14122688
169. Choueiri TK, Labaki C, Bakouny Z, et al. Breakthrough SARS-CoV-2 infections among patients with cancer following two and three doses of COVID-19 mRNA vaccines: a retrospective observational study from the COVID-19 and Cancer Consortium. *Lancet Reg Health Am*. Mar 2023;19:100445. doi:10.1016/j.lana.2023.100445
170. Wang Q, Guo Y, Iketani S, et al. Antibody evasion by SARS-CoV-2 Omicron subvariants BA.2.12.1, BA.4 and BA.5. *Nature*. Aug 2022;608(7923):603-608. doi:10.1038/s41586-022-05053-w
171. Antonelli M, Pujol JC, Spector TD, Ourselin S, Steves CJ. Risk of long COVID associated with delta versus omicron variants of SARS-CoV-2. *Lancet*. Jun 18 2022;399(10343):2263-2264. doi:10.1016/S0140-6736(22)00941-2
172. Hwang MJ, Hwang I, Park C, Park H, Son T, Kim JH. Clinical severity according to the primary infection variant in patients with suspected SARS-CoV-2 reinfection in Korea. *Epidemiol Health*. Dec 21 2023;45:e2023007. doi:10.4178/epih.e2023007

173. Riemersma KK, Haddock LA, 3rd, Wilson NA, et al. Shedding of infectious SARS-CoV-2 despite vaccination. *PLoS Pathog.* Sep 2022;18(9):e1010876. doi:10.1371/journal.ppat.1010876
174. MacIntyre CR, Costantino V, Heslop DJ. The potential impact of a recent measles epidemic on COVID-19 in Samoa. *BMC Infect Dis.* Oct 7 2020;20(1):735. doi:10.1186/s12879-020-05469-7
175. Wang Q, Bowen A, Valdez R, et al. Antibody Response to Omicron BA.4-BA.5 Bivalent Booster. *N Engl J Med.* Feb 9 2023;388(6):567-569. doi:10.1056/NEJMc2213907
176. Lewis NM, Murray N, Adams K, et al. Absolute and Relative Vaccine Effectiveness of Primary and Booster Series of COVID-19 Vaccines (mRNA and Adenovirus Vector) Against COVID-19 Hospitalizations in the United States, December 2021-April 2022. *Open Forum Infect Dis.* Jan 2023;10(1):ofac698. doi:10.1093/ofid/ofac698
177. Ma Y, Zhu P, Zhong G, et al. Serial negative response after standard and third (Booster) dose of COVID-19 inactivated vaccine is associated with low vitamin D levels in patients with solid cancers. *Front Med (Lausanne).* 2022;9:898606. doi:10.3389/fmed.2022.898606
178. Salter SM, Li D, Trentino K, et al. Safety of Four COVID-19 Vaccines across Primary Doses 1, 2, 3 and Booster: A Prospective Cohort Study of Australian Community Pharmacy Vaccinations. *Vaccines (Basel).* Nov 25 2022;10(12)doi:10.3390/vaccines10122017
179. CDC U. Effectiveness of bivalent mRNA vaccines in preventing symptomatic SARS-CoV-2 infection—increasing community access to testing program, United States: CDC-MMWR Report, Sept.–Nov. 2022. CDC. Accessed Jan 6, 2023. https://www.cdc.gov/mmwr/volumes/71/wr/mm7148e1.htm?s_cid=mm7148e1_w#T2_down
180. Tamanaha E, Zhang Y, Tanner NA. Profiling RT-LAMP tolerance of sequence variation for SARS-CoV-2 RNA detection. *PLoS One.* 2022;17(3):e0259610. doi:10.1371/journal.pone.0259610
181. Dingemans J, van der Veer B, Gorgels KMF, et al. Investigating SARS-CoV-2 breakthrough infections per variant and vaccine type. *Front Microbiol.* 2022;13:1027271. doi:10.3389/fmicb.2022.1027271
182. Uraki R, Ito M, Furusawa Y, et al. Humoral immune evasion of the omicron subvariants BQ.1.1 and XBB. *Lancet Infect Dis.* Jan 2023;23(1):30-32. doi:10.1016/S1473-3099(22)00816-7
183. Shrestha NK, Burke PC, Nowacki AS, Terpeluk P, Gordon SM. Necessity of coronavirus disease 2019 (COVID-19) vaccination in persons who have already had COVID-19. *Clin Infect Dis.* Aug 24 2022;75(1):e662-e671. doi:10.1093/cid/ciac022
184. Peng D, Gao Y, Liu Z, Zheng J, Liu Y. Interactive effect of booster vaccination and vitamin D status on antibody production of Omicron variant-infected adults: A real-world cohort study. *Clin Respir J.* Oct 2023;17(10):1067-1076. doi:10.1111/crj.13694
185. Das R, Jobayer Chisti M, Ahshanul Haque M, et al. Evaluating association of vaccine response to low serum zinc and vitamin D levels in children of a birth cohort study in Dhaka. *Vaccine.* Jan 3 2021;39(1):59-67. doi:10.1016/j.vaccine.2020.10.048
186. Connolly CM, Paik JJ. SARS-CoV-2 vaccination in the immunocompromised host. *J Allergy Clin Immunol.* Jul 2022;150(1):56-58. doi:10.1016/j.jaci.2022.05.001
187. Wimalawansa SJ. Global epidemic of coronavirus—COVID-19: What can we do to minimize risks? *European J Biomed & Pharma Sci.* 2020;7(3):432-438. doi:https://storage.googleapis.com/journal-uploads/ejbps/article_issue/volume_7_march_issue_3/1584436192.pdf
188. Zhou X, Wang H, Luo Y, Cui L, Guan Y, Zhang Y. Single-injection COVID-19 subunit vaccine elicits potent immune responses. *Acta Biomater.* Oct 1 2022;151:491-500. doi:10.1016/j.actbio.2022.08.006
189. Zhang Q, Wang M, Han C, et al. Intraduodenal Delivery of Exosome-Loaded SARS-CoV-2 RBD mRNA Induces a Neutralizing Antibody Response in Mice. *Vaccines (Basel).* Mar 16 2023;11(3)doi:10.3390/vaccines11030673
190. Polack FP, Thomas SJ, Kitchin N, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med.* Dec 31 2020;383(27):2603-2615. doi:10.1056/NEJMoa2034577
191. Baden LR, El Sahly HM, Essink B, et al. Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N Engl J Med.* Feb 4 2021;384(5):403-416. doi:10.1056/NEJMoa2035389
192. Thompson MG, Burgess JL, Naleway AL, et al. Interim Estimates of Vaccine Effectiveness of BNT162b2 and mRNA-1273 COVID-19 Vaccines in Preventing SARS-CoV-2 Infection Among Health Care Personnel, First Responders, and Other Essential and Frontline Workers - Eight U.S. Locations, December 2020-March 2021. *MMWR Morbidity and mortality weekly report.* Apr 2 2021;70(13):495-500. doi:10.15585/mmwr.mm7013e3
193. Anonymus. Coronavirus disease (COVID-19): Herd immunity, lockdowns and COVID-19. WHO-Questins and answers. Accessed 20 february 2024, <https://www.who.int/news-room/questions-and-answers/item/herd-immunity-lockdowns-and-covid-19>
194. Singh V. Can Vitamins, as Epigenetic Modifiers, Enhance Immunity in COVID-19 Patients with Non-communicable Disease? *Curr Nutr Rep.* Sep 2020;9(3):202-209. doi:10.1007/s13668-020-00330-4

195. Weixel N. CDC: More vaccinated adults hospitalized with omicron, but fewer admitted to ICU. <https://thehill.com/policy/healthcare/592847-cdc-more-vaccinated-adults-hospitalized-with-omicron-but-fewer-admitted-to/>
196. Uversky VN, Redwan EM, Makis W, Rubio-Casillas A. IgG4 antibodies induced by repeated vaccination may generate immune tolerance to the SARS-CoV-2 Spike protein. *Vaccines (Basel)*. May 17 2023;11(5)doi:10.3390/vaccines11050991
197. Chamie JJ, Hibberd JA, Scheim DE. COVID-19 excess deaths in Peru's 25 States in 2020: Nationwide trends, confounding factors, and correlations with the extent of ivermectin treatment by State. *Cureus*. Aug 2023;15(8):e43168. doi:10.7759/cureus.43168
198. Mustafa A, Shekhar C. Concentration levels of serum 25-Hydroxyvitamin-D and vitamin D deficiency among children and adolescents of India: a descriptive cross-sectional study. *BMC Pediatr*. Aug 6 2021;21(1):334. doi:10.1186/s12887-021-02803-z
199. Yin Z, Fang Q, Wen T, et al. Effectiveness of COVID-19 vaccines against SARS-CoV-2 Omicron variants during two outbreaks from March to May 2022 in Quzhou, China. *Hum Vaccin Immunother*. Dec 31 2023;19(1):2163813. doi:10.1080/21645515.2022.2163813
200. Tsiakalos A, Ziakas PD, Polyzou E, Schinas G, Akinosoglou K. Early fluvoxaminereduces the risk for clinical deterioration in symptomatic outpatients with COVID-19: A real-world, retrospective, before-after analysis. *Microorganisms*. Aug 12 2023;11(8)doi:10.3390/microorganisms11082073
201. Lenze EJ, Mattar C, Zorumski CF, et al. Fluvoxamine vs placebo and clinical deterioration in outpatients with symptomatic COVID-19: A randomized clinical trial. *JAMA*. Dec 8 2020;324(22):2292-2300. doi:10.1001/jama.2020.22760
202. Mocanu ML, Amariei S. Elderberries-A source of bioactive compounds with antiviral action. *Plants (Basel)*. Mar 10 2022;11(6)doi:10.3390/plants11060740
203. Saber-Moghaddam N, Salari S, Hejazi S, et al. Oral nano-curcumin formulation efficacy in management of mild to moderate hospitalized coronavirus disease-19 patients: An open label nonrandomized clinical trial. *Phytother Res*. May 2021;35(5):2616-2623. doi:10.1002/ptr.7004
204. Koshak AE, Koshak EA, Mobeireek AF, et al. Nigella sativa for the treatment of COVID-19: An open-label randomized controlled clinical trial. *Complement Ther Med*. Sep 2021;61:102769. doi:10.1016/j.ctim.2021.102769
205. Lan SH, Lee HZ, Chao CM, Chang SP, Lu LC, Lai CC. Efficacy of melatonin in the treatment of patients with COVID-19: A systematic review and meta-analysis of randomized controlled trials. *J Med Virol*. May 2022;94(5):2102-2107. doi:10.1002/jmv.27595
206. Cosentino M, Vernocchi V, Martini S, et al. Early outpatient treatment of COVID-19: A retrospective analysis of 392 cases in Italy. *J Clin Med*. Oct 18 2022;11(20)doi:10.3390/jcm11206138
207. Wan EYF, Wang B, Mathur S, et al. Molnupiravir and nirmatrelvir-ritonavir reduce mortality risk during post-acute COVID-19 phase. *J Infect*. Jun 2023;86(6):622-625. doi:10.1016/j.jinf.2023.02.029
208. Fung KW, Baye F, Baik SH, McDonald CJ. Nirmatrelvir and Molnupiravir and post-COVID-19 condition in older patients. *JAMA Intern Med*. Dec 1 2023;183(12):1404-1406. doi:10.1001/jamainternmed.2023.5099
209. Najjar-Debbiny R, Gronich N, Weber G, et al. Effectiveness of Paxlovid in reducing severe coronavirus disease 2019 and mortality in high-risk patients. *Clin Infect Dis*. Feb 8 2023;76(3):e342-e349. doi:10.1093/cid/ciac443
210. Ioannou GN, Berry K, Rajeevan N, et al. Effectiveness of Nirmatrelvir-Ritonavir against the development of post-COVID-19 conditions among U.S. Veterans : A target trial emulation. *Ann Intern Med*. Nov 2023;176(11):1486-1497. doi:10.7326/M23-1394
211. Ader F, Bouscambert-Duchamp M, Hites M, et al. Remdesivir plus standard of care versus standard of care alone for the treatment of patients admitted to hospital with COVID-19 (DisCoVeRy): a phase 3, randomised, controlled, open-label trial. *Lancet Infect Dis*. Feb 2022;22(2):209-221. doi:10.1016/S1473-3099(21)00485-0
212. Burhan E, Liu K, Marwali EM, et al. Characteristics and outcomes of patients with severe COVID-19 in Indonesia: Lessons from the first wave. *PLoS One*. 2023;18(9):e0290964. doi:10.1371/journal.pone.0290964
213. Yang S, Shen S, Hou N. Is ivermectin effective in treating COVID-19? *Frontiers in pharmacology*. 2022;13:858693. doi:10.3389/fphar.2022.858693
214. Bitterman A, Martins CP, Cices A, Nadendla MP. Comparison of trials using ivermectin for COVID-19 between regions with high and low prevalence of strongyloidiasis: A meta-analysis. *JAMA Netw Open*. Mar 1 2022;5(3):e223079. doi:10.1001/jamanetworkopen.2022.3079

215. Ravikirti, Roy R, Pattadar C, et al. Evaluation of ivermectin as a potential treatment for mild to moderate COVID-19: A double-blind randomized placebo Controlled trial in eastern India. *J Pharm Pharm Sci.* 2021;24:343-350. doi:10.18433/jpps32105
216. Mahmud R, Rahman MM, Alam I, et al. Ivermectin in combination with doxycycline for treating COVID-19 symptoms: a randomized trial. *J Int Med Res.* May 2021;49(5):3000605211013550. doi:10.1177/03000605211013550
217. Rago Z, Toth B, Szalenko-Tokes A, et al. Results of a systematic review and meta-analysis of early studies on ivermectin in SARS-CoV-2 infection. *Geroscience.* Aug 2023;45(4):2179-2193. doi:10.1007/s11357-023-00756-y
218. Bryant A, Lawrie TA, Dowswell T, et al. Ivermectin for prevention and treatment of COVID-19 infection: A systematic review, meta-analysis, and trial sequential analysis to inform clinical guidelines. *Am J Ther.* Jun 21 2021;28(4):e434-e460. doi:10.1097/MJT.0000000000001402
219. Navarro M, Camprubi D, Requena-Mendez A, et al. Safety of high-dose ivermectin: a systematic review and meta-analysis. *J Antimicrob Chemother.* Apr 1 2020;75(4):827-834. doi:10.1093/jac/dkz524
220. Guzzo CA, Furtek CI, Porras AG, et al. Safety, tolerability, and pharmacokinetics of escalating high doses of ivermectin in healthy adult subjects. *J Clin Pharmacol.* Oct 2002;42(10):1122-33. doi:10.1177/009127002401382731
221. Varnaseri M AF, Jamshididan R, Dargahi M, Gheibi N, Abolghasemi S, Dayer M, Varnasseri N, Hoseinynejad K, Kheradhoosh S, Nazari P, Babadi E, Mousavinezhad SM, Ebrahimi P. Ivermectin as a potential addition to the limited anti-COVID-19 arsenal: A double-blinded clinical trial. *Jundishapur J Health Sci.* 2024;16(2):e146703. doi:10.5812/jjhs-146703
222. Hayward G, Yu LM, Little P, et al. Ivermectin for COVID-19 in adults in the community (PRINCIPLE): An open, randomised, controlled, adaptive platform trial of short- and longer-term outcomes. *J Infect.* Apr 2024;88(4):106130. doi:10.1016/j.jinf.2024.106130
223. Reis G, Silva E, Silva DCM, et al. Effect of early treatment with Ivermectin among patients with Covid-19. *N Engl J Med.* May 5 2022;386(18):1721-1731. doi:10.1056/NEJMoa2115869
224. Naggie S, Boulware DR, Lindsell CJ, et al. Effect of ivermectin vs placebo on time to sustained recovery in outpatients with mild to moderate COVID-19: A randomized clinical trial. *JAMA.* Oct 25 2022;328(16):1595-1603. doi:10.1001/jama.2022.18590
225. Bramante CT, Huling JD, Tignanelli CJ, et al. Randomized trial of metformin, ivermectin, and fluvoxamine for Covid-19. *N Engl J Med.* Aug 18 2022;387(7):599-610. doi:10.1056/NEJMoa2201662
226. Bramante CT, Buse JB, Liebovitz DM, et al. Outpatient treatment of COVID-19 and incidence of post-COVID-19 condition over 10 months (COVID-OUT): a multicentre, randomised, quadruple-blind, parallel-group, phase 3 trial. *Lancet Infect Dis.* Oct 2023;23(10):1119-1129. doi:10.1016/S1473-3099(23)00299-2
227. Popp M, Reis S, Schiesser S, et al. Ivermectin for preventing and treating COVID-19. *Cochrane Database Syst Rev.* Jun 21 2022;6(6):CD015017. doi:10.1002/14651858.CD015017.pub3
228. Marcolino MS, Meira KC, Guimaraes NS, et al. Systematic review and meta-analysis of ivermectin for treatment of COVID-19: evidence beyond the hype. *BMC Infect Dis.* Jul 23 2022;22(1):639. doi:10.1186/s12879-022-07589-8
229. Naggie S, Boulware DR, Lindsell CJ, et al. Effect of higher-dose ivermectin for 6 Days vs placebo on time to sustained recovery in outpatients with COVID-19: A randomized clinical trial. *JAMA.* Mar 21 2023;329(11):888-897. doi:10.1001/jama.2023.1650
230. Lopez-Medina E, Lopez P, Hurtado IC, et al. Effect of ivermectin on time to resolution of symptoms Among adults with mild COVID-19: A randomized clinical trial. *JAMA.* Apr 13 2021;325(14):1426-1435. doi:10.1001/jama.2021.3071
231. Roman YM, Burela PA, Pasupuleti V, Piscocoya A, Vidal JE, Hernandez AV. Ivermectin for the treatment of coronavirus disease 2019: A systematic review and Meta-analysis of randomized controlled trials. *Clin Infect Dis.* Mar 23 2022;74(6):1022-1029. doi:10.1093/cid/ciab591
232. E OM, Spillane S, Byrne P, O'Neill M, Harrington P, Ryan M. Interventions in an ambulatory setting to prevent progression to severe disease in patients with COVID-19: A systematic review. *Ann Pharmacother.* Mar 2022;56(3):309-318. doi:10.1177/10600280211028242
233. Xu Y, Li H, Kirui B, et al. Effectiveness of COVID-19 Vaccines over 13 Months Covering the Period of the Emergence of the Omicron Variant in the Swedish Population. *Vaccines (Basel).* Dec 5 2022;10(12)doi:10.3390/vaccines10122074
234. Bergman P, Lindh AU, Bjorkhem-Bergman L, Lindh JD. Vitamin D and respiratory tract iInfections: A systematic review and meta-analysis of randomized controlled trials. *PLoS One.* 2013;8(6):e65835. doi:10.1371/journal.pone.0065835

235. Zhang X, Song Y, Ci X, et al. Ivermectin inhibits LPS-induced production of inflammatory cytokines and improves LPS-induced survival in mice. *Inflamm Res*. Nov 2008;57(11):524-9. doi:10.1007/s00011-008-8007-8
236. Vincent MJ, Bergeron E, Benjannet S, et al. Chloroquine is a potent inhibitor of SARS coronavirus infection and spread. *Virology*. Aug 22 2005;2:69. doi:10.1186/1743-422X-2-69
237. Cao M, He C, Gong M, Wu S, He J. The effects of vitamin D on all-cause mortality in different diseases: an evidence-map and umbrella review of 116 randomized controlled trials. *Front Nutr*. 2023;10:1132528. doi:10.3389/fnut.2023.1132528
238. Caly L, Druce JD, Catton MG, Jans DA, Wagstaff KM. The FDA-approved drug ivermectin inhibits the replication of SARS-CoV-2 in vitro. *Antiviral Res*. Jun 2020;178:104787. doi:10.1016/j.antiviral.2020.104787
239. Meeus G, Van Coile F, Pottel H, et al. Efficacy and safety of in-hospital treatment of Covid-19 infection with low-dose hydroxychloroquine and azithromycin in hospitalized patients: A retrospective controlled cohort study. *New Microbes New Infect*. Oct 2023;55:101172. doi:10.1016/j.nmni.2023.101172
240. Anonymus. COVID-19 Treatment Guidelines-Vitamin D. NIH. Accessed March 17, 2024, <https://www.covid19treatmentguidelines.nih.gov/therapies/supplements/vitamin-d/>
241. Marks PW, Califf RM. Bivalent Covid-19 Vaccines. *N Engl J Med*. Mar 23 2023;388(12):1151. doi:10.1056/NEJMc2301323

Highlights:

- COVID-19 outbreaks coincided with each winter when population vitamin D levels are lowest.
- In 2021, the COVID-19 vaccine reduced SARS-CoV-2 hospitalizations; however, with emerging mutants and immune evasion, the rate of infection and hospitalization increased from late 2021.
- The efficacy of SARS-CoV-2 mRNA and bivalent vaccines against Omicron significantly reduced from late 2022 due to new mutants and immune evasion causing outbreaks.
- Vitamin D is a potent anti-inflammatory and anti-oxidant, which helps negate the adverse effects of viruses and vaccines.
- Exploring alternative cost-effective therapies, like vitamin D and ivermectin, is essential to prepare for future pandemics.

Declartation foi interests:

Conflicts of interest The author declares no conflicts of interest.
He did not receive any funding or writing assistance

Funding: None

Journal Pre-proof