

Prevalence of vitamin D deficiency among Turkish, Moroccan, Indian and sub-Sahara African populations in Europe and their countries of origin: an overview

I. M. van der Meer · B. J. C. Middelkoop ·
A. J. P. Boeke · P. Lips

Received: 25 February 2010 / Accepted: 7 April 2010 / Published online: 12 May 2010
© The Author(s) 2010. This article is published with open access at Springerlink.com

Abstract

Summary Vitamin D status of nonwestern immigrants in Europe was poor. Vitamin D status of nonwestern populations in their countries of origin varied, being either similar to the immigrant populations in Europe or higher than in European indigenous populations. Vitamin D concentrations in nonwestern immigrant populations should be improved.

Purpose The higher the latitude, the less vitamin D is produced in the skin. Most European countries are located at higher latitudes than the countries of origin of their

nonwestern immigrants. Our aim was to compare the serum 25-hydroxyvitamin D (25(OH)D) concentration of nonwestern immigrant populations with those of the population in their country of origin, and the indigenous population of the country they migrated to.

Methods We performed literature searches in the “PubMed” and “Embase” databases, restricted to 1990 and later. The search profile consisted of terms referring to vitamin D or vitamin D deficiency, prevalence or cross-sectional studies, and countries or ethnicity. Titles and abstracts were reviewed to identify studies on population-based mean serum 25(OH)D concentrations among Turkish, Moroccan, Indian, and sub-Sahara African populations in Europe, Turkey, Morocco, India, and sub-Sahara Africa.

Results The vitamin D status of immigrant populations in Europe was poor compared to the indigenous European populations. The vitamin D status of studied populations in Turkey and India varied and was either similar to the immigrant populations in Europe (low) or similar to or even higher than the indigenous European populations (high).

Conclusions In addition to observed negative consequences of low serum 25(OH)D concentrations among nonwestern populations, this overview indicates that vitamin D status in nonwestern immigrant populations should be improved. The most efficacious strategy should be the subject of further study.

Keywords Indian · Moroccan · Prevalence · Serum 25-hydroxyvitamin D · Sub-Sahara African · Turkish

I. M. van der Meer · B. J. C. Middelkoop
Department of Epidemiology,
Municipal Health Service of The Hague,
The Hague, The Netherlands

B. J. C. Middelkoop
Department of Public Health and Primary Care,
Leiden University Medical Center,
Leiden, The Netherlands

A. J. P. Boeke
Department of General Practice,
Institute for Health and Care Research (EMGO Institute),
VU University Medical Center,
Amsterdam, The Netherlands

P. Lips
Department of Internal Medicine,
Endocrine Section and Institute for Health and Care Research
(EMGO Institute), VU University Medical Center,
Amsterdam, The Netherlands

I. M. van der Meer (✉)
GGD Den Haag,
Postbus 12 652,
2500 DP The Hague, The Netherlands
e-mail: irene.vandermeer@denhaag.nl

Introduction

Vitamin D status has been found to be poor among nonwestern immigrant populations in European countries

compared to indigenous European populations [1–4]. The lower serum 25(OH)D concentrations among nonwestern immigrants compared to indigenous European populations may lead to differences in health. Consequences of vitamin D deficiency include bone- and muscle-related symptoms (e.g., bone and muscle pain), decreased muscle strength, and diseases (e.g., rickets in children; osteomalacia in adults) [5, 6]. Other possible consequences are diabetes mellitus, infectious diseases, and cancer [7].

Direct sunlight stimulates the production of vitamin D in the skin from 7-dehydrocholesterol. Other sources of vitamin D include some natural foods (e.g., fatty fish), fortified foods (e.g., margarine), and supplements. The amount of vitamin D produced through exposure to UVB radiation depends on skin type: the darker the skin, the more sunlight is required to produce a given amount of vitamin D [8–10]. Nonwestern immigrants usually have darker skin than indigenous European subjects. Therefore, they have a higher risk of lower serum 25-hydroxyvitamin D (25(OH)D) concentrations when living at the same latitude.

The duration of UVB irradiation needed to produce a certain quantity of vitamin D in a particular skin surface depends on season, time of day, and geographical location [11]. The higher the latitude, the lower the UVB intensity, and the fewer months and hours per day during which vitamin D is produced. Most European countries are located at a higher latitude than the countries of origin of nonwestern immigrants.

The threshold for vitamin D deficiency should—ideally—be based on its consequences. However, most studies of the consequences of vitamin D deficiency have been performed among older western populations in Europe and North America, rather than among adult nonwestern immigrant populations in these countries. Another means of establishing a deficiency threshold is through the use of reference values within a population [12]. For that purpose, a comparison of the vitamin D status of nonwestern immigrant populations with the populations in their countries of origin might be more suitable than a comparison with the indigenous western populations. Our aim was to compare the vitamin D status of nonwestern immigrant populations with both the populations in their countries of origin and the populations in the country they migrated to. Additionally, we wanted to identify what determinants were mentioned to explain differences in vitamin D status between subgroups in the studied populations.

Methods

We performed literature searches in the “PubMed” and “Embase” databases. The search profile consisted of terms referring to vitamin D or vitamin D deficiency, prevalence

or cross-sectional studies, and countries or ethnicity. The search was restricted to publications from 1990 onwards; about 1,000 were returned. Titles and abstracts were reviewed to identify studies on population-based mean serum 25(OH)D concentrations among Turkish, Moroccan, Indian, and sub-Sahara African populations in Europe, Turkey, Morocco, India, or sub-Sahara Africa. We accepted the definitions of ethnicity as used in the identified articles.

We extracted data for the Turkish, Moroccan, Indian, and sub-Sahara African populations and for the indigenous European populations if this group was included in the studies performed in Europe. From suitable publications, we extracted information about geographical location and season of data collection, age and gender of the study population, duration of pregnancy if applicable, number of subjects, mean serum 25(OH)D concentration with standard deviation, percentage of subjects with serum 25(OH)D <25 nmol/l, and determinants of serum 25(OH)D concentration. Specific characteristics of the study population which could influence the vitamin D status, such as clothing habits, were also extracted. Of identified intervention studies, we used only data from baseline measurements. Serum 25(OH)D concentrations presented in nanogram per milliliter or microgram per liter were transformed into nanomole per liter. Data variances presented as standard errors or 95% confidence intervals were converted to standard deviations. When either vitamin D status parameter (mean and % <25 nmol/l) was not presented, another measure for vitamin D status (such as median concentration or % below another threshold) was extracted.

Results

Prevalence

The identified studies on Turkish populations in Europe are presented in Table 1 and on Turkish populations in Turkey in Table 2. The vitamin D status was lower in the Turkish groups in Europe than in the indigenous European groups. Vitamin D status in the Turkish groups in Turkey varied widely. The subgroups with covering clothes had the lowest serum 25(OH)D concentrations (mean 10 nmol/l) [13, 14]. Turkish elderly living in their own homes (mean 158 nmol/l for males and 103 nmol/l for females) and Turkish unveiled adult women (mean 135 nmol/l)—all of whom were measured at the end of summer—had the highest serum 25(OH)D concentrations [15, 16].

Studies on Moroccan populations in Europe are presented in Table 3. Table 4 presents the only study found on the vitamin D status of a Moroccan population in Morocco. As was the result among Turkish populations, the Moroccan populations in Europe had lower serum 25(OH)D concentrations than the indigenous European populations. The

Table 1 Studies among Turkish populations in Europe

Study	Study characteristics	Study population	Serum 25(OH)D (nmol/l) Mean±SD ^a	Determinants for lower serum 25(OH)D
Adults				
Madar et al. [39]	Norway, Oslo (60° N), all year round	Turkish F, mean 27 years (<i>n</i> =25)	26±14, 56% < 25	No daily use of vitamin D supplementation, education <10 years
Holvik et al. [33]	Norway, Oslo (60° N), in spring	Turkish M, mean 39 years for all men (<i>n</i> =87) Turkish F, mean 37 years for all women (<i>n</i> =101)	median 33, 23% < 25 median 26, 46% < 25	Female gender, younger age, April/May blood sample (compared to June), lower use of cod liver oil supplements, lower intake of fatty fish; in females: higher BMI, shorter educational length
Van der Meer et al. [1]	The Netherlands, Amsterdam, The Hague, Amersfoort, and Haarlem (52° N), all year round	Dutch M (40%)+F, median 45 years (<i>n</i> =102) Turkish M (41%)+F, median 35 years (<i>n</i> =121)	median 67, 06% < 25 median 27, 41% < 25	Autumn or winter season, pregnant or breastfeeding, lower consumption of fatty fish, no use of vitamin D supplements, smaller area of uncovered skin, no use of tanning bed, lower consumption of margarine, no preference for sun
Grooijans-Geerts and Wielders [25]	The Netherlands, Amersfoort, end of winter	Dutch F, mean 44 years (<i>n</i> =32) Turkish veiled F, mean 30 years (<i>n</i> =51)	28% < 30 90% < 30	–
Erkal et al. [2]	Germany, Giessen (50° N), end of winter	German M (50%)+F, 19–63 years (<i>n</i> =101) Turkish M, 18–69 years (<i>n</i> =270)	29% < 50 Median 40	Female gender, veiling, having three or more children, living at higher latitude, higher BMI
Moreno-Reyes et al. [3]	Belgium, Brussels, all year round.	Turkish F, 16–67 years (<i>n</i> =296) Belgian M (50%)+F, mean 52 years (<i>n</i> =100) Turkish M (50%)+F, mean 49 years, first-generation immigrants (<i>n</i> =101)	Median 31 49±22, 13% < 25 31±20, 53% < 25	Winter season, male gender
Pregnant women				
Van der Meer et al. [26]	The Netherlands, The Hague (52° N), at the first antenatal visit (12th week), all year round	Western, mean 30 years (<i>n</i> =105) Turkish, mean 24 years (<i>n</i> =79)	53±22, 08% < 25 15±12, 84% < 25	–
Children				
Madar et al. [39]	Norway, Oslo (60° N), all year round	Turkish M+F, mean 7 weeks (<i>n</i> =25)	37±38, 56% < 25	Exclusively breastfed infants (no supplements)
Meulmeester et al. [27]	The Netherlands, The Hague, or Rotterdam, at the end of winter or the end of spring	Caucasian M (50%)+F, 8 years, The Hague, end of winter (<i>n</i> =39) Turkish M (50%)+F, 8 years, The Hague, end of winter (<i>n</i> =40) Caucasian M (50%)+F, 8 years, Rotterdam, end of spring (<i>n</i> =40) Turkish M (50%)+F, 8 years, Rotterdam, end of spring (<i>n</i> =40)	57±16 23±10 73±14 37±13	End of winter measurement, lower cumulative global sun radiation

SD standard deviation

^a Unless mentioned otherwise

Table 2 Studies among Turkish populations in Turkey

Study	Study characteristics	Study population	Serum 25(OH)D (nmol/l) Mean±SD ^a	Determinants for lower serum 25(OH)D
Adults				
Erkal et al. [2]	Turkey, Mersin (36° N), Ankara (40° N), Istanbul and Unye (42° N), end of winter	Turkish M, 21–66 years (<i>n</i> =85) Turkish F, 17–69 years (<i>n</i> =242)	Median 47 Median 36	Female gender, veiling, having three or more children, living at higher latitude, higher BMI
Guzel et al. [16]	Turkey (37° N), end of summer	Turkish F, mean 25 years, veiled (<i>n</i> =30) Turkish F, mean 25 years, unveiled (<i>n</i> =30)	83±40 135±68	Veiling, lower exposure to sunlight, longer duration of being veiled
Alagol et al. [13]	Turkey, Istanbul (41° N), in summer	Turkish F, mean 24 years, dressed with usual areas exposed to the sun (<i>n</i> =18) Turkish F, mean 28 years, traditional clothing, hands and face uncovered (<i>n</i> =15) Turkish F, mean 26 years, traditional clothing, covering whole body including hands and face (<i>n</i> =15)	32±24 10±06	Covering clothes/veiling
Atli et al. [15]	Turkey, Ankara (40° N), at the end of summer	Turkish M, mean 73 years, own home (<i>n</i> =24) Turkish F, mean 72 years, own home (<i>n</i> =171) Turkish M, mean 76 years, old age home (<i>n</i> =87) Turkish F, mean 75 years, old age home (<i>n</i> =138)	158±108 103±98 94±72 62±74	Female gender, living in old age home, older age, lower benefit from ultraviolet index (ratio of points for sunlight exposure and covering clothes)
Pregnant women				
Pehlivan et al. [14]	Turkey, Last trimester	Turkish, total group (<i>n</i> =78) Turkish, with covered head and hands, not the face (<i>n</i> =4) Turkish, with covered head, not the hands or face (<i>n</i> =49) Turkish, with no cover on head, hands or face (<i>n</i> =25)	18±10, 80% < 25 10±05 17±10 20±10	Low educational level, insufficient intake of vitamin D within diet, “covered” dressing habits
Children				
Olmez et al. [34]	Turkey, Izmir, end of summer or end of winter	Turkish F, 14–18 years, low socioeconomic status, end of summer (<i>n</i> =32) Turkish F, 14–18 years, high socioeconomic status, end of summer (<i>n</i> =32) Turkish F, 14–18 years, low socioeconomic status, end of winter (<i>n</i> =30) Turkish F, 14–18 years, high socioeconomic status, end of winter (<i>n</i> =30)	52±23 65±29 34±16 59±24	End of winter measurement, low socioeconomic status

SD standard deviation

^a Unless mentioned otherwise

Table 3 Studies among Moroccan populations in Europe

Study	Study characteristics	Study population	Serum 25(OH)D (nmol/l) Mean±SD ^a	Determinants for lower serum 25(OH)D
Adults				
Van der Meer et al. [1]	The Netherlands, Amsterdam, The Hague, Amersfoort, and Haarlem (52° N)	Dutch M (40%)+F, median 45 years (<i>n</i> =102) Moroccan M (41%)+F, median 38 years (<i>n</i> =96) Median 30, 37% < 25	Median 67, 06% < 25	Autumn or winter season, pregnant or breastfeeding, lower consumption of fatty fish, no use of vitamin D supplements, smaller area of uncovered skin, no use of tanning bed, lower consumption of margarine, no preference for sun
Moreno-Reyes et al. [3]	Belgium, Brussels, all year round	Belgian M (50%)+F, mean 52 years (<i>n</i> =100) Moroccan M (50%)+F, mean 49 years, first-generation immigrants (<i>n</i> =100)	49±22, 13% < 25 27±17, 54% < 25	Winter season, male gender
Pregnant women				
Van der Meer et al. [26]	The Netherlands, The Hague (52° N), at the first antenatal visit (12th week)	Western, mean 30 years (<i>n</i> =105) Moroccan, mean 26 years (<i>n</i> =69)	53±22, 08% < 25 20±14, 81% < 25	–
Children				
Meulmeester et al. [27]	The Netherlands, The Hague or Rotterdam, at the end of winter or the end of spring	Caucasian M (50%)+F, 8 years, The Hague, end of winter (<i>n</i> =39) Moroccan M (50%)+F, 8 years, The Hague, end of winter (<i>n</i> =38) Caucasian M (50%)+F, 8 years, Rotterdam, end of spring (<i>n</i> =40) Moroccan M (50%)+F, 8 years, Rotterdam, end of spring (<i>n</i> =42)	57±16 30±14 73±14 38±14	End of winter measurement, lower cumulative global sun radiation

SD standard deviation

^a Unless mentioned otherwise

Table 4 Studies among Moroccan populations in Morocco

Study	Study characteristics	Study population	Serum 25(OH)D (nmol/l) Mean±SD ^a	Determinants for lower serum 25(OH)D
Adults				
Allali et al. [17]	Morocco, Rabat, in the end of winter	Moroccan F, mean 50 years, total group (n=415)	45±20	Age>55 years, calcium intake<700 mg/d, wearing a veil, sunlight exposure<30 min/day
		Moroccan F, mean 43 years, premenopausal (n=108)	47±19	
		Moroccan F, mean 56 years, postmenopausal (n=307)	44±20	

SD standard deviation

^a Unless mentioned otherwise

Moroccan adult women in Morocco, who were measured at the end of winter, had a mean serum 25(OH)D concentration of 45 nmol/l [17]. This was lower than the indigenous population in the Netherlands (median 67 nmol/l) and in Belgium (mean 49 nmol/l) [1, 3]. The Dutch and Belgian populations consisted of both men and women, and these were measured year-round, which might explain the difference.

Studies on adult Indian (or South Asian) populations in Europe also found lower serum 25(OH)D concentrations in comparison to indigenous European populations (Table 5). Concerning pregnant women and children, we did not identify any studies which included an indigenous European population. The vitamin D status among various Indian populations in India differed (Table 6). Some populations with limited sunlight exposure, such as physicians and nurses (mean 8 nmol/l in winter and 18 nmol/l in summer) or Delhi-based males (mean 18 nmol/l) and females (mean 17 nmol/l) measured in winter, had low serum 25(OH)D concentrations, similar to Indian populations in Europe [18, 19]. Other, mainly rural, Indian adult populations in India had higher serum 25(OH)D concentrations [20, 21].

Sub-Saharan Africans in the Netherlands—consisting predominantly of Ghanaians and Somalis—had a median serum 25(OH)D concentration of 33 nmol/l (n=57) [1]. Congolese immigrants in Belgium had a mean serum 25(OH)D concentration of 38 nmol/l (standard deviation (SD) of 14 nmol/l). We did not identify any studies on vitamin D status in Ghana, Somalia, or the Democratic Republic of Congo. Studies in sub-Saharan countries include a study in Cameroon, with a mean serum 25(OH)D concentration of 53 nmol/l (SD 19 nmol/l) among an older population (aged 60–86 years), a study on Nigerian children (6–35 months) with a mean serum 25(OH)D concentration of 64 nmol/l (sd 23 nmol/l), and a study on Gambian women aged 25 or older, with a mean serum 25(OH)D concentration between 73 nmol/l (SD 20 nmol/l) and 113 nmol/l (SD 27 nmol/l), varying with age [22–24].

In all studies performed in Europe where both groups were included, immigrant groups in European countries had significantly lower serum 25(OH)D concentrations than indigenous European groups [1–4, 25–32].

Determinants

In the last column of each table, the determinants for a lower 25(OH)D concentration are presented. As expected, many studies found a lower exposure to sunlight (e.g., behavior or season) [1–3, 13–18, 27, 32–38] or a restricted intake of vitamin D (via food or supplements) [1, 14, 17, 33, 39, 40], to be associated with a lower serum 25(OH)D concentration.

Neither gender nor age were unambiguously associated with the serum 25(OH)D concentration. Female gender was found to be a determinant for a low serum 25(OH)D concentration [2, 4, 15, 33, 35, 36, 41, 42], but not in all studies that compared males and females [3, 19, 20, 31, 41, 43]. Both a younger age [33] and an older age [15, 17] were associated with a lower serum 25(OH)D concentration.

Other determinants of lower serum 25(OH)D concentrations—explained by association with exposure to sunlight or dietary habits—are a lower socioeconomic position [34, 42], a shorter duration of education [33, 39], or a lower educational level [14], living in an urban environment [20, 21], and an earlier start time to the workday [44].

In newborn children, a mother's lower serum 25(OH)D concentration was associated with a lower serum 25(OH)D concentration in the child [18, 45, 46].

Discussion

The vitamin D status of Turkish, Moroccan, Indian, and sub-Sahara African immigrant populations in Europe was poor compared to the indigenous European populations. The vitamin D states of studied populations in Turkey, Morocco, and India varied between concentrations similar

Table 5 Studies among Indian populations in Europe

Study	Study characteristics	Study population	Serum 25(OH)D (nmol/l) Mean±SD ^a	Determinants for lower serum 25(OH)D
Adults				
Brooke-Wavell et al. [28]	United Kingdom	White European F, mean 59 years (<i>n</i> =23) South Asian F, mean 59 years (Bangladeshi, Indian <i>n</i> =24)	76±18 33±13	–
Ward et al. [29]	United Kingdom, Manchester	White Caucasian European F, mean 30 years (<i>n</i> =96) Pakistani muslim or Gujarati Hindu F, mean 29 years (<i>n</i> =95)	67±34 20±12	–
Ford et al. [4]	United Kingdom, Birmingham, end of summer.	Caucasian M+F, mean 59 years (1–92 years; <i>n</i> =317) Asian M+F, mean 47 years (2–87years) (<i>n</i> =251)	58±31, 12% < 25 36±26, 31% < 25	In the Asian group: female gender
Hamson et al. [30]	United Kingdom, Leicester	White M, 33 years (<i>n</i> =37) White F, 32 years (<i>n</i> =51)	3% < 12.5 0% < 12.5	–
Solanki et al. [31]	United Kingdom, Birmingham, end of winter.	Gujarati M, 34 years (Gujarat region India; <i>n</i> =42)	60% < 12.5	–
		Gujarati F, 34 years (Gujarat region India; <i>n</i> =71)	51% < 12.5	–
Finch et al. [32]	United Kingdom, London, all year round.	White M, <65 years, mean 30 years men and women (<i>n</i> =4)	28±12	–
		White F, <65 years, mean 30 years men and women (<i>n</i> =12)	48±29	–
		White M, >65 years, mean 74 years men and women (<i>n</i> =4)	55±14	–
		White F, >65 years, mean 74 years men and women (<i>n</i> =14)	40±21	–
		Asian M, <65 years, mean 31 years men and women (<i>n</i> =3)	16±08	–
		Asian F, <65 years, mean 31 years men and women (<i>n</i> =3)	21±07	–
		Asian M, >65 years, mean 72 years men and women (<i>n</i> =21)	13±09	–
		Asian F, >65 years, mean 72 years men and women (<i>n</i> =16)	23±20	–
		White M (50%)±F, mean 39 years, winter (<i>n</i> =30)	39±18	Winter season (March/April), vegetarian, Hindu religion, Muslim religion (only in winter); Hindus seasonal responses are blunted, resulting in significantly lower peak values than for whites or non-vegetarian (Muslim) Asians
		White M (50%)±F, mean 39 years, summer (<i>n</i> =18)	65±27	–
Van der Meer et al. [1]	The Netherlands, Amsterdam, The Hague, Amersfoort and Haarlem (52°N)	Asian M (70%)±F, mean 42 years, non-vegetarians, winter (<i>n</i> =116)	19±13	–
		Asian M (70%)±F, mean 42 years, non-vegetarians, summer (<i>n</i> =22)	45±24	–
		Asian M (40%)±F, mean 42 years, vegetarians, winter (<i>n</i> =29)	10±8	–
		Asian M (40%)±F, mean 42 years, vegetarians, summer (<i>n</i> =16)	27±21	–
Pregnant women	The Netherlands, Amsterdam, The Hague, Amersfoort and Haarlem (52°N)	Dutch M (40%)±F, median 45 years (<i>n</i> =102)	Median 67, 06% < 25	Autumn or winter season, pregnant or breastfeeding, lower consumption of fatty fish, no use of vitamin D supplements, smaller area of uncovered skin, no use of tanning bed, lower consumption of margarine, no preference for sun
		Surinam South Asian M (37%)±F, median 41 years (<i>n</i> =107)	Median 24, 51% < 25	–
Datta et al. [63]	United Kingdom, Cardiff (51.5°N), at booking visit	Indian subcontinent (<i>n</i> =100)	52% < 20	Being in Britain for more than 3 years (compared to less than 3 years and to being born in Britain)
Children				
Lawson and Thomas [40]	UK, autumn	Bangladeshi M±F, 2 years (<i>n</i> =139) Pakistani M±F, 2 years (<i>n</i> =200)	42±21, 20% < 25 36±20, 34% < 25	Failure to take a vitamin supplement.
Koch and Burmeister [64]	Germany, in summer	Indian M±F, 2 years (<i>n</i> =279)	42±23, 25% < 25	–
		Asian M (33%)±F, 3–17 years (Birma, Sri Lanka, India; <i>n</i> =9)	28±09, 44% < 25	–

SD standard deviation

^a Unless mentioned otherwise

Table 6 Studies among Indian populations in India

Study	Study characteristics	Study population	Serum 25(OH)D (nmol/l) Mean±SD ^a	Determinants for lower serum 25(OH)D
Adults				
Goswami et al. [19]	India, Delhi, in winter	Adult M, mean 31 years (<i>n</i> =244) Adult F, mean 35 years (<i>n</i> =398)	18±9 17±11	–
Goswami et al. [41]	India, Agota village (29° N), in winter	Adult M, rural, mean 43 years (<i>n</i> =32) Adult F, rural, mean 43 years (<i>n</i> =25)	44±24 27±16	Female gender
Harinarayan et al. [20]	India, Tirupati (13° N)	Adult M, urban, mean 46 years for urban M+F (<i>n</i> =134) Adult M, urban, mean 43 years for urban M+F (<i>n</i> =109) Adult F, urban, mean 46 years for urban M+F (<i>n</i> =807) Adult F, rural, mean 43 years for rural M+F (<i>n</i> =96)	46±22 59±20 39±20 48±22	Urban subject Lower exposure to sunlight, female gender
Zargar et al. [35]	India, Kashmir valley, all year round	Indian M, mean 29 years (<i>n</i> =64) Indian F, mean 27 years (<i>n</i> =28)	38±30, 41% < 25 14±11, 96% < 25	–
Gulvady et al. [44]	India, Mumbai	Indian M, 40–68 years, senior executives (indoor workers; <i>n</i> =86)	28% < 19	Earlier start of the workday
Vupputuri et al. [43]	India, Delhi (28° N)	Asian Indian M, mean 43 years (for both men and women), urban, middle income, mostly housewives (<i>n</i> =54) Indian F, mean 54 years, postmenopausal (<i>n</i> =164)	27±17 22±12	–
Harinarayan [65]	India, Tirupati (13° N), all year round	Indian F, mean 43 years (for both men and women), urban, middle income, mostly housewives (<i>n</i> =54) Indian F, mean 54 years, postmenopausal (<i>n</i> =164)	37±18, 30% < 25	Higher dietary calcium intake, higher dietary phytate intake, higher phytate to calcium ratio
Harinarayan et al. [21]	India, around Tirupati (13° N), winter to summer (Jan–Jul)	Indian, mean 44 years, rural (<i>n</i> =191) Indian, mean 46 years, urban (<i>n</i> =125)	53±06, 03% < 25 34±07, 35% < 25	Urban subject, lower dietary calcium intake, higher phytate to calcium ratio
Goswami et al. [18]	India, Delhi (28° N), in winter or summer	Indian M, mean 25 years, soldiers, winter (<i>n</i> =31) Indian M (58%)+F, mean 23 years, physicians and nurses, winter (<i>n</i> =19) Indian M (67%)+F, mean 43 years, depigmented persons, winter (<i>n</i> =15) Indian M (58%)+F, mean 24 years, physicians and nurses, summer (<i>n</i> =19)	47±12 08±03 18±11 18±08	Less exposure to sunlight, more skin pigmentation, winter season
Pregnant women				
Sahu et al. [36]	India, Barabanki district, 32 km from Lucknow (27°), all year round	Indian, rural, mean 27 years (<i>n</i> =139)	38±20, 32% < 25	Lower summer sun exposure, measurement in winter
Farrant et al. [66]	India, Mysore (South India) at the 30th week of pregnancy	Indian, mean 24 years (<i>n</i> =559)	Median 38, 31% < 28 nmol/l	Taking calcium and vitamin D at recruitment, measurement in Mar–Aug
Bhalala et al. [45]	Western India, at the 37th week of pregnancy, all year round	Indian, 20–35 years, middle income group (<i>n</i> =42) Cord blood (<i>n</i> =42)	57±27 48±24	Lower serum 25(OH)D in mother → lower serum 25(OH)D in cord blood
Sachan et al. [46]	India, Lucknow (27° N), before labor, autumn	Indian, total group (<i>n</i> =207) Indian, urban (<i>n</i> =140) Indian, rural (<i>n</i> =67)	43% < 25 35±24 35±22	–
Goswami et al. [18]	India, Delhi (28° N), in summer	Indian, mean 23 years, poor socioeconomic class (<i>n</i> =29)	22±11	–
Children				
Sahu et al. [36]	India, Barabanki district, 32 km from Lucknow (27° N), all year round	Indian F, rural, mean 14 years, total group (<i>n</i> =121) Indian M, mean 14 years, brothers of the 28 girls, in winter (<i>n</i> =34)	33±16, 34% < 25 68±29, 36% < 25	Lower summer sun exposure, female gender, measurement in winter

Puri et al. [37]	India, Dehli (28° N), in summer	Indian F, mean 14 years, sisters of the 34 boys, in winter (<i>n</i> =28)	31±14, 03% < 25	Higher BMI, lower sun exposure, smaller percentage of body surface area exposed
Harinarayan et al. [20]	India, Tirupati (13° N)	Indian F, mean 12 years (6–18), lower socioeconomic strata (<i>n</i> =193)	35±17, 31% < 25	
Bhalala et al. [45]	Western India, all year round	Indian F, mean 12 years (6–18), upper socioeconomic strata (<i>n</i> =211)	29±13, 39% < 25	
Khadilkar et al. [67]	India, Pune (18° N), in winter	Indian M, urban, mean 13 years for urban M+F (<i>n</i> =30)	39±17	
Sivakumar et al. [68, 69]	India, Hyderabad, end of winter, summer (Mar and Jul)	Indian M, rural, mean 13 years for rural M+F (<i>n</i> =34)	43±22	
Marwaha et al. [42]	India, New Dehli (28° N)	Indian F, urban, mean 13 years for urban M+F (<i>n</i> =39)	46±28	
Sachan et al. [46]	India, Lucknow (27° N), autumn	Indian F, rural, mean 13 years for rural M+F (<i>n</i> =36)	48±23	Lower serum 25(OH)D in mother
Tiwari and Puliyeel [70]	India, Dehli, in winter or summer	Indian, 3 months, exclusively breast fed, from middle income mothers (<i>n</i> =35)	45±24	
Agarwal et al. [38]	India, Dehli (28° N), end of winter	Post-menarchal F, mean 15 years (<i>n</i> =50)	70% < 30	
Goswami et al. [18]	India, Dehli (28° N), in summer	Indian M+F, 6–18 years, middle income, semi-urban (<i>n</i> =328)	26% < 25	Female gender, lower socioeconomic status
		Indian M, 10–18 years (<i>n</i> =325)	27% < 22.5	
		Indian F, 10–18 years (<i>n</i> =435)	42% < 22.5	
		Indian M (39%)+F, 10–18 years, low socioeconomic group (<i>n</i> =430)	42% < 22.5	
		Indian M (48%)+F, 10–18 years, upper socioeconomic group (<i>n</i> =330)	27% < 22.5	
		Indian neonates (cord blood, <i>n</i> =207)	21±14	Lower serum 25(OH)D in mother
		9–30 months, Sundermagari area, winter (<i>n</i> =47)	96±26	
		9–30 months, Rajiv Colony area, winter (<i>n</i> =49)	24±27	
		9–30 months, Rajiv Colony area, summer (<i>n</i> =48)	18±22	
		9–30 months, Gurgaon area, summer (<i>n</i> =52)	19±20	Atmospheric pollution
		Mean 16 months (9–24), Mori Gate area (high pollution; <i>n</i> =26)	31±18	
		Mean 16 months (9–24), Gurgaon area (low pollution; <i>n</i> =31)	68±18	
		Indian M (55%)+F, newborns from mothers from poor socioeconomic class (<i>n</i> =29) Cord blood	17±05	Lower serum 25(OH)D in mother

SD standard deviation

^a Unless mentioned otherwise

to the immigrant populations in Europe (low) and concentrations similar to or higher than the European indigenous populations (high). Determinants of the serum 25(OH)D concentration included both sources of vitamin D: exposure to sunlight and intake of vitamin D.

Gender and age were each associated with serum 25(OH)D concentration in both directions. Differences according to gender and age group could be the result of biological differences but might also reflect behavioral differences; dress style (e.g., wearing a veil) is often mentioned as a reason for a higher prevalence of vitamin D deficiency among women than men. A lower serum 25(OH)D concentration among older participants can partly be the result of the lower capacity of the skin to produce vitamin D after exposure to sunlight. The study that found lower serum 25(OH)D concentrations at younger ages [33] might have had a study population that was too young to find an effect of a lower skin capacity (their mean age was below 40 years).

As the described studies were observational, not all determinants could be studied due to a lack of variation in the determinants. For instance, Sahu et al. described the dietary intake in rural India as remarkably monotonous from meal to meal, with a low consumption of dairy and foods containing reasonable amounts of vitamin D [36]. As a consequence, it is difficult to find an association between dietary intake and serum 25(OH)D.

The darker skin types of the immigrant populations are a suitable protection against the intensity and amount of sunlight in their countries of origin, while they are a risk factor for vitamin D deficiency in northerly European countries. The serum 25(OH)D concentrations of the populations in the country of origin may, therefore, indicate normal or reference concentrations. However, those populations may themselves be deficient or suffer from insufficient concentrations as a whole. Given that until recently, mankind lived and worked outside, the serum 25(OH)D concentrations of groups who currently spend much of their time outdoors might, therefore, be considered “normal” [47]. Serum 25(OH)D concentrations of rural populations, who are expected to have a greater exposure to sunlight as a result of their agricultural occupation than urban populations [20, 21], might be a more suitable indicator of normal concentrations than those of total populations.

The high (>100 nmol/l) serum 25(OH)D concentrations in subgroups of two Turkish studies, which were performed at the end of the summer, suggest a large impact of sunlight. As sun exposure does not lead to toxic vitamin D concentrations due to a feedback mechanism, these serum 25(OH)D concentrations are expected to be within the normal or reference range, which is an additional argument that the low serum 25(OH)D concentrations (found in immigrant populations) can be interpreted as a deficiency.

Of course, assay differences might also explain part of the difference with other studies.

Symptomatic vitamin D deficiency is also suggested by the prevalence of rickets in Turkey, India, and some African countries [48–53]. The incidence of rickets in Eastern Turkey declined from 6.09% to 0.099% after a nationwide free vitamin D supplementation program [54]. Within European countries, rickets is not highly prevalent, but immigrant populations are groups at risk [55–57]. Additionally, although most nonwestern immigrant populations are younger than the indigenous European populations, cases of osteomalacia in nonwestern immigrants have been observed [58, 59]. Finch et al. found all but one case of osteomalacia within the vegetarian Asian group in England, the group with lowest vitamin D concentrations in their study [32]. Furthermore, osteoporotic and peripheral fractures were found in the vitamin-D-deficient subgroup in Morocco [17]. Erkal et al. found that 61% of the Turkish group (in Turkey) and 55% of the Turkish immigrant group in Germany complained of bone pain and/or nonspecific generalized muscle aches and pain, while it was 15% within the German group with higher serum 25(OH)D concentrations [2]. However, one should keep in mind that serum 25(OH)D is not the sole determinant of rickets; calcium intake is also important [48, 60, 61].

The comparison of serum 25(OH)D concentrations of the various populations in this article has some limitations.

First, several studies present the prevalence of vitamin D deficiency but have excluded individuals using drugs or medication known to affect bone metabolism, those recently treated for vitamin D deficiency, or those who used vitamin D supplements [1, 2, 4, 14–17, 19, 28, 35, 37, 41–43]. Medications that affect bone metabolism include, among others, vitamin D and calcium. One can argue whether the presented values represent the real prevalence in the respective populations when these individuals are excluded. However, we expect the number of excluded individuals to be small and, therefore, not of great influence on the prevalence. Furthermore, it implies that the prevalence is applicable for an apparently healthy population.

Second, the season of blood sampling varies, and this might account for a part of the observed differences between studies, because the intensity of sunlight and the amount of sunlight per day varies between seasons. These differences may be larger when studies in European countries are part of the comparison, because seasonal differences in sunlight are expected to be higher in countries at higher latitudes. For that reason, the time of year was mentioned in the tables.

Third, the comparison is hampered because the serum 25(OH)D assessment methods differ, which may influence differences between groups [62]. In addition, the level of accuracy of studies within Europe and in the country of

origin might differ. However, although we could not adjust for this type of bias, we presume that the differences will not be systematic or large enough to substantially alter the conclusions.

Finally, in comparing the various populations, it is important to realize that the social conditions of the immigrants might not be the same as those of the original populations. The cultural habits (skin-covering clothes, sun exposure, diet) might also change after immigration, particularly among the second generation.

Serum 25(OH)D concentrations of nonwestern immigrants in Europe and of subgroups of Turkish, Moroccan, Indian, and sub-Saharan countries are low. Ways to increase the serum 25(OH)D concentration include increased exposure to sunlight and increased intake of products that contain vitamin D. The strategy to effectuate these increases will differ in the various countries and populations and should be the subject of further study. These studies should ideally include measures of health to support the need for this increase in serum 25(OH)D.

Acknowledgement We gratefully acknowledge René Otten of the VU University Medical Library for his assistance in searching the PubMed and Embase databases.

Conflicts of interest None

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- van der Meer IM, Boeke AJ, Lips P, Grootjans-Geerts I, Wuister JD, Deville WL, Wienders JP, Bouter LM, Middelkoop BJ (2008) Fatty fish and supplements are the greatest modifiable contributors to the serum 25-hydroxyvitamin D concentration in a multiethnic population. *Clin Endocrinol (Oxf)* 68:466–472
- Erkal MZ, Wilde J, Bilgin Y, Akinci A, Demir E, Bodeker RH, Mann M, Bretzel RG, Stracke H, Holick MF (2006) High prevalence of vitamin D deficiency, secondary hyperparathyroidism and generalized bone pain in Turkish immigrants in Germany: identification of risk factors. *Osteoporos Int* 17:1133–1140
- Moreno-Reyes R, Carpentier YA, Boelaert M, El Moumni K, Dufourny G, Bazelmans C, Leveque A, Gervy C, Goldman S (2009) Vitamin D deficiency and hyperparathyroidism in relation to ethnicity: a cross-sectional survey in healthy adults. *Eur J Nutr* 48:31–37
- Ford L, Graham V, Wall A, Berg J (2006) Vitamin D concentrations in an UK inner-city multicultural outpatient population. *Ann Clin Biochem* 43:468–473
- Lips P (2006) Vitamin D physiology. *Prog Biophys Mol Biol* 92:4–8
- Eriksen EF, Glerup H (2002) Vitamin D deficiency and aging: implications for general health and osteoporosis. *Biogerontology* 3:73–77
- Holick MF (2007) Vitamin D deficiency. *N Engl J Med* 357:266–281
- Holick MF, MacLaughlin JA, Doppelt SH (1981) Regulation of cutaneous previtamin D3 photosynthesis in man: skin pigment is not an essential regulator. *Science* 211:590–593
- Clemens TL, Adams JS, Henderson SL, Holick MF (1982) Increased skin pigment reduces the capacity of skin to synthesise vitamin D3. *Lancet* 1:74–76
- Matsuoka LY, Wortsman J, Haddad JG, Hollis BW (1990) Skin types and epidermal photosynthesis of vitamin D3. *J Am Acad Dermatol* 23:525–526
- Holick MF (1995) Environmental factors that influence the cutaneous production of vitamin D. *Am J Clin Nutr* 61:638S–645S
- Lips P (2005) How to define normal values for serum concentrations of 25-hydroxyvitamin D? An overview. In: Feldman D, Pike W, Glorieux FH (eds) *vitamin D*, 2nd edn. Elsevier, Amsterdam, pp 1019–1028
- Alagol F, Shihadeh Y, Boztepe H, Tanakol R, Yarman S, Azizlerli H, Sandalci O (2000) Sunlight exposure and vitamin D deficiency in Turkish women. *J Endocrinol Invest* 23:173–177
- Pehlivan I, Hatun S, Aydogan M, Babaoglu K, Gokalp AS (2003) Maternal vitamin D deficiency and vitamin D supplementation in healthy infants. *Turk J Pediatr* 45:315–320
- Atli T, Gullu S, Uysal AR, Erdogan G (2005) The prevalence of Vitamin D deficiency and effects of ultraviolet light on Vitamin D levels in elderly Turkish population. *Arch Gerontol Geriatr* 40:53–60
- Guzel R, Kozanoglu E, Guler-Uysal F, Soyupak S, Sarpel T (2001) Vitamin D status and bone mineral density of veiled and unveiled Turkish women. *J Womens Health Gend Based Med* 10:765–770
- Allali F, El Aichaoui S, Khazani H, Benyahia B, Saoud B, El Kabbaj S, Bahiri R, Abouqal R, Hajjaj-Hassouni N (2009) High prevalence of hypovitaminosis D in Morocco: relationship to lifestyle, physical performance, bone markers, and bone mineral density. *Semin Arthritis Rheum* 38:444–451
- Goswami R, Gupta N, Goswami D, Marwaha RK, Tandon N, Kochupillai N (2000) Prevalence and significance of low 25-hydroxyvitamin D concentrations in healthy subjects in Delhi. *Am J Clin Nutr* 72:472–475
- Goswami R, Marwaha RK, Gupta N, Tandon N, Sreenivas V, Tomar N, Ray D, Kanwar R, Agarwal R (2009) Prevalence of vitamin D deficiency and its relationship with thyroid autoimmunity in Asian Indians: a community-based survey. *Br J Nutr* 102:382–386
- Harinarayan CV, Ramalakshmi T, Prasad UV, Sudhakar D (2008) Vitamin D status in Andhra Pradesh: a population based study. *Indian J Med Res* 127:211–218
- Harinarayan CV, Ramalakshmi T, Venkataprasad U (2004) High prevalence of low dietary calcium and low vitamin D status in healthy south Indians. *Asia Pac J Clin Nutr* 13:359–364
- Njemini R, Meyers I, Demanet C, Smits J, Sosso M, Mets T (2002) The prevalence of autoantibodies in an elderly sub-Saharan African population. *Clin Exp Immunol* 127:99–106
- Pfützner MA, Thacher TD, Pettifor JM, Zoakah AI, Lawson JO, Isichei CO, Fischer PR (1998) Absence of vitamin D deficiency in young Nigerian children. *J Pediatr* 133:740–744
- Aspray TJ, Yan L, Prentice A (2005) Parathyroid hormone and rates of bone formation are raised in perimenopausal rural Gambian women. *Bone* 36:710–720
- Grootjans-Geerts I, Wienders JP (2002) A pilot study of hypovitaminosis D in apparently healthy, veiled, Turkish women: severe vitamin D deficiency in 82% [In Dutch: Pilotonderzoek naar hypovitaminose D bij ogenschijnlijk gezonde gesluerde Turkse vrouwen: ernstige vitamine D-deficiëntie bij 82%]. *Ned Tijdschr Geneesk* 146:1100–1101
- van der Meer IM, Karamali NS, Boeke AJ, Lips P, Middelkoop BJ, Verhoeven I, Wuister JD (2006) High prevalence of vitamin D

- deficiency in pregnant non-Western women in The Hague, Netherlands. *Am J Clin Nutr* 84:350–353
27. Meulmeester JF, van den Berg H, Wedel M, Boshuis PG, Hulshof KF, Luyken R (1990) Vitamin D status, parathyroid hormone and sunlight in Turkish, Moroccan and Caucasian children in The Netherlands. *Eur J Clin Nutr* 44:461–470
 28. Brooke-Wavell K, Khan AS, Taylor R, Masud T (2008) Lower calcaneal bone mineral density and broadband ultrasonic attenuation, but not speed of sound, in South Asian than white European women. *Ann Hum Biol* 35:386–393
 29. Ward KA, Roy DK, Pye SR, O'Neill TW, Berry JL, Swarbrick CM, Silman AJ, Adams JE (2007) Forearm bone geometry and mineral content in UK women of European and South-Asian origin. *Bone* 41:117–121
 30. Hamson C, Goh L, Sheldon P, Samanta A (2003) Comparative study of bone mineral density, calcium, and vitamin D status in the Gujarati and white populations of Leicester. *Postgrad Med J* 79:279–283
 31. Solanki T, Hyatt RH, Kemm JR, Hughes EA, Cowan RA (1995) Are elderly Asians in Britain at a high risk of vitamin D deficiency and osteomalacia? *Age Ageing* 24:103–107
 32. Finch PJ, Ang L, Colston KW, Nisbet J, Maxwell JD (1992) Blunted seasonal variation in serum 25-hydroxy vitamin D and increased risk of osteomalacia in vegetarian London Asians. *Eur J Clin Nutr* 46:509–515
 33. Holvik K, Meyer HE, Haug E, Brunvand L (2005) Prevalence and predictors of vitamin D deficiency in five immigrant groups living in Oslo, Norway: the Oslo Immigrant Health Study. *Eur J Clin Nutr* 59:57–63
 34. Olmez D, Bober E, Buyukgebiz A, Cimrin D (2006) The frequency of vitamin D insufficiency in healthy female adolescents. *Acta Paediatr* 95:1266–1269
 35. Zargar AH, Ahmad S, Masoodi SR, Wani AI, Bashir MI, Laway BA, Shah ZA (2007) Vitamin D status in apparently healthy adults in Kashmir Valley of Indian subcontinent. *Postgrad Med J* 83:713–716
 36. Sahu M, Bhatia V, Aggarwal A, Rawat V, Saxena P, Pandey A, Das V (2009) Vitamin D deficiency in rural girls and pregnant women despite abundant sunshine in northern India. *Clin Endocrinol (Oxf)* 70:680–684
 37. Puri S, Marwaha RK, Agarwal N, Tandon N, Agarwal R, Grewal K, Reddy DH, Singh S (2008) Vitamin D status of apparently healthy schoolgirls from two different socioeconomic strata in Delhi: relation to nutrition and lifestyle. *Br J Nutr* 99:876–882
 38. Agarwal KS, Mughal MZ, Upadhyay P, Berry JL, Mawer EB, Puliyel JM (2002) The impact of atmospheric pollution on vitamin D status of infants and toddlers in Delhi, India. *Arch Dis Child* 87:111–113
 39. Madar AA, Stene LC, Meyer HE (2009) Vitamin D status among immigrant mothers from Pakistan, Turkey and Somalia and their infants attending child health clinics in Norway. *Br J Nutr* 101:1052–1058
 40. Lawson M, Thomas M (1999) Vitamin D concentrations in Asian children aged 2 years living in England: population survey. *BMJ* 318:28
 41. Goswami R, Kochupillai N, Gupta N, Goswami D, Singh N, Dudha A (2008) Presence of 25(OH) D deficiency in a rural North Indian village despite abundant sunshine. *J Assoc Physicians India* 56:755–757
 42. Marwaha RK, Tandon N, Reddy DR, Aggarwal R, Singh R, Sawhney RC, Saluja B, Ganie MA, Singh S (2005) Vitamin D and bone mineral density status of healthy schoolchildren in northern India. *Am J Clin Nutr* 82:477–482
 43. Vupputuri MR, Goswami R, Gupta N, Ray D, Tandon N, Kumar N (2006) Prevalence and functional significance of 25-hydroxyvitamin D deficiency and vitamin D receptor gene polymorphisms in Asian Indians. *Am J Clin Nutr* 83:1411–1419
 44. Gulvady C, Pingle S, Shanbhag S (2007) Incidence of vitamin B12 / D3 deficiency among company executives. *Indian J Occup Environ Med* 11:83–85
 45. Bhalala U, Desai M, Parekh P, Mokal R, Chheda B (2007) Subclinical hypovitaminosis D among exclusively breastfed young infants. *Indian Pediatr* 44:897–901
 46. Sachan A, Gupta R, Das V, Agarwal A, Awasthi PK, Bhatia V (2005) High prevalence of vitamin D deficiency among pregnant women and their newborns in northern India. *Am J Clin Nutr* 81:1060–1064
 47. Hollis BW (2005) Circulating 25-hydroxyvitamin D levels indicative of vitamin D sufficiency: implications for establishing a new effective dietary intake recommendation for vitamin D. *J Nutr* 135:317–322
 48. Thacher TD, Fischer PR, Strand MA, Pettifor JM (2006) Nutritional rickets around the world: causes and future directions. *Ann Trop Paediatr* 26:1–16
 49. Girish M, Subramanian G (2008) Rickets in exclusively breast fed babies. *Indian J Pediatr* 75:641–643
 50. el Hag AI, Karrar ZA (1995) Nutritional vitamin D deficiency rickets in Sudanese children. *Ann Trop Paediatr* 15:69–76
 51. Tezer H, Siklar Z, Dallar Y, Dogankoc S (2009) Early and severe presentation of vitamin D deficiency and nutritional rickets among hospitalized infants and the effective factors. *Turk J Pediatr* 51:110–115
 52. Echarri JJ, Bazebo JA, Guillem-Grima F (2008) Rachitic deformities of lower members in congolese children. *An Sist Sanit Navar* 31:235–240
 53. Prentice A (2008) Vitamin D deficiency: a global perspective. *Nutr Rev* 66:S153–S164
 54. Ozkan B, Doneray H, Karacan M, Vancelik S, Yildirim ZK, Ozkan A, Kosan C, Aydin K (2009) Prevalence of vitamin D deficiency rickets in the eastern part of Turkey. *Eur J Pediatr* 168:95–100
 55. Beck-Nielsen SS, Brock-Jacobsen B, Gram J, Brixen K, Jensen TK (2009) Incidence and prevalence of nutritional and hereditary rickets in southern Denmark. *Eur J Endocrinol* 160:491–497
 56. Mallet E, Gaudelus J, Reinert P, Le Luyer B, Lecointre C, Leger J, Loirat C, Quinet B, Benichou JJ, Furioli J, Loeuille GA, Roussel B, Larchet M, Freycon F, Vidailhet M, Varet I (2004) Symptomatic rickets in adolescents. *Arch Pediatr* 11: 871–878
 57. Yeste D, Carrascosa A (2003) Nutritional rickets in childhood: analysis of 62 cases. *Med Clin (Barc)* 121:23–27
 58. Jensen JE, Hitz MF (2000) Osteomalacia—a frequently overlooked condition among refugees and immigrants. *Ugeskr Laeger* 162:6250–6251
 59. Coster A, Ringe JD (2000) Osteomalacia in immigrants: therapeutic management of two cases. *Med Klin (Munich)* 95:451–456
 60. Balasubramanian K, Rajeswari J, Gulab GYC, Agarwal AK, Kumar A, Bhatia V (2003) Varying role of vitamin D deficiency in the etiology of rickets in young children vs. adolescents in northern India. *J Trop Pediatr* 49:201–206
 61. Pettifor JM (2004) Nutritional rickets: deficiency of vitamin D, calcium, or both? *Am J Clin Nutr* 80:1725S–1729S
 62. Lips P, Chapuy MC, Dawson-Hughes B, Pols HA, Holick MF (1999) An international comparison of serum 25-hydroxyvitamin D measurements. *Osteoporos Int* 9:394–397
 63. Datta S, Alfaham M, Davies DP, Dunstan F, Woodhead S, Evans J, Richards B (2002) Vitamin D deficiency in pregnant women from a non-European ethnic minority population—an interventional study. *Bjog* 109:905–908

64. Koch HC, Burmeister W (1993) Vitamin D status of children and adolescents of African and Asian diplomats in Germany. *Klin Padiatr* 205:416–420
65. Harinarayan CV (2005) Prevalence of vitamin D insufficiency in postmenopausal south Indian women. *Osteoporos Int* 16:397–402
66. Farrant HJ, Krishnaveni GV, Hill JC, Boucher BJ, Fisher DJ, Noonan K, Osmond C, Veena SR, Fall CH (2009) Vitamin D insufficiency is common in Indian mothers but is not associated with gestational diabetes or variation in newborn size. *Eur J Clin Nutr* 63:646–652
67. Khadilkar A, Das G, Sayyad M, Sanwalka N, Bhandari D, Khadilkar V, Mughal MZ (2007) Low calcium intake and hypovitaminosis D in adolescent girls. *Arch Dis Child* 92:1045
68. Sivakumar B, Vijayaraghavan K, Vazir S, Balakrishna N, Shatrugna V, Sarma KV, Nair KM, Raghuramulu N, Krishnaswamy K (2006) Effect of micronutrient supplement on health and nutritional status of schoolchildren: study design. *Nutrition* 22: S1–S7
69. Sivakumar B, Nair KM, Sreeramulu D, Suryanarayana P, Ravinder P, Shatrugna V, Kumar PA, Raghunath M, Rao VV, Balakrishna N, Kumar PU, Raghuramulu N (2006) Effect of micronutrient supplement on health and nutritional status of schoolchildren: biochemical status. *Nutrition* 22:S15–S25
70. Tiwari L, Puliye JM (2004) Vitamin D level in slum children of Delhi. *Indian Pediatr* 41:1076–1077