



Food and health

Higher fish consumption and lower risk of central nervous system demyelination

Lucinda J. Black¹ · Yun Zhao¹ · Yee Cheng Peng¹ · Jill L. Sherriff¹ · Robyn M. Lucas^{2,3} · Ingrid van der Mei⁴ · the Ausimmune Investigator GroupGavin Pereira¹

Received: 21 May 2019 / Revised: 25 June 2019 / Accepted: 5 July 2019
© The Author(s), under exclusive licence to Springer Nature Limited 2019

Abstract

Background/Objectives The evidence for diet as a risk factor for multiple sclerosis (MS) is inconclusive. We examined the associations between fish consumption and risk of a first clinical diagnosis of central nervous system demyelination (FCD), a common precursor to MS.

Methods The 2003–2006 Ausimmune Study was a case-control study examining environmental risk factors for FCD, with participants recruited from four regions of Australia and matched on age, sex, and study region. Dietary intake data were collected using a food frequency questionnaire. We used conditional logistic regression models to test associations between fish consumption (total, tinned, grilled, and fried) and risk of FCD (249 cases and 438 controls), adjusting for history of infectious mononucleosis, smoking, serum 25-hydroxyvitamin D concentrations, socio-economic status, omega-3 supplement use, dietary under-reporting, and total energy intake.

Results Higher total fish consumption (per 30 g/day, equivalent to two serves/week) was associated with an 18% reduced risk of FCD (AOR 0.82; 95% CI 0.70, 0.97). While we found no statistically significant associations between grilled and fried fish consumption and risk of FCD, higher tinned fish consumption (per 30 g/day) was associated with a 41% reduced risk of FCD (AOR 0.59; 95% CI 0.39, 0.89).

Conclusions Tinned fish is predominantly oily, whereas grilled and fried fish are likely to be a combination of oily and white types. Oily fish is high in vitamin D and very long chain polyunsaturated omega-3 fatty acids, both of which may be beneficial in relation to MS.

Introduction

Multiple sclerosis (MS) is a chronic inflammatory and neurodegenerative disease of the central nervous system (CNS)

characterised by demyelination and episodes of neurological disability or progressive neurologic deterioration [1]. The disease affects more than two million people globally, and is two to three times more common in females than males. A number of genetic and environmental risk factors have been described, the latter including past history of infectious mononucleosis, adolescent obesity, low past sun exposure, smoking, and low vitamin D status [1]. By the time of peak MS risk—early to middle adulthood—very few of these risk factors are modifiable [2]. There is, however, some evidence to indicate that diet is a potentially modifiable risk factor for MS [3–9].

A number of studies have shown that oily fish consumption associates with reduced risk of MS [8, 10, 11]. Oily fish is the best natural source of both dietary vitamin D [12] and very long-chain omega-3 polyunsaturated fatty acids (VLCn3PUFAs) [13], both of which may have a beneficial role in MS. The association between vitamin D status and risk of MS is well established [2], while VLCn3PUFAs—namely eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA),

Deceased: Tony McMichael

✉ Lucinda J. Black
lucinda.black@curtin.edu.au

- ¹ School of Public Health, Curtin University, Bentley, WA 6102, Australia
- ² National Centre for Epidemiology and Population Health, The Australian National University, Canberra, ACT 1200, Australia
- ³ Centre for Ophthalmology and Visual Science, University of Western Australia, Perth, WA 6009, Australia
- ⁴ Menzies Institute for Medical Research, University of Tasmania, Hobart, TAS, Australia

and docosapentaenoic acid (DPA)—play critical roles in the CNS, exhibiting anti-inflammatory, and neuroprotective effects [14].

The 2003–2006 Australian Multi-centre Study of Environment and Immune Function (the Ausimmune Study) [15] is a multicentre, incident case-control study investigating the environmental risk factors for a first clinical diagnosis of CNS demyelination (FCD), a common precursor to MS. The Ausimmune Study is one of the largest, most well-characterised samples of people with early MS worldwide. Previously, higher vitamin D status [16], higher VLCn3PUFA intakes (largely from fish, but also including small amounts from commonly consumed meats, such as beef and ham) [17], a higher healthy dietary pattern score [9], and higher unprocessed red meat consumption [18] have been associated with reduced risk of FCD in the Ausimmune Study. We build on these studies by examining the associations between fish consumption and risk of FCD using data from the Ausimmune Study.

Methods

Study population

Between November 2003 and December 2006, participants were recruited from four regions of Australia: Brisbane city, Newcastle region, Geelong and the Western districts of Victoria, and the island of Tasmania [15]. Case participants ($n = 282$; 18–59 years) were referred to the study as previously described [15]. Following a full history and neurological examination, a study neurologist confirmed the date of onset and presenting symptoms suggestive of CNS demyelination [15]. Within the study period, case participants were diagnosed with CNS demyelination for the first time. The diagnoses included: a classic first demyelinating event (FDE; defined as a single, first, episode of clinical symptoms suggestive of CNS demyelination; $n = 216$); a first recognised event, but past history revealed a prior, undiagnosed event, that, on review was highly suggestive of CNS demyelination ($n = 48$); first presentation of primary progressive MS (based on neurological assessment on study entry ($n = 18$)). It is unlikely that a prior unrecognised demyelinating event that had not been ascribed to CNS demyelination would have triggered any changes in dietary behaviour; thus these participants are considered to have an ‘incident’ FCD. The date of MRI scan, which was available for most participants, was used as a proxy for the date of FCD. There was a time lag between the date of MRI scan by the neurologist (the date of the diagnosis which brought the participants into the study) and the study interview. For participants with dietary intake data, the median (interquartile range (IQR)) time lag was 101 (147) days.

The Australian Electoral Roll (compulsory registration for citizens ≥ 18 years) was used to randomly select control participants ($n = 558$) from the general population. Control participants were matched to case participants on age (within 2 years), sex, and study region. Between one and four controls were matched to each case to maximise power, with more controls per case in regions with a lower expected number of cases due to being either at lower latitude (and lower expected incidence) or a smaller source population. Ethics approval was obtained from the nine Human Research Ethics Committees of the participating institutions [15]. All participants gave written informed consent for the use of their data. All participant information was anonymised and de-identified prior to analysis.

Dietary assessment

Information on habitual dietary intakes in the 12 months prior to the study interview was collected using the Cancer Council Victoria Dietary Questionnaire for Epidemiological Studies version 2 (DQESv2) [19]. The DQES v2 is a self-administered, semi-quantitative, food frequency questionnaire (FFQ) developed for use in the ethnically diverse adult Australian population [20]. The consumption of food items from four food groups (cereals, sweets, and snacks; dairy, meats, and fish; fruit; vegetables) was recorded on a scale from “never” to “three or more times per day”. Portion size diagrams of four commonly consumed foods (potato, other vegetables, steak, and casserole) were used to determine respondents’ average portion size factor, and responses were used to scale standard portion sizes up or down for different foods [19]. Fish consumption (g/day) was reported for total (sum of tinned, grilled, and fried fish), tinned, grilled, and fried fish.

Covariates

Self-reported questionnaires were used to collect information on history of infectious mononucleosis (yes, no, and do not know), highest level of education (year 10 or less, year 12 and Technical and Further Education, university), and total number of years smoked minus any periods of absence. Socio-economic status was assessed as quintiles of the Index of Relative Socio-economic Advantage and Disadvantage (IRSAD) using postal area code and data from the 2006 Census of Population and Housing: Socio-Economic Indexes for Areas (SEIFA), Australia [21]. IRSAD summarises information about the economic and social conditions of households within an area, including relative advantage and disadvantage measures: a low score indicates relatively greater disadvantage; a high score indicates greater advantage in general [21]. Supplement use was captured, as follows: “In the last 12 months, have you

used any dietary or vitamin supplements on a regular basis?" If yes, participants recorded the name, type, dose, and frequency of use. Those who were using any fish oil, omega-3, or cod liver oil supplements were considered to be using an omega-3 supplement.

A study nurse measured height and weight. Basal metabolic rate (BMR) was calculated using the equations developed by Harris and Benedict [22]. Under-reporters were classified using a Goldberg cut-off below $BMR \times 1.05$ [23] and a two-category variable was defined: under-reporter vs. plausible reporter. Most participants (94%) provided a blood sample: serum aliquots (1 mL) were stored at -80°C . Serum samples were analysed at study completion for 25-hydroxyvitamin D (25(OH)D) concentrations using liquid chromatography tandem mass spectrometry [16]. Using the seasonal patterns, serum 25(OH)D concentrations for control participants were adjusted to match the date of the case blood draw [16]. This was done to account for blood samples of cases and controls being taken at different times of the year.

Statistical analysis

Characteristics of participants were described as frequency and percentage for categorical variables, mean, and standard deviation (SD) for continuous variables with a Normal distribution, median, and IQR for continuous variables with a non-Normal distribution.

We used conditional logistic regression models (with cases and controls matched on age, sex, and study region) to estimate odds ratios (ORs), 95% confidence intervals (95% CI), and *p* values for associations between fish consumption (total, tinned, grilled, and fried) and risk of FCD. Fish consumption was analysed as a continuous variable per 30 g/day, to reflect a clinically relevant level of consumption (equivalent to ~2 serves/week). Model 1 was unadjusted; model 2 was adjusted for known environmental risk factors for MS (history of infectious mononucleosis, serum 25(OH)D concentrations, and total years of smoking), along with socio-economic status, omega-3 supplement use, dietary under-reporting, and total energy intake.

We tested for non-linearity using quadratic terms for total, tinned, grilled, and fried fish consumption. To test for any sex differences, we included an interaction term for fish consumption and sex in the adjusted models. We conducted the following sensitivity analyses: (a) within the smaller group of those with a classic FDE; and (b) including only participants with plausible energy intakes (500–5000 kcal/day, based on a daily energy intake associated with survival [23], which excludes those with extreme energy intakes as previously described [24, 25]).

The Paramed command in Stata [26] was used to assess the potential mediating effects (natural indirect effect, NIE)

of serum 25(OH)D concentrations on the relationship between fish consumption and risk of FCD, adjusting for the same environmental risk factors. Statistically, the NIE compares the risks of FCD when the values of the serum 25(OH)D concentrations vary from the one realised at fish consumption ≥ 30 g/day to the one realised at fish consumption < 30 g/day, assuming all relevant participants with fish consumption ≥ 30 g/day. The mediation analyses were performed for all participants, and further for the subgroup of people who had plausible energy intakes. Data were analysed using Stata 14 software [27].

Results

A total of 791 participants (272 cases and 519 controls) provided dietary intake data. Missing data for covariates were as follows: total years of smoking $n = 3$; serum 25(OH)D concentrations, $n = 38$; dietary misreporting, $n = 3$; socio-economic status, $n = 10$. A total of 687 participants (249 cases and 438 controls) provided complete data on dietary intake and all covariates, and were part of at least a matched control pair. As expected, case participants were more likely than controls to have a history of infectious mononucleosis, lower serum 25(OH)D concentrations, and a greater total number of years of smoking (Table 1). Median total, tinned, and grilled fish consumption were lower in case than control participants, while fried fish consumption was higher. Total fish consumption ≥ 30 g/day was reported by 40% of participants (10, 12, and 3% of participants consumed ≥ 30 g/day tinned, grilled, and fried fish, respectively).

In adjusted models, higher total fish consumption (per 30 g/day, equivalent to two serves/week) was associated with an 18% reduced risk of FCD, and higher tinned fish consumption (per 30 g/day) was associated with a 41% reduced risk of FCD (Table 2). There were no statistically significant associations between grilled or fried fish consumption and risk of FCD (Table 2). The effect estimates for all models were similar when limited to those with plausible energy intakes and in the smaller subgroup of those with a classic FDE (Table 2). There was no evidence of non-linearity in any models and no evidence of any interactions between fish consumption and sex. We found no evidence of statistically significant mediating effects of the serum 25(OH)D concentrations (total fish intake: NIE = 1.00, 95% CI: 0.94–1.05, and $p = 0.846$; tinned fish intake: NIE = 0.99, 95% CI: 0.89–1.09, and $p = 0.808$). The results were similar for the subgroup of people who had plausible energy intakes (total fish intake: NIE = 1.00, 95% CI: 0.95–1.05, and $p = 0.941$; tinned fish intake: NIE = 1.00, 95% CI: 0.92–1.08, and $p = 0.918$).

Table 1 Characteristics of Ausimmune participants included in the current study (249 cases and 438 controls)

	Case	Control
Sex, <i>n</i> (%) ^a		
Male	63 (25.3)	108 (24.7)
Female	186 (74.7)	330 (75.3)
Age (years), mean (SD) ^a	38.7 (9.7)	40.0 (9.6)
Study region, <i>n</i> (%) ^a		
Brisbane (27°S)	83 (33.3)	159 (36.3)
Newcastle (33°S)	32 (12.9)	65 (14.8)
Geelong (37°S)	59 (23.7)	108 (24.7)
Tasmania (43°S)	75 (30.1)	106 (24.2)
History of infectious mononucleosis, <i>n</i> (%)		
No	163 (65.5)	345 (78.8)
Yes	70 (28.1)	71 (16.2)
Do not know	16 (6.4)	22 (5.0)
Serum 25(OH)D concentrations (nmol/L), mean (SD)	75.6 (29.5)	82.1 (30.4)
Total years of smoking, median (IQR)	5.7 (18.6)	2 (14.8)
Socio-economic status, <i>n</i> (%)		
Quintile 1 (lowest)	28 (11.2)	57 (13.0)
Quintile 2	38 (15.3)	49 (11.2)
Quintile 3	43 (17.3)	103 (23.5)
Quintile 4	83 (33.3)	136 (31.1)
Quintile 5 (highest)	57 (22.9)	93 (21.2)
Energy intake (kcal/day), median (IQR)	1658.4 (858.1)	1743.9 (894.4)
Dietary misreporting, <i>n</i> (%)		
Under-reporter	106 (42.6)	176 (40.2)
Plausible reporter	143 (57.4)	262 (59.8)
Fish consumption (g/day), median (IQR)		
Total fish	21.7 (28.1)	23.7 (30.0)
Tinned fish	4.7 (10.2)	6.6 (13.0)
Grilled fish	8.5 (14.2)	10.6 (16.1)
Fried fish	3.5 (8.6)	2.9 (8.2)

^aCase and control participants were matched for sex, age (within 2 years), and study region

SD standard deviation, 25(OH)D 25-hydroxyvitamin D, IQR interquartile range

Discussion

Our results demonstrate an association between higher fish consumption and lower risk of FCD, particularly for tinned fish. An increment of two serves/week of tinned fish associated with ~40% reduced risk of FCD. Two serves of fish per week is in line with the Australian Dietary Guidelines (one serve = 100 g cooked fish fillet (115 g raw) or one small can of fish) [28]. Tinned fish is primarily oily fish (e.g. tuna, salmon, sardines, and mackerel), which is the

richest dietary source of both vitamin D [12] and VLCn3PUFAs [13]. We found no association between consumption of grilled or fried fish and risk of FCD. Grilled fish is likely to include a combination of oily and white fish. The latter has lower levels of both vitamin D and VLCn3PUFAs than oily fish. Fried fish, particularly from take-away outlets, is likely to be white fish, again with lower levels of both vitamin D and VLCn3PUFAs than oily fish.

Low vitamin D status is a known risk factor for MS [16]. The major source of vitamin D for humans is cutaneous synthesis from sun exposure; dietary intake of vitamin D becomes important when sun exposure is limited, with oily fish considered one of the best natural sources. However, in this study, we have not been able to demonstrate that serum 25(OH)D concentration is a significant mediator on the relationship between the fish consumption and the risk of FCD. This contradicts a previous hypothesis that intake of oily fish may compensate for vitamin D deficiency that is associated with increased MS risk [8].

VLCn3PUFAs have been shown to suppress pro-inflammatory T-helper cells [29]; to inhibit the migration of T-helper cells across the blood brain barrier [30]; and to inhibit matrix metalloproteinases which are toxic to myelin [31]. DHA is a major constituent of neuronal membranes [32] and appears to play a role in synaptic signal transduction [33], while DPA is emerging as an important bioactive fatty acid, with a role in brain function and mental health [34]. Previous analysis of data from the Ausimmune Study showed a reduced risk of FCD with increasing intake of VLCn3PUFAs [17]. Furthermore, a small number of clinical trials have investigated the hypothesis that a higher intake of omega-3 fatty acids reduces disease activity in those with clinically diagnosed MS; however, the evidence is inconclusive [35–37]. We were not able to test the mediating effects of VLCn3PUFAs due to the high uptake of fish oil supplements after FCD and the strong possibility of reverse causation in measuring VLCn3PUFAs in blood after diagnosis.

Fish is also an important source of the sulfur-containing amino acid, taurine [38–40], which has therapeutic potential against neurological disorders [41]. Taurine has been shown to have anti-inflammatory and neuroprotective properties in a mouse Parkinson's disease model [42], and has been identified as a dysregulated metabolite in animal models of MS, including in non-human primates [43]. Using a global metabolomics approach, taurine has also been shown to enhance remyelination through increasing differentiation of oligodendrocyte precursor cells. On the basis of these findings, the authors suggested that taurine supplementation, in combination with existing treatment strategies, could be a feasible strategy to improve remyelination [44]. To our knowledge, the role of taurine in MS risk has not been investigated.

Table 2 Conditional logistic regression models showing associations between total, tinned, grilled, and fried fish consumption and (a) risk of FCD (249 cases and 438 controls); (b) risk of true FDE (191 cases and 328 controls); (c) risk of FCD in those with plausible energy intakes (500–5000 kcal/day) (245 cases and 428 controls)

	Model 1: unadjusted		Model 2: adjusted ^a	
	OR (95% CI)	<i>P</i>	AOR (95% CI)	<i>P</i>
(a) Risk of FCD				
Total fish, per 30 g/day	0.88 (0.76, 1.02)	0.088	0.82 (0.70, 0.97)	0.024
Tinned fish, per 30 g/day	0.64 (0.44, 0.94)	0.022	0.59 (0.39, 0.89)	0.012
Grilled fish, per 30 g/day	0.89 (0.71, 1.10)	0.284	0.83 (0.65, 1.07)	0.143
Fried fish, per 30 g/day	0.84 (0.55, 1.28)	0.421	0.79 (0.49, 1.26)	0.321
(b) Risk of FDE				
Total fish, per 30 g/day	0.92 (0.79, 1.08)	0.295	0.89 (0.73, 1.07)	0.208
Tinned fish, per 30 g/day	0.71 (0.47, 1.09)	0.115	0.67 (0.42, 1.07)	0.096
Grilled fish, per 30 g/day	0.92 (0.74, 1.14)	0.444	0.89 (0.68, 1.16)	0.389
Fried fish, per 30 g/day	0.96 (0.60, 1.53)	0.87	0.96 (0.55, 1.69)	0.898
(c) Risk of FCD in those with plausible energy intakes (500–5000 kcal/day)				
Total fish, per 30 g/day	0.87 (0.74, 1.02)	0.094	0.84 (0.70, 1.00)	0.054
Tinned fish, per 30 g/day	0.66 (0.45, 0.97)	0.033	0.61 (0.40, 0.93)	0.023
Grilled fish, per 30 g/day	0.90 (0.69, 1.18)	0.461	0.88 (0.65, 1.18)	0.385
Fried fish, per 30 g/day	0.82 (0.51, 1.32)	0.415	0.76 (0.46, 1.28)	0.305

^aAdjusted for history of infectious mononucleosis, serum 25-hydroxyvitamin D concentrations, total years of smoking, socio-economic status, omega-3 supplement use, dietary under-reporting and total energy intake

Our results concur with the few other studies investigating fish consumption and risk of MS. In a 2005–2012 population-based case-control study (1879 cases and 4135 controls) of incident MS in Sweden, participants were asked to specify how often, on average, they had eaten oily or white fish during the last 5 years [8]. Consumption of oily, but not white, fish was associated with decreased occurrence of MS. A case-control study conducted in Norway in 2003 (152 cases and 402 controls) investigated sun exposure and dietary vitamin D in people with MS residing at latitudes above the Arctic Circle [10]. The results supported a protective effect of consuming boiled or fried fish (unspecified type) three or more times a week. In a case-control study of incident MS (197 cases, 202 age- and sex-matched controls) conducted between 1992 and 1995 in Montreal, Canada, dietary intake data were collected using a 164-item FFQ [11]. Fish consumption was protective in women only, with borderline statistical significance; however, the type of fish consumed was unspecified and likely to reflect consumption of both oily and white fish.

A major strength of the Ausimmune Study is its strong study design (multicentre, matched case-control), along with the recruitment of participants with an incident FCD (rather than participants with established MS). Although not all participants had a classic FDE during the study period, we performed sensitivity analyses in the smaller group of those with a classic FDE. Results were similar to the main models, albeit with wider CI. We used an established FFQ to collect information on dietary intake over the previous 12 months, and we were able to account for a number of

important potential confounders, including serum 25(OH)D concentrations measured by an accurate and reliable methodology [45].

Although the widely acknowledged under-reporting of energy intake [46] is a limitation of our study, we attempted to account for dietary under-reporting by adjusting for a misreporting variable. Further, we performed a sensitivity analysis including only participants with plausible energy intakes (500–5000 kcal/day), with minimal change in effect estimates compared with the main model. Although case participants may be more likely to recall exposure to risk factors than control participants [47], this bias is likely to be minimal in our study because diet is not commonly considered a cause of CNS demyelination. The association between higher fish consumption and reduced risk of FCD in our study may be due to lifestyle or environmental characteristics not captured in our analyses, and we cannot rule out the possibility of residual confounding. However, we adjusted for the main known environmental risk factors for MS. Other lifestyle characteristics, including current body mass index and physical activity, were not associated with risk of FCD in previous analysis of the Ausimmune Study [48]. Since the study consisted of Australian participants who were predominantly Caucasian, the findings may not be generalisable to other populations with differing diets.

Our results suggest a protective effect of higher fish consumption, particularly tinned (oily) fish consumption, on risk of FCD. The equivalent of two serves per week of tinned fish was associated with ~40% reduced risk of FCD.

This level of fish consumption is in line with the Australian Dietary Guidelines. Given the higher levels of VLCn3PU-FAs in oily fish compared with white fish, future studies would benefit from separating the consumption of these two types of fish.

Acknowledgements We thank the participants of the Ausimmune Study. We would like to acknowledge and thank the physicians who notified case participants to the Ausimmune Study: Jeffrey Blackie FRACP, Richard Bourke FRACGP, John Cameron MD, Ross Carne MD, Ben Clark FRANZCO, Steven Collins MD, Diana Conrad FRANZCO, Michael Coroneos FRACS, Nicholas Downie FRANZCO, David Floate FRACP, Peter Gates FRACP, Kerry Green FRACP, Erwin Groeneveld FRANZCO, John Harrison FRANZCO, Michael Haybittel FRANZCO, Robert Henderson FRACP, John Henshaw MMed, James Hurley MD, Dean Jones FRACP, Michael Katekar MBBS, Anthony Kemp FRACP, Mark King FRACP, George Kiroff FRACS, Brett Knight FRACP, Thomas Kraemer FRACP, Cecile Lander FRACP, Jeannette Lechner-Scott FRACP, Andre Loisselle FRACP, Paul McCartney FRANZCO, Pamela McCombe PhD, Mark McGree FRANZCO, David McKnight FRANZCO, Daniel McLaughlin PhD, Satish Nagarajah MBBS, Rob Nightingale FRACP, Terence O'Brien MD, John O'Sullivan MD, Gregory Outteridge FRANZCO, Anthony Pane FRANZCO, Mark Parsons FRACP, Melinda Pascoe FRACP, David Prentice PhD, Richard Ralph FRACGP, Stephen Read FRACP, John Richmond FRACP, Ian Routley FRANZCO, Timothy Ruddle FRANZCO, Noel Saines FRACP, Stan Siejka MBBS (dec), Christopher Staples FRACP, Paul Talman FRACP, Don Todman FRACP, Nitin Verma FRANZCO, Brendan Vote FRANZCO, Michael Waldie FRANZCO, Michael Weetch FRACP, Rodney Westmore FRANZCO, Andrew Wong FRACP; and the local research officers: Susan Agland BN, Barbara Alexander BN, Marcia Davis MD, Zoe Dunlop BN, Rosalie Scott BN, Marie Steele RN, Catherine Turner MPH&TM, Brenda Wood RN; and the Ausimmune Study project officers during the course of the study: Jane Gresham MA(Int Law), Camilla Jozwick BSc(Hons), Helen Rodgers RN.

Funding Funding for the Ausimmune Study was provided by the National Multiple Sclerosis Society of the United States of America, the National Health and Medical Research Council of Australia and Multiple Sclerosis Research Australia. LJB is supported by a Multiple Sclerosis Western Australia Postdoctoral Fellowship and a Curtin University Research Fellowship. RML is supported by a National Health and Medical Research Council of Australia Senior Research Fellowship. Funding bodies had no role in the design or conduct of the study; collection, management, analysis, or interpretation of data; or preparation, review, or approval of the paper.

⁵Barwon Health, Geelong, VIC, Australia; ⁶Royal Brisbane and Women's Hospital and The University of Queensland, Brisbane, QLD, Australia; ⁷School of Public Health, University of Adelaide, Adelaide, SA, Australia; ⁸Murdoch Childrens Research Institute, University of Melbourne, Melbourne, VIC, Australia; ⁹Centre for Neuroscience, University of Melbourne, Melbourne, VIC, Australia; ¹⁰QIMR Berghofer Medical Research Institute, Brisbane, QLD, Australia; ¹¹Hunter Health, Newcastle, NSW, Australia

Author contributions The Ausimmune Investigator Group and LJB designed the study; LJB analysed and interpreted the data and wrote the paper; YCP contributed to data analysis; GP provided statistical support and contributed to data interpretation; JLS, RML, GP, and the Ausimmune Investigator Group provided critical revision of the paper for important intellectual content; LJB had primary responsibility for

the final content. All the authors read and approved the final version of the paper.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. Reich DS, Lucchinetti CF, Calabresi PA. Multiple sclerosis. *N Engl J Med*. 2018;378:169–80.
2. O'Gorman C, Lucas R, Taylor B. Environmental risk factors for multiple sclerosis: a review with a focus on molecular mechanisms. *Int J Mol Sci*. 2012;13:11718–52.
3. Lauer K. Notes on the epidemiology of multiple sclerosis, with special reference to dietary habits. *Int J Mol Sci*. 2014;15:3533–45.
4. Mandia D, Ferraro O, Nosari G, Montomoli C, Zardini E, Bergamaschi R. Environmental factors and multiple sclerosis severity: a descriptive study. *Int J Environ Res Public Health*. 2014;11:6417.
5. Riccio P, Rossano R. Nutrition facts in multiple sclerosis. *ASN Neuro*. 2015;7. <https://doi.org/10.1177/1759091414568185>.
6. Jahromi SR, Toghae M, Jahromi MJ, Aloosh M. Dietary pattern and risk of multiple sclerosis. *Iran J Neurol*. 2012;11:47–53.
7. Sedaghat F, Jessri M, Behrooz M, Mirghotbi M, Rashidkhani B. Mediterranean diet adherence and risk of multiple sclerosis: a case-control study. *Asia Pac J Clin Nutr*. 2016;25:377–84.
8. Baarnhielm M, Olsson T, Alfredsson L. Fatty fish intake is associated with decreased occurrence of multiple sclerosis. *Mult Scler J*. 2014;20:726–32.
9. Black LJ, Rowley C, Sherriff J, Pereira G, Ponsonby A-L, Ausimmune Investigator Group et al. A healthy dietary pattern associates with a lower risk of a first clinical diagnosis of central nervous system demyelination. *Mult Scler J*. 2018. <https://doi.org/10.1177/1352458518793524>.
10. Kampman MT, Wilsgaard T, Mellgren SI. Outdoor activities and diet in childhood and adolescence relate to MS risk above the Arctic Circle. *J Neurol*. 2007;254:471–7.
11. Ghadirian P, Jain M, Ducic S, Shatenstein B, Morisset R. Nutritional factors in the aetiology of multiple sclerosis: a case-control study in Montreal, Canada. *Int J Epidemiol*. 1998;27:845–52.
12. Liu J, Arcot J, Cunningham J, Greenfield H, Hsu J, Padula D, et al. New data for vitamin D in Australian foods of animal origin: impact on estimates of national adult vitamin D intakes in 1995 and 2011–13. *Asia Pac J Clin Nutr*. 2015;24:464–71.
13. Meyer BJ. Australians are not meeting the recommended intakes for omega-3 long chain polyunsaturated fatty acids: results of an analysis from the 2011-2012 National Nutrition and Physical Activity Survey. *Nutrients*. 2016;8:111.
14. DeFilippis AP, Sperling LS. Understanding omega-3s. *Am Heart J*. 2006;151:564–70.
15. Lucas R, Ponsonby AL, McMichael A, van der Mei I, Chapman C, Coulthard A, et al. Observational analytic studies in multiple sclerosis: controlling bias through study design and conduct. The Australian Multicentre Study of Environment and Immune Function. *Mult Scler J*. 2007;13:827–39.
16. Lucas RM, Ponsonby A-L, Dear K, Valery PC, Pender MP, Taylor BV, et al. Sun exposure and vitamin D are independent risk factors for CNS demyelination. *Neurology*. 2011;76:540–8.

17. Hoare S, Lithander F, van der Mei I, Ponsonby AL, Lucas R. Higher intake of omega-3 polyunsaturated fatty acids is associated with a decreased risk of a first clinical diagnosis of central nervous system demyelination: results from the Ausimmune Study. *Mult Scler J*. 2016;22:884–92.
18. Black LJ, Bowe GS, Pereira G, Lucas RM, Dear K, van der Mei I, et al. Higher non-processed red meat consumption is associated with a reduced risk of central nervous system demyelination. *Front Neurol*. 2019;10:125.
19. Cancer Council Victoria. Dietary questionnaire for epidemiological studies (DQES v2) user information guide. Victoria, Australia: Carlton; 2009.
20. Ireland P, Jolley D, Giles G, O’Dea K, Powles J, Rutishauser I, et al. Development of the Melbourne FFQ: a food frequency questionnaire for use in an Australian prospective study involving an ethnically diverse cohort. *Asia Pac J Clin Nutr*. 1994;3:19–31.
21. Australian Bureau of Statistics. Census of Population and Housing, 2006. <http://www.abs.gov.au>. Accessed 24 Jan 2019.
22. Harris J, Benedict F. A biometric study of basal metabolism in man. Washington D.C.: Carnegie Institute of Washington; 1919.
23. Goldberg GR, Black AE, Jebb SA, Cole TJ, Murgatroyd PR, Coward WA, et al. Critical evaluation of energy intake data using fundamental principles of energy physiology: derivation of cut-off limits to identify under-reporting. *Eur J Clin Nutr*. 1991;45:569–81.
24. Willett W. Nutritional epidemiology. 3rd ed New York: Oxford University Press; 2013.
25. Ambrosini GL, Fritschi L, de Klerk NH, Mackerras D, Leavy J. Dietary patterns identified using factor analysis and prostate cancer risk: a case control study in Western Australia. *Ann Epidemiol*. 2008;18:364–70.
26. Emsley R, Liu H. Stata module to perform causal mediation analysis using parametric regression models, 2013. <https://ideas.repec.org/c/boc/bocode/s457581.html>. Accessed 16 Feb 2019.
27. StataCorp. Stata Statistical Software: Release 14. College Station, TX: StataCorp LP; 2015.
28. Australian Government Department of Health. Lean Meat and poultry, fish, eggs, tofu, nuts and seeds and legumes/beans, 2017. <https://www.eatforhealth.gov.au/food-essentials/five-food-groups/lean-meat-and-poultry-fish-eggs-tofu-nuts-and-seeds-and>. Accessed 24 Jan 2019.
29. Allen MJ, Fan YY, Monk JM, Hou TY, Barhoumi R, McMurray DN, et al. n-3 PUFAs reduce T-helper 17 cell differentiation by decreasing responsiveness to interleukin-6 in isolated mouse splenic CD4(+) T cells. *J Nutr*. 2014;144:1306–13.
30. Shinto L, Marracci G, Bumgarner L, Yadav V. The effects of omega-3 fatty acids on matrix metalloproteinase-9 production and cell migration in human immune cells: implications for multiple sclerosis. *Autoimmune Dis*. 2011;2011:134592.
31. Liuzzi GM, Latronico T, Rossano R, Viggiani S, Fasano A, Riccio P. Inhibitory effect of polyunsaturated fatty acids on MMP-9 release from microglial cells-implications for complementary multiple sclerosis treatment. *Neurochem Res*. 2007;32:2184–93.
32. Yehuda S, Rabinovitz S, Mostofsky DI. Essential fatty acids are mediators of brain biochemistry and cognitive functions. *J Neurosci Res*. 1999;56:565–70.
33. Jones CR, Arai T, Rapoport SI. Evidence for the involvement of docosahexaenoic acid in cholinergic stimulated signal transduction at the synapse. *Neurochem Res*. 1997;22:663–70.
34. Kaur G, Guo XF, Sinclair AJ. Short update on docosapentaenoic acid: a bioactive long-chain n-3 fatty acid. *Curr Opin Clin Nutr Metab Care*. 2016;19:88–91.
35. Ramirez-Ramirez V, Macias-Islas MA, Ortiz GG, Pacheco-Moises F, Torres-Sanchez ED, Sorto-Gomez TE, et al. Efficacy of fish oil on serum of TNF alpha, IL-1 beta, and IL-6 oxidative stress markers in multiple sclerosis treated with interferon beta-1b. *Oxid Med Cell Longev*. 2013;2013:709493.
36. Bates D, Cartledge NE, French JM, Jackson MJ, Nightingale S, Shaw DA, et al. A double-blind controlled trial of long chain n-3 polyunsaturated fatty acids in the treatment of multiple sclerosis. *J Neurol Neurosurg Psychiatry*. 1989;52:18–22.
37. Torkildsen O, Wergeland S, Bakke S, Beiske AG, Bjerve KS, Hovdal H, et al. omega-3 fatty acid treatment in multiple sclerosis (OFAMS Study): a randomized, double-blind, placebo-controlled trial. *Arch Neurol*. 2012;69:1044–51.
38. Gormley T, Neumann T, Fagan J, Brunton N. Taurine content of raw and processed fish fillets/portions. *Eur Food Res Technol*. 2007;225:837–42.
39. Tørris C, Småstuen MC, Molin M. Nutrients in fish and possible associations with cardiovascular disease risk factors in metabolic syndrome. *Nutrients*. 2018;10:E952.
40. Wójcik OP, Koenig KL, Zeleniuch-Jacquotte A, Costa M, Chen Y. The potential protective effects of taurine on coronary heart disease. *Atherosclerosis*. 2010;208:19–25.
41. Jakaria M, Azam S, Haque ME, Jo S-H, Uddin MS, Kim I-S, et al. Taurine and its analogs in neurological disorders: focus on therapeutic potential and molecular mechanisms. *Redox Biol*. 2019;24:101223.
42. Che Y, Hou L, Sun F, Zhang C, Liu X, Piao F, et al. Taurine protects dopaminergic neurons in a mouse Parkinson’s disease model through inhibition of microglial M1 polarization. *Cell Death Dis*. 2018;9:435–435.
43. ‘t Harst BA, Vogels JTWE, Spijksma G, Brok HPM, Polman C, van Der Greef J. 1H-NMR spectroscopy combined with pattern recognition analysis reveals characteristic chemical patterns in urines of MS patients and non-human primates with MS-like disease. *J Neurol Sci*. 2003;212:21–30.
44. Beyer BA, Fang M, Sadrian B, Montenegro-Burke JR, Plaisted WC, Kok BPC, et al. Metabolomics-based discovery of a metabolite that enhances oligodendrocyte maturation. *Nat Chem Biol*. 2017;14:22–31.
45. Black LJ, Anderson D, Clarke MW, Ponsonby A-L, Lucas RM, Ausimmune Investigator Group. Analytical bias in the measurement of serum 25-hydroxyvitamin D concentrations impairs assessment of vitamin D status in clinical and research settings. *PLoS ONE*. 2015;10:e0135478.
46. Black AE, Prentice AM, Goldberg GR, Jebb SA, Bingham SA, Livingstone MBE, et al. Measurements of total energy expenditure provide insights into the validity of dietary measurements of energy intake. *J Am Diet Assoc*. 1993;93:572–9.
47. Althubaiti A. Information bias in health research: definition, pitfalls, and adjustment methods. *J Multidiscip Healthc*. 2016;9:211–7.
48. Ponsonby AL, Lucas RM, Dear K, van der Mei I, Taylor B, Chapman C, et al. The physical anthropometry, lifestyle habits and blood pressure of people presenting with a first clinical demyelinating event compared to controls: the Ausimmune study. *Mult Scler J*. 2013;19:1717–25.