ABSTRACT: The innate immune system utilizes many approaches for defense against invading microorganisms, including complement-mediated lysis, engulfment, formation of neutrophil extracellular traps (NETs) and release of antimicrobial peptides (AMPs).1

Recent evidence demonstrates that macrophages produce the AMP LL-37 in response to endogenously produced 1,25(OH)2D to enhance innate immunity. Additional evidence shows 1,25(OH)2D modulates the adaptive immune system as well through direct effects on T cell activation and on the phenotype and function of antigen-presenting cells (APCs), particularly of dendritic cells (DCs).2

This paper hypothesizes that well-tolerated Vitamin D3 supplementation merits investigation in combating coronaviruses and their diseases, such as COVID-19.

DISCUSSION: The immune system defends the body from foreign, invading organisms, promoting protective immunity while maintaining tolerance to self. The implications of vitamin D deficiency on the immune system have become clearer in recent years and in the context of vitamin D deficiency, there appears to be an increased susceptibility to infection and a diathesis, in a genetically susceptible host to autoimmunity.3

There are two major classes of amphipathic AMPs present in human respiratory lining fluids: defensins and cathelicidins. There is evidence that both of these classes of AMPs play a role during Influenza A Virus (IAV) infection. One representative of the class of cathelicidins is LL-37. Recent reviews have discussed the extraordinary range of activities of LL-37, which include direct antimicrobial and antiviral activities, chemotactic activities for various immune cells, modulation of macrophage responses to inflammatory stimuli, and modulation of dendritic cell responses.4

Deaths caused by IAV infection mostly resulted from acute lung injury, systemic inflammation or bacterial superinfection, suggesting that new treatments with anti-viral, anti-bacterial and anti-inflammation effects would be ideal. AMPs are antimicrobial peptides that not only play important roles as host defense against pathogens but also modulate inflammatory responses, and thus they are potential candidates for IAV

Previous studies have reported that 1,25(OH)2D3, the hormonal form of vitamin D, is a negative endocrine regulator of the RAS and inhibits renin biosynthesis.. The results of the present study demonstrated that vitamin D inhibited renin, ACE and Ang II expression, and induced ACE2 levels in LPS-induced ALI. Therefore, vitamin D may attenuate LPS-Induced ALI by, at least partially, inducing ACE2/Ang-(1-7) axis activity and inhibiting renin and the ACE/Ang II/AT1R cascade.5

Angiotensin-converting enzyme 2 (ACE2) is the cellular receptor for severe acute respiratory syndromecoronavirus (SARS-CoV) and the new coronavirus (SARS-CoV-2) that is causing the serious coronavirus disease 2019 (COVID-19) epidemic.6

Observational studies report consistent independent associations between low serum concentrations of 25hydroxyvitamin D (the major circulating vitamin D metabolite) and susceptibility to acute respiratory tract infection. 25-hydroxyvitamin D supports induction of antimicrobial peptides in response to both viral and bacterial stimuli, suggesting a potential mechanism by which vitamin D inducible protection against respiratory pathogens might be mediated.7

The recent discovery that vitamin D induces antimicrobial peptide gene expression explains, in part, the 'antibiotic' effect of vitamin D and has greatly renewed interest in the ability of vitamin D to improve immune function. Subsequent work indicates that this regulation is biologically important for the response of the innate immune system to wounds and infection and that deficiency may lead to suboptimal responses toward bacterial and viral infections.8

To assess the overall effect of vitamin D supplementation on risk of acute respiratory tract infection, and to identify factors modifying this effect. 25 eligible randomised controlled trials (total 11 321 participants, aged 0 to 95 years) were identified. Vitamin D supplementation was safe and it protected against acute respiratory tract infection overall. Patients who were very vitamin D deficient and those not receiving bolus doses experienced the most benefit.9

CONCLUSION: Whereas, in consideration, it is reasonable to investigate the role of Vitamin D deficiency in COVID-19 (SARS-CoV-2) disease (via ACE2 receptor entry) and reducing the innate immune system's effective defense. Further, exogenous Vitamin D supplementations may reasonably lessen susceptibility to the disease progression and complications, via AMP promotion and immune sytem modulation, specifically related to inflammatory response mechanisms.

- and Autoimmune Disease J. Michelle Kahlenberg and Mariana J. Kaplan J Immunol November 15, 2013.
- 2 Vitamin D and molecular actions on the immune system: modulation of innate and autoimmunity. DLKamen, V.Tangpricha
- 3 Vitamin D and the Immune System Cynthia Aranow, MD, Investigator
- Little Peptide, Big Effects: The Role of LL-37 in Inflammation 4 The Role of Antimicrobial Peptides in Influenza Virus Infection and Their Potential as Antiviral and Immunomodulatory Therapy, IHsieh, KHartshorn.
 - 5 Vitamin D alleviates lipopolysaccharide-induced acute lung injury via regulation of the renin-angiotensin system. Jun Xu Jialai Ya et al
 - 6Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. Renhong Yan et al.
- 7Vitamin D supplementation to prevent acute respiratory tract infections; systematic review and meta-analysis of individual participant data. Adrian R Martineau et al.
- 8The vitamin D-antimicrobial peptide pathway and its role in protection against infection Adrian F Gombart, Future Microbio. 2009.
- 9 Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. Adrian R Martineau et al. 2016.

Why Vitamin D3 is the KEY to fighting Coronavirus like COVID-19!

The innate immune system is the best protector when supported. **D**3 "...vitamin D induces antimicrobial peptide gene expression."2

- kills virus
- kills bacteria
- blocks ARDS (Immune over reaction)

Cytokine Storm

CATHELICIDIN C205H340N60O53

Vitamir

"AMPs (anti-microbial peptides Cathelicidin and Defensin) are endogenous proteins playing important roles in host defense through direct antimicrobial and antiviral activities and through immunomodulatory effects."3


Secondary **Bacteria**

> ³ Calcitriol-modulated human antibiotics: New pathophysiological aspects of vitamin D Carlos Antonio Amado Diagoa, et al. 2016.

Vitamin D deficiency contributes directly to the acute respiratory distress syndrome (ARDS) Rachel C A Dancer, Dhruv Parekh, [...], and David R Thickett, Thorax, 2015.

COVID-19 Coronavirus

> ² The vitamin D-antimicrobial peptide pathway and its role in protection against infection Adrian F Gombart, Future Microbio, 2009.



The main role of ACE2 (angiotensin-converting enzyme) is the degradation of Ang II resulting in the formation of angiotensin 1–7 (Ang 1–7) which opposes the actions of Ang II. Increased Ang II levels are thought to upregulate ACE2 activity, thus, ACE2 plays a crucial role in the RAS because it opposes the actions of Ang II. Consequently, it has a beneficial role in many diseases such as hypertension, diabetes, and cardiovascular disease.³


¹ https://www.drugtargetreview.com/news/ 56895/scientists-demonstrate-how-covid-19

^{2 1,25-}Dihydroxyvitamin D3 Suppresses Renin Gene Transcription by Blocking the Activity of the Cyclic AMP Response Element in the Renin Gene Promoter Welhua Yuan ‡, Wei Pan ‡, Juan Kong ‡, Wei Zheng ‡, Frances L, Szato §, Kan E. Wong §, Ronald Cohen ‡, Anna Klopot ‡, Zhongyi Zhang ‡ and Yan Chun Li ‡

³ Angiotensin-Converting Enzyme 2 (ACE2) Is a Key Modulator of the Renin Angiotensin System in Health and Disease Chris Tikellis1 and M. C. Thomas1

Mechanism of Vitamin D3 and ARDS* — Page 4 of 6-

(*Acute Respiratory Distress Syndrome)



Mechanism through which vitamin D modulates the secretion of cathelicidin and beta-2-defensin. Binding of some microbial antigens (Ag) to toll-like receptors (TLRs) activates 25-hydroxyvitamin D-1α-hydroxylase (CYP27b1), so that, depending on intracellular concentrations of 25-hydroxyvitamin D, local synthesis of 1,25-dihydroxyvitamin D is enhanced. The latter in turn binds to its receptor (VDR), and the intranuclear hormone-receptor complex activates the transcription of the cathelicidin and beta-2-defensin genes. Intracellular concentrations of 25-hydroxyvitamin D depend on plasma concentrations of this molecule, which is the most reliable marker of the nutritional status of this vitamin.3

Edema

Fluid

Vitamin D deficiency contributes directly to the acute respiratory distress syndrome (ARDS) Rachel C A Dancer, Dhruv Parekh, [...], and David R Thickett, Thorax, 2015.

² The vitamin D-antimicrobial peptide pathway and its role in protection against infection Adrian F Gombart, Future Microbio, 2009.

³ Calcitriol-modulated human antibiotics: New pathophysiological aspects of vitamin D Carlos Antonio Amado Diagoa, et al. 2016.

