

Journal Pre-proof

Treatment of SARS-CoV-2 with high dose oral zinc salts: A report on four patients

Eric Finzi



PII: S1201-9712(20)30441-0

DOI: <https://doi.org/10.1016/j.ijid.2020.06.006>

Reference: IJID 4311

To appear in: *International Journal of Infectious Diseases*

Received Date: 19 May 2020

Revised Date: 2 June 2020

Accepted Date: 3 June 2020

Please cite this article as: Finzi E, Treatment of SARS-CoV-2 with high dose oral zinc salts: A report on four patients, *International Journal of Infectious Diseases* (2020), doi: <https://doi.org/10.1016/j.ijid.2020.06.006>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Treatment of SARS-CoV-2 with high dose oral zinc salts: A report on four patients

Eric Finzi, MD, PhD¹

George Washington School of Medicine¹, 2120 L St, Washington, DC, 20037

efinzi@mfa.gwu.edu

240 353 0611

Summary line: Treatment of COVID-19 with Zinc Salts

Abstract: 46 words

Text: 1194 words

Highlights

Treatment of COVID-19 with high dose zinc salts

Improvement of oxygenation in COVID-19 after zinc gluconate

Rapid resolution of COVID-19 shortness of breath after zinc salts

Abstract

Coronavirus 2019 (COVID-19) is a pandemic with substantial mortality and no accepted therapy. We report here on four consecutive outpatients with clinical characteristics (CDC case definition) of and/or laboratory-confirmed COVID-19

who were treated with high dose zinc salt oral lozenges. All four patients experienced significant improvement in objective and symptomatic disease measures after one day of high dose therapy suggesting that zinc therapy was playing a role in clinical recovery. A mechanism for zinc's effects is proposed based on previously published studies on SARS-CoV-1, and randomized controlled trials assessing zinc shortening of common cold duration. The limited sample size and study design preclude a definitive statement about the effectiveness of zinc as a treatment for COVID-19 but suggest the variables to be addressed to confirm these initial findings in future trials.

Keywords: COVID-19, Zinc ,Treatment, Case Series

COVID-19 is a pandemic with substantial mortality and no accepted therapy. Ideally, effective therapy would be both readily available and minimally toxic. Coronaviruses are positive sense RNA viruses of which SARS-Coronavirus-2 (SARS-CoV-2) is the most recently identified member and cause of COVID-19. Previous work has shown that increased intracellular Zn^{2+} concentrations can interfere with proteolytic processing of polyproteins in many RNA viruses¹. In addition, all coronaviruses are unified in requiring a RNA-dependent RNA polymerase (RdRp) that is a core enzyme in their RNA-synthesizing machinery. Increased Zn^{2+} levels directly inhibit isolated RdRp complexes and purified recombinant RdRps, SARS-CoV-1 virus replication in tissue culture, and replication of other positive sense RNA viruses².

It is unknown but plausible that Zn^{2+} could work by the mechanisms above to inhibit SARS-CoV-2 virus replication. Of note, coronaviruses are part of the larger subset of RNA viruses that cause the common cold. A recent meta-analysis of 7 randomized controlled trials (RCT's) showed that zinc lozenges shortened the mean duration of the common cold by 33% (95% CI, 21-45%)³. Zinc has also been proposed to be a helpful cofactor in treatment of COVID-19 with hydroxychloroquine⁴⁻⁷.

We asked whether zinc salt lozenges could be therapeutic for COVID-19.

Four consecutive outpatients with clinical characteristics (CDC case definition) of and/or laboratory-confirmed COVID-19 were treated by the author with high dose zinc salt oral lozenges, and clinical symptoms followed on an outpatient basis. Patients only received treatments as specified in each case. All patients provided verbal informed consent and were treated on a compassionate basis. The reported efficiency of zinc lozenges for the common cold ranked zinc acetate>zinc gluconate>zinc citrate⁸. Patients 1,2 were treated with zinc citrate lozenges (23mg of elemental zinc); patient 3, zinc citrate/zinc gluconate(23mg); patient 4, zinc acetate (15 mg). Patients were started on zinc therapy at different times in their disease course,

depending on when they were referred for treatment. They were instructed to take lozenges every 2-4 hours, dissolving lozenges on their tongue over 20 to 30 minutes, but not to exceed 200mg. Patient 2 did not take the higher dose until day 10 of her illness. No side effects from zinc therapy were reported, other than a chalky taste (case 2). Elemental zinc doses up to 216mg/day have been well tolerated in clinical trials for the common cold⁹, without any significant side effects. Short term zinc use at these doses is considered very safe, with occasional gastric distress and metallic taste being the only reported side effects¹⁰.

Case 1

The subject was a 63-year-old man who developed symptoms 5 days after contact with a household member (A) who had direct exposure to a laboratory-confirmed COVID-19 case. At 5 pm on his first day of illness he developed intensely painful muscle aches in the mid- and lower back, headache, shaking chills and a fever to 100.5°F/38.1°C. He assumed he had COVID-19, so he took three 23mg zinc citrate lozenges. Over the next 24 hours, he took a total of 9 lozenges (207mg). His improvement in fever, headaches, and muscle pain began on day 2. On day 3 he became afebrile. His symptoms continued to improve over the next 10 days while taking 184 mg of Zn²⁺ per day.

Case 2

A 57-year-old female partner of case 1. She developed mild symptoms of diarrhea, fatigue, and low grade fever (under 100.5), 7 days after contact with household member A. She took only one or two 23mg zinc lozenges daily. On day 5 she developed a dry, nonproductive cough, sore throat, chest tightness and with mild dyspnea while walking. On day 10, she suddenly developed severe dry cough, chest pain, intense neck muscle pain, fever, headache, and shortness of breath at rest. Over the next 5 hours, she took a 23 mg lozenge every hour. After a total of 7 zinc lozenges (161mg), her cough and shortness of breath was much improved. On day 11, she had reduced her zinc dose to two daily and her cough started again. After 3 more lozenges her coughing decreased. She increased her regimen to taking 5 lozenges (115mg) daily for 10 days, during which time she gradually improved.

Case 3

A 41-year-old female healthcare worker developed body aches, cough and sinus pain on day 1. On day 3 she had a fever to 101.5, severe body aches, worsening cough and tested positive for COVID-19. On day 4 her PaO₂ dropped to 94% on room air, (Figure 1,2). On day 6 she developed shortness of breath (PaO₂- 93/94%) and was started on hydroxychloroquine, 200mg po bid. On days 7-9 she experienced an overall worsening in her cough, body aches and shortness of breath.

On the evening of day 9, she started zinc, 23mg, q 4 hours, 138 mg daily. On day 10 she felt better; her PaO₂ and fever began improving one day after zinc was initiated (Figure 1,2). She continued zinc q 4 hours and was well by day 19.

Case 4

A 26 year-old female acquired COVID-19 through direct contact with an infected health care worker. During week 1 she had fever, cough and severe body aches. In the second week she developed shortness of breath and severe fatigue. During week 3 her fever abated but she had a significant cough, fatigue and body aches, and slept more than 14 hours daily. After three weeks of symptoms the patient began 15 mg of zinc, q 2 hours, 10 lozenges a day, 150mg daily for 14 days. One day later the cough and body aches and fatigue began to improve, and two weeks later she felt fully recovered.

We report four consecutive outpatients with clinical characteristics of and /or laboratory-confirmed COVID-19 who were treated with high dose zinc salt lozenges. Two cases lacked laboratory confirmation but had direct exposure to confirmed cases and had characteristic clinical features (CDC case definition). To the best of our knowledge, this is the first report of COVID-19 cases treated with high dose zinc salts. In all 4 cases symptomatic and objective measures of disease began to improve within 24 hours after initiation of zinc lozenges.

Twenty percent of common colds are caused by coronaviruses. Zinc acetate and zinc gluconate appear to be the most effective in RCT's to treat common colds^{8,9}, and they also have the most theoretical effectiveness as they release the most free Zn^{2+} . The effectiveness also appears to be influenced by the time course of dissolution of the lozenge on the tongue, and exact chemical composition of the lozenge^{8,9}. In successful zinc clinical trials, total ionic zinc release from lozenges was found to be highly correlated ($P < 0.001$) with reduction in common cold symptoms⁹.

In this uncontrolled case series, initiation of high dose zinc salt lozenges was followed by symptomatic and objective improvement in 4 consecutive patients with clinical signs and symptoms and/or laboratory confirmation of COVID-19. However, the limited sample size and study design preclude a definitive statement about the potential effectiveness of this treatment. The day within the disease course that each patient initiated zinc therapy varied, and due to difficulty in obtaining zinc lozenges during the pandemic, three different zinc salts were used. Therefore, controlled clinical trials with single formulation will be required to properly assess the role of zinc therapy in CoVID-19. Particular care is needed in the selection of zinc lozenges such that sufficient ionic zinc is released. In addition, frequent dosing (i.e- q 2-4hours), adequate dosing (i.e. close to 200mg daily), and slow dissolution on the tongue (20 -30 min) will be essential for meaningful trials^{8,9}.

Given the low toxicity and ease of administration of zinc lozenges it would seem prudent to begin testing of zinc in a controlled setting as a potential therapy for COVID-19.

Figure 1 Legend.

Case 3 self-monitored her PaO₂ by pulse oximetry. The arrow shows the start of high dose zinc lozenges.

Journal Pre-proof

Fig 1





Figure 2 Legend.

Case 3 self-monitored daily oral temperature. The arrow shows the start of high dose zinc lozenges.



The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The work described has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans.

Funding: No funding was provided for this study.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- 1) Lanke K, Krenn BM, Melchers WJ, Seipelt J, van Kuppeveld FJ. PDTC inhibits picornavirus polyprotein processing and RNA replication by transporting zinc ions into cells. *J Gen Virol*. 2007 Apr;88(Pt 4):1206-17.
- 2) te Velthuis AJW, van den Worm SHE, Sims AC, Baric RS, Snijder EJ, van Hemert MJ. Zn(2+) inhibits coronavirus and arterivirus RNA polymerase activity in vitro and zinc ionophores block the replication of these viruses in cell culture. *PLOS Pathog*. 2010; 6:e1001176.
- 3) Hemilä H. Zinc lozenges and the common cold: a meta-analysis comparing zinc acetate and zinc gluconate, and the role of zinc dosage. *JRSM Open*. 2017 May 2;8(5):2054270417694291. doi: 10.1177/2054270417694291.
- 4) Skalny AV, Rink L, Ajsuvakova OP, et al. Zinc and respiratory tract infections: Perspectives for COVID-19 (Review). *Int J Mol Med*. 2020 Apr 14.
- 5) Shittu MO, Afolami OI. Improving the efficacy of Chloroquine and Hydroxychloroquine against SARS-CoV-2 may require Zinc additives - A better synergy for future COVID-19 clinical trials. *Infez Med*. 2020 Ahead of print Jun 1;28(2):192-197.
- 6) Derwand R, Scholz M. Does zinc supplementation enhance the clinical efficacy of chloroquine/hydroxychloroquine to win today's battle against COVID-19? *Med Hypotheses*. 2020 May 6;142:109815.
- 7) Carlucci P, Tania T, Petrilli et al. Hydroxychloroquine and azithromycin plus zinc vs hydroxychloroquine and azithromycin alone: outcomes in hospitalized COVID-19 patients *Medrxiv* doi: <https://doi.org/10.1101/2020.05.02.20080036>
- 8) Eby GA. Zinc lozenges: cold cure or candy? Solution chemistry determinations. *Biosci Rep*. 2004 Feb;24(1):23-39.
- 9) Eby GA. Zinc lozenges as cure for the common cold – A review and hypothesis. *Med Hypothesis*. 2010 Mar; 74(3): 482–492.
- 10) Prasad AS, Fitzgerald JT, Bao B, et al. Duration of symptoms and plasma cytokine levels in patients with the common cold treated with zinc acetate. A randomized, double-blind, placebo-controlled trial. *Ann Int Med* 2000;133:245–52.