

# **A Basic Review of the Preliminary Evidence that COVID-19 Risk and Severity is Increased in Vitamin D Deficiency**

**Linda L Benskin**<sup>1,2\*</sup>

<sup>1</sup> Independent researcher for improving health in rural areas of tropical developing countries, Austin, TX, USA

<sup>2</sup> Ferris Mfg. Corp., makers of PolyMem<sup>®</sup> multifunctional dressings, Ft. Worth, TX, USA

\* **Correspondence:** LindaBenskin@utexas.edu

**Keywords: vitamin D; COVID-19; health disparities; minority health; vitamin D deficiency; preventive medicine; inflammation; cytokine storm;**

## **Abstract**

As the world's attention has been riveted upon the growing COVID-19 pandemic, many researchers have written brief reports supporting the hypothesis that vitamin D deficiency is related to the incidence and severity of COVID-19.

The clear common thread among the top risk groups - vitamin D deficiency – may be being overlooked because of previous overstated claims of vitamin D benefits. However, the need to decrease COVID-19 fatalities among high-risk populations is urgent.

Early researchers reported three striking patterns. Firstly, the innate immune system is impaired by vitamin D deficiency, which would predispose sufferers to viral infections such as COVID-19. Vitamin D deficiency also increases the activity of the X-chromosome-linked 'Renin-Angiotensin' System, making vitamin D deficient individuals (especially men) more susceptible to COVID-19's deadly "cytokine storm" (dramatic immune system overreaction). Secondly, the groups who are at highest risk for severe COVID-19 match those who are at highest risk for severe vitamin D deficiency. This includes the elderly, men, ethnic groups whose skin is naturally rich in melanin (if living outside the tropics), those who avoid sun exposure for cultural and health reasons, those who live in institutions, the obese, and/or those who suffer with hypertension, cardiovascular disease, or diabetes. And thirdly, the pattern of geographical spread of COVID-19 reflects higher population vitamin D deficiency. Both within the USA and throughout the world, COVID-19 fatality rates parallel vitamin D deficiency rates.


A literature search was performed on PubMed, Google Scholar, and RSMLDS, with targeted Google searches providing additional sources. Although randomized controlled trial results may be available eventually, the correlational and causal study evidence supporting a link between vitamin D deficiency and COVID-19 risks is already so strong that it supports action.

The 141 author groups writing primarily about biological plausibility detailed how vitamin D deficiency can explain every risk factor and every complication of COVID-19, but agreed that other factors are undoubtedly at work. COVID-19 was compared with dengue fever, for which oral vitamin D supplements of 4000IU for 10 days were significantly more effective than 1000IU in reducing virus replication and controlling the "cytokine storm" responsible for fatalities.

Among the 47 original research studies summarized here, chart reviews found that serum vitamin D levels predicted COVID-19 mortality rates (16 studies) and linearly predicted COVID-19 illness severity (8 studies). Two causal modeling studies and several analyses of variance strongly supported the hypothesis that vitamin D deficiency is a causal, rather than a bystander factor, in COVID-19 outcomes. Three of the four studies whose findings opposed the hypothesis relied upon disproven assumptions.

The literature review also found that prophylactically correcting possible vitamin D deficiency during the COVID-19 pandemic is extremely safe. Widely recommending 2000IU of vitamin D daily for all populations with limited ability to manufacture vitamin D from the sun has virtually no potential for harm and is reasonably likely to save many lives.

**Search and Reporting Method:**



(Note: Because this is such a rapidly evolving topic, some of the references for this article were still in the preprint stage of publication when this review was finalized. These citations are marked with the superscript “p” and the journal to which they have been submitted is listed in the references, where available. In addition, three studies (refs 46, 47, and 205) from Southeast Asia, all with results consistent with that of other studies, may have relied upon data from unofficial sources.)

## Contents

1	Introduction .....	4
1.1	The Vitamin D Debate .....	4
1.1.1	Irish Medical Journal debate on vitamin D supplements during the COVID-19 pandemic ..	5
1.1.2	Debate over reports using “big data” (the UK Biobank and EPIC in the USA, see Results, 3.4.6)	5
1.2	Defining appropriate serum vitamin D levels and appropriate supplementation dosages.....	7
1.2.1	Controversy concerning risk of overdose .....	7
1.2.2	Controversy over appropriate 25(OH)D goals .....	8
1.2.3	Controversy over recommended supplement doses.....	9
2	Methods: Literature Search .....	10
3	Results of Searches .....	12
3.1	Planned Clinical Trials .....	12
3.2	Biological Plausibility Discussions .....	12
3.2.1	Vitamin D enhances resistance to viral illnesses .....	12
3.2.2	How vitamin D may decrease serious COVID-19-associated complications.....	13
3.3	Risk for Severe COVID-19 Parallels Risk for Vitamin D Deficiency .....	14
3.4	Evidence Informing the Hypotheses that Vitamin D Deficiency Influences COVID-19.....	15
3.4.1	Analyses of population data .....	15
3.4.2	Causal inference modeling reports.....	20
3.4.3	Case Studies and Case Series in which vitamin D is mentioned .....	21
3.4.4	Prospective correlational study, case-controlled survey, and cohort observational study	22
3.4.5	Retrospective chart reviews favoring the hypothesis.....	22
3.4.6	Retrospective chart reviews that are neutral or strongly oppose the hypothesis .....	27
3.4.7	Rapid systematic review and meta-analysis with an ecological approach .....	28
4	COVID-19-Specific Recommendations of Experts.....	28
5	Discussion.....	29
6	Conclusion.....	30
7	Limitations.....	31
8	Afterward .....	31
8.1	Conflict of Interest .....	31
8.2	Author Contributions .....	31
8.3	Funding .....	31
8.4	Acknowledgments.....	31
8.5	Author Brief Biography .....	31
8.6	Abbreviations:.....	32
9	References .....	32
10	Appendices.....	57
10.1	Table 2.....	57
10.2	Table 3.....	65
10.3	The Irish Medical Journal Debate .....	75

# 1 Introduction

COVID-19 was first recognized in December of 2019.(1,2) By January of 2020 it was clear the elderly are by far the most likely succumb to COVID-19 pneumonia, which is caused by a “cytokine storm.”(3,4) Later, male sex, obesity, and possessing naturally melanin-rich skin while living outside of the tropics came to be known as the highest risk factors after older age.(2<sup>p</sup>,5-10,11<sup>p</sup>,12,13) Unlike influenza, children under age ten are almost completely spared in COVID-19.(14,15) This unusual risk factor pattern presented a mystery that spawned studies showing that COVID-19 fatalities are especially high in areas with lower levels of sunshine due to latitude or air pollution, except when population vitamin D intake is high.(6,7,9,16–21) In fact, the risk groups for severe COVID-19 match the risk groups for vitamin D deficiency exactly, and there is biological plausibility: vitamin D is known to modulate the immune system, helping prevent both an under-reaction that allows upper respiratory infections to be contracted, and the over-reaction referred to in COVID-19 as the “cytokine storm” (see **Section 3.2**) (16,22,23) This review explores the evidence related to the hypothesis that vitamin D deficiency increases both COVID-19 rates and illness severity.

## 1.1 The Vitamin D Debate

The discussion of the idea that the top risk groups for severe COVID-19 complications tend to have vitamin D deficiency [**Table 1**] was initially popularized, not by the scientific community or governmental bodies, but rather, by some entertainers and bloggers, who recommended supplements to their audiences.(24–26) This led some in the scientific community to respond with either agreement or disapproval. Trinity College Dublin researchers quickly issued a news release urging the Irish government to change their recommendations for vitamin D supplements in light of evidence of an association between vitamin D levels and COVID-19 mortality.(27) However, most governments, medical organizations, and key opinion leaders give one or more of these four reasons not to recommend vitamin D supplements: past claims for vitamin D benefits were overstated, evidence for a link to COVID-19 is insufficient, overdoses are theoretically possible, and the public might believe that taking vitamin D supplements will make them “immune” to COVID-19.(28–36)

**Table 1 : Classification of vitamin D levels (serum 25(OH)D levels):(9,37,38<sup>p</sup>,39–51)**

Classification	Nanograms	Nanomoles	Recommended D
danger of toxicity	> 100 ng/ml*	> 250 nmol/l	
normal or optimal	> 30 ng/ml	> 75 nmol/l	400 - 4000 IU/day
insufficient	21 - 29 ng/ml	51 - 74 nmol/l	4000 - 6000 IU/day
deficient	11 - 20 ng/ml	26 - 50 nmol/l	7000 IU/day
severely deficient (often not distinguished from deficient)	< 10ng/ml	25 nmol/l	10,000 IU/day x 1 month or 500,000IU x 1

\*some sources found that 150ng/ml was not harmful

Although the International Association for Gerontology and Geriatrics (IAGG) Asia/Oceania Region COVID-19 Prevention Statement acknowledged that increasing vitamin D levels could

reduce infection risks in elderly individuals whose levels are insufficient they recommended only “getting enough sunlight in the morning” without mentioning supplements.(52) Two May 2020 Centre for Evidence-Based Medicine rapid reviews concluded, without discussing any of the recent studies, that there is no evidence to support a role for vitamin D in prevention or treatment of COVID-19 or the cytokine storm.(29,53) Alarmed by the media response to a literature review suggesting a link between COVID-19 and vitamin D, two Brazilian medical associations jointly published a note stating that vitamin D supplements are only approved for bone health.(54–56) The high mortality rates among minorities are providing momentum for various public health program expansions, which could diminish if vitamin D deficiency, rather than access to care and economic disparities, were found to be even a partial explanation.(57–60) In addition, previous studies of dubious quality suggesting that vitamin D can “cure” various chronic illnesses and may be influencing the reluctance to recommend supplements for COVID-19.(61<sup>P</sup>)

However, despite these concerns, former CDC Chief Dr. Tom Frieden recommended sunshine and up to 2000IU/day of vitamin D as a potential preventative for COVID-19, the British Dietetic Association recommended sunshine (or 400IU/day for those are not able to go outside due to self-isolation), and former vitamin D skeptic Dr. JoAnn Manson’s calls for daily vitamin D supplements (1000 - 2000IU/day) during the COVID-19 pandemic - if vitamin D intake is low and going outdoors is not feasible - were published on both Medscape and WebMD.(16,62–65) Medscape published a second review of the topic by McCall, in which correcting possible vitamin D deficiency was characterized as “low hanging fruit” that has no downside.(66) Mitchell’s brief review (20 May 2020) in a Lancet-affiliated online journal also supported the vitamin D hypothesis.(67) Authors of an early meta-analysis of nine studies found that a high percentage of COVID-19 patients are either vitamin D insufficient or deficient, and that countries with lower population vitamin D status have somewhat higher COVID-19 mortality rates and somewhat lower COVID-19 recovery rates.(68<sup>P</sup>) In Qatar, vitamin D for prevention of COVID-19 is being proactively delivered to the homes of high-risk diabetics.(69)

### **1.1.1 Irish Medical Journal debate on vitamin D supplements during the COVID-19 pandemic**

The Irish Medical Journal hosted a six-article formal debate on the topic in response to three published reports, including one by the researchers managing Ireland’s part of the 26-country longitudinal study on aging (TILDA), in their May 2020 issue.(70–73) All three reports strongly recommended vitamin D supplements to help protect all adults in Ireland from COVID-19 while they are “cocooning” (not going outdoors).(70,71) [details in Appendix 10.3]

### **1.1.2 Debate over reports using “big data” (the UK Biobank and EPIC in the USA, see Results, 3.4.6)**

Three research teams relied on the 2006-2010 UK Biobank data for the vitamin D levels included in their analyses of the relationship between COVID-19 incidence and vitamin D status.(10,74<sup>P</sup>,75) Roy, et al., challenged the assertion that vitamin D levels are stable over time (important since levels were assessed 10-14 years prior to the pandemic) noting that the cited study only included women and followed up for only 5 years.(10,76,77) In fact, in cited study (Meng, et al.) found that, rather than being stable, the mean 25(OH)D increased significantly

( $p < 0.05$ ) over the five years, and that the increase was driven by significant ( $p < 0.05$ ) increases among participants who were initially at risk for deficiency; supplement intake and overall vitamin D intake increased significantly ( $p < 0.05$ ).<sup>(76)</sup> Etsy criticized a UK Biobank study's assumption that, despite government's recommendations to supplement, the participants failed to correct any vitamin D deficiency revealed by their participation.<sup>(78)</sup> In their preprint, Darling, et al., cited a different study to support their assertion that vitamin D levels are stable over time.<sup>(10,74<sup>P</sup>,75)</sup> However, the Norwegians in the cited study had far higher vitamin D levels than the UK Biobank participants at their initial evaluation, a subset increased their 25(OH)D levels significantly ( $0 < 0.001$ ) by initiating supplements, and, as in the study by Meng, et al., vitamin D levels did increase significantly for the group as a whole over time ( $p < 0.01$ ).<sup>(79)</sup> The authors of the most recent article using the UK BioBank study data did not address the issue of the use of potentially no longer accurate 25(OH)D levels in their preprint, but added a reference to the same Norwegian study as Darling, et al., in their published article.<sup>(75)</sup>

During the time frame of these three studies, COVID-19 testing in the UK was extremely limited (Raisi-Estrabragh, et al., stated that most were tested only if hospitalized).<sup>(75,77)</sup> Such limited testing, Roy asserted, raises the possibility that the authors of the first UK Biobank study accidentally included COVID-19 positive patients who were only moderately ill in their negative group.<sup>(77)</sup> It is also likely that the two UK Biobank studies authored later compared COVID-19 patients with patients who had serious illnesses such as influenza pneumonia, rather than with healthy individuals.<sup>(74<sup>P</sup>,75)</sup> If vitamin D deficiency increases viral infection risk and severity as hypothesized, the patients in both arms of these studies could be expected to have higher deficiency and insufficiency rates than the general population. In fact, the research teams found high rates of vitamin D insufficiency and deficiency in both of their study groups.<sup>(74<sup>P</sup>,75)</sup>

Three authors all pointed out that the UK Biobank studies failed to address the severity of the COVID-19 the patients experienced, which is critical to the question of whether or not vitamin D deficiency contributes to the potentially fatal cytokine storm.<sup>(10,74<sup>P</sup>,75,77,78,80)</sup> Boucher cautioned against adjusting data for obesity or dark-skinned ethnicity, providing empirical evidence that both directly lower 25(OH)D.<sup>(81)</sup>

Grant and McDonnell formally responded to the first UK Biobank study, asserting that their multivariable model was over-adjusted because causal factors were treated as confounds, suggesting that the authors provide multiple analyses for transparency, including simple and complex models.<sup>(82)</sup> They also requested that the analysis be stratified by ethnicity, citing a previous study in which low 25(OH)D increased risk for preterm birth equally across ethnicities.<sup>(82)</sup> They echoed Roy's concern that lack of a positive COVID-19 test result did not assure lack of infection in the UK at the time.<sup>(82)</sup> Grant and McDonnell concluded by pointing out that few UK Biobank participants had 25(OH)D levels in the immune-protective range ( $>40\text{ng/ml}$ ), which would decrease the effect.<sup>(82)</sup>

In their response, rather than addressing the question of vitamin D deficiency being caused by old age, disability, obesity, etc., the authors of the first UK Biobank study stated that 25(OH)D cannot be a mediator because it is not the "cause" of old age, disability, etc.<sup>(82)</sup> They asserted that impaired health is more likely associated with reduced outdoor activity than with vitamin D

status.(82) They presented the non-significant results of an “intermediate” model that still included the deficiency-related variables of age, sex, ethnicity, and obesity, omitting only “health-related covariates” (e.g., BP, diabetes) as proof that inclusion of potential mediators did not influence their initial study results.(82) The original study authors also reiterated their assertions that there is no statistical interaction between ethnicity and vitamin D deficiency and a positive COVID-19 test would ascertain more severe infections, and concluded by stating that 40ng/ml is not deficient because in the adult UK population mean 25(OH)D levels are only 17.4ng/ml for men under 65 years and 18.9ng/ml for women under 65 years.(82)


Although vitamin D levels are not drawn routinely, Fox used data from EPIC, a database with 15,000,000 patients across 26 US states, in his analysis of the relationship between vitamin D status and COVID-19 infection, hospitalization, and mortality rates.(83) DeFilipps commented succinctly, pointing out because vitamin D deficiency is ubiquitous, assuming patients with no 25(OH)D in their charts were vitamin D sufficient renders the study results unreliable.(84) DeFilipps recommended evaluating only the subgroup who were hospitalized for COVID-19 who had pre-existing conditions and known vitamin D deficiency to determine if there was a relationship between their level of deficiency, illness severity, complications, or length of stay.(84)

## **1.2 Defining appropriate serum vitamin D levels and appropriate supplementation dosages**

The 25(OH)D (serum vitamin D level) is the most reliable indicator of functional vitamin D status, but until recently, the test assays varied.(85) However, past research study results can be compared by mathematically harmonizing them, and increasingly, labs are adopting LC/MS (D<sub>2</sub>+D<sub>3</sub>) as the standard, increasing consistency.(85)

### **1.2.1 Controversy concerning risk of overdose**

Food fortification was introduced shortly after the discovery of vitamin D. However, there was a dramatic increase in infants with hypercalcemia in the UK, leading to an abrupt scaling back of fortification.(86) Later, a rare genetic defect, Williams–Beuren syndrome, was found to be responsible for the hypercalcemia.(86) However, vitamin D toxicity concerns remain heightened, with a reluctance to recommend supplements (**Figure 1**).



Dietary sources provide UK adults with only about 100IUs of vitamin D per day.(87,88) During the COVID-19 pandemic, after concluding that vitamin D is likely to reduce acute respiratory tract infection risk, and that 10,000IU/day is safe, the NHS paradoxically recommended only 400IU/day “to protect bone and muscle health.”(34,88)

It is not considered possible to achieve toxic levels via the sun alone, and supplementation for prolonged periods brings 25(OH)D to toxic levels only if the dose is consistently extraordinarily high (40,000IU/day for many months).(38<sup>p</sup>,86,89,90) The average naturally acquired 25(OH)D among equatorial tribal groups is 46ng/ml.(91) Healthy lifeguards typically have 25(OH)D levels of 100-125ng/ml.(39)

The Endocrine Society found toxicity symptoms only at levels above 150ng/ml.(91) Toxicity is related to high calcium levels; 25(OH)D levels higher than 150ng/ml in conjunction with high calcium levels produce weakness, GI symptoms and accompanying weight loss, arrhythmias, confusion, and kidney damage.(38,86,90) Historically, toxic levels of vitamin D (>150ng/ml) have almost exclusively been the result of industrial errors (inaccurate doses in supplements or fortified foods), and the few cases of toxicity from extremely high doses being intentionally taken for prolonged periods of time (sometimes under the direction of a health care practitioner) were rarely severe.(92,93)

### 1.2.2 Controversy over appropriate 25(OH)D goals

In 2014, Veugelers and Ekwaru asserted that the statistical calculations to determine recommendations for vitamin D were incorrectly interpreted, leading to a US RDA (600IU, or 700IU/day for those over 70) that is off by a factor of more than 10.(85,94) Heaney, et al., supported the higher level in a reply, citing a recent supplementation study which supported an RDA closer to 7000IU/day.(95) All three groups used the goal of 20ng/ml for musculoskeletal health.(85). In contrast, the Endocrine Society, aiming to optimize immune health and other



aspects of vitamin D function, recommends adults take in 1500IU - 2000IU per day to maintain a 25(OH)D level of 30ng/ml; 30ng/ml is the NIH target level as well.(39,85)

Controversies regarding appropriate 25(OH)D, are also informed by studies of parathyroid hormone levels.(39) Parathyroid hormone levels were not reduced in participants taking 15,000IU/day, even with 25(OH)D levels above 60ng/dl, in a study with a goal of bringing 25(OH)D levels up to at least 40ng/dl.(91) Mean serum calcium levels were not increased from baseline.(91) 25(OH)D levels of up to 120ng/dl appeared safe, and hypercalcemia and hypercalciuria were least common in participants with the highest 25(OH)D levels (calcium was not supplemented).(91) Goal 25(OH)D levels were achieved by 70% of the participants with 6000IU/day for normal weight participants, but 7000IU and 8000IU was required for overweight and obese participants, respectively.(91)

Growing research suggests that 40-60ng/ml is needed for prevention of respiratory infections, and 50-80ng/ml is required to favorably influence hypertension and cardiovascular disease.(38) In a 2019 randomized controlled trial, subjects without deficiency (initial 25(OH)D < 25ng/ml) who took 10,000IU/day for 3 years were slightly less likely to suffer a serious adverse event than those taking 400IU/day.(96) Mean 25(OH)D levels in the 400IU/day group did not increase, while 25(OH)D for the 10,000IU/day and 4000IU/day groups rose and then plateaued at 58ng/dl and 53ng/dl, respectively.(96)

### **1.2.3 Controversy over recommended supplement doses**

Recommended upper limits of vitamin D supplements in the USA were relaxed after several studies demonstrated that 4000IU of vitamin D daily is safe.(38,73,89<sup>P</sup>) One review showed that 10,000IU daily seemed to be the upper limit of tolerability.(73) The Endocrine Society recommends up to 10,000IU/day, particularly for obese individuals.(91,97<sup>P</sup>) However, some study participants have taken 15,000IU to 40,000IU daily for at least 6 months without apparent adverse effects.(89<sup>P</sup>)

The European Society for Clinical Nutrition and Metabolism recommends a one-time dose of 500,000IU IV for ICU patients who are vitamin D deficient (25(OH)D less than 20ng/ml), based upon evidence that this practice decreases length of stay.(9,16,98) Giving 500,000IU enterally over 5 days increased 25(OH)D levels and decreased ICU length of stay, but giving the entire 500,000IU in one bolus enterally did not improve 90-day mortality rates.(99,100)

Grant, et al. authored an early article positing a relationship between COVID-19 and vitamin D which recommended 10,000IU/day for one month, followed by 5000IU/day, with a goal 25(OH)D of 40-60ng/ml.(16) Although some other researchers agreed, many were outraged [**Tables 2 and 3, appendix**]. Kow, et al., questioned both the dose and the goal, citing a robust study in which supplements decreased the incidence of acute respiratory tract infections only when 25(OH)D levels were less than 10ng/ml, and 800IU/day was sufficient.(101,102) Grant, et al., replied with several additional studies to support their recommendation of 40-60ng/ml as a goal, but included an example of significantly decreased incidence of respiratory infections with lesser vitamin D<sub>3</sub> doses (although 800IU was inferior to 2000IU, it still provided significant benefits over the placebo).(103,104)

Sharma, et al., reviewed the literature informing decisions about COVID-19 and vitamin D<sub>3</sub>, finding compelling evidence for 10,000IU/day for a month, followed by 5000IU/day to bring 25(OH)D levels up to the target of 40-60ng/ml, then recommended a more modest 1000-2000IU/day.(105) One group, Quesada-Gomez, et al., posited that vitamin D supplementation should be with oral calcifediol.(106) However, the majority of researchers and commenters recommend vitamin D<sub>3</sub> supplements of 1000IU - 4000IU during “COVID-19 times,” with a goal of achieving 25(OH)D levels of 30ng/ml (see citations for **Table 1** and **Section 4**). (107)

A meta-analysis of vitamin D supplementation to prevent acute respiratory infections found that daily vitamin D supplementation was safe and provided modest protective benefits, rising to a 70% protective effect when deficiency was corrected.(102,108) However, studies also found that large bolus doses are not particularly beneficial.(102) Effective study doses of vitamin D were most often in the range of 400 – 2000IU (10 - 50mcg), with the higher doses being given to adults.(102) A 2020 study of pregnant women also found that daily supplementation is superior to boluses, that 2000IU/day was sufficient to resolve deficiency over time, and that up to 5000IU/day is safe.(109)

## 2 Methods: Literature Search

“COVID-19” is the MeSH term for SARS-CoV-2 disease, coronavirus 2019, COVID-19, and derivative terms. The topic of COVID-19 is a relatively new one, with the first reports published only 6 months ago (January 2020). In addition, because vitamin D supplementation is controversial, publication bias is a significant concern. Consequently, a significant percentage of the pertinent literature is found only on preprint services, most of which are captured by Google Scholar. PubMed casts a wider net than MEDLINE. Therefore, initially, PubMed and Google Scholar were searched for “COVID-19 AND “Vitamin D”” (date range, 2020, omitting citations and patents, no language limitations). Repeated searches confirmed the growing interest in the hypothesis that vitamin D deficiency may play an important role in the COVID-19 pandemic.(7,9,16,22,40<sup>P</sup>,108,110<sup>P</sup>,111-116,117<sup>P</sup>,118–125) From May 2 to May 19, Google Scholar hits increased from 49 to 88 and PubMed hits increased from 17 publications to 32. By June 16, the Google Scholar search retrieved 158 possible references and the PubMed publications on the topic had increased to 69. Using the same search terms, the author also accessed the Royal Society of Medicine Library Discovery Service, which, on 16 June, 2020, provided 144 results from academic journals, reports, magazines, and electronic resources.

Duplicates were deleted and full texts obtained for every publication from all three sources as of June 16, 2020, references were scanned for additional sources, and appropriate articles found on ResearchGate and through other internet sources were added to capture how the topic is being addressed in the popular press. Authors of perspectives and studies on this topic span the globe (**Figure 2a & b**). Most of the research publications are quite brief, and many of the PubMed indexed articles are expert summaries of relevant data supporting the biological plausibility of the hypothesis, rather than reports of original research. Therefore, the author deemed it premature to limit this review to the “best evidence” as one would do in a formal systematic review of the literature. Rather, every publication discussing vitamin D with respect to COVID-19 found by the three formal searches as of June 16, 2020 is included in **Table 2** (Original Research, n=47) or **Table 3** (Plausibility and In Vitro Studies, n=141). All original research

studies (excepting in vitro) are summarized in the Results (Section 3). However, due to space limitations, while many of the Table 3 documents are cited, few are summarized individually.

Figure 2a Interest in the Topic - Plausibility Article Authors' Locations (141 Articles)





Figure 2a Interest in the Topic - Study Article Authors' Locations (47 Articles)



(Blank maps from wiki images: Creative Commons, credit: Canuckguy and many others)

### 3 Results of Searches

#### 3.1 Planned Clinical Trials

Formal clinical trials that include COVID-19 and vitamin D (including 16 on [clinicaltrials.gov](https://clinicaltrials.gov)) include: vitamin D boluses plus other medications for COVID-19 positive patients, boluses or moderate daily doses to prevent severe complications in at-risk populations, low-dose vitamin D as a placebo in a drug trial, genetic variant studies focused upon the interaction between vitamin D and COVID-19, and studies of vitamin D levels in patients with differing severities of COVID-19 illness.(29,126–143) As late as May 19, 2019, few studies had begun recruiting. Although most of these studies are not designed to determine if daily modest vitamin D supplementation decreases either the risk of contracting COVID-19 or its severity, Dr. Manson plans to test this hypothesis.(64)

Testing the hypothesis that vitamin D deficiency prior to contracting the virus increases COVID-19 rates and severity necessitates screening participants for deficiency at enrollment. Failing to correct a deficiency being considered as a potentially significant risk factor for fatal COVID-19 complications would be unethical.(144<sup>P</sup>) Therefore, before and after population study designs (recommending supplements to groups known to be vitamin D deficient and observing if the groups' fatality rates decline) might be more feasible than randomized controlled trials.(144<sup>P</sup>)

#### 3.2 Biological Plausibility Discussions

Many of the reports examining the relationship between vitamin D and COVID-19 present biological plausibility arguments. These reports are summarized in both **Table 2** and **Table 3**. The main arguments are presented concisely here, citing both COVID-19 specific and primary sources.

##### 3.2.1 Vitamin D enhances resistance to viral illnesses

Early studies of vitamin D supplementation for acute respiratory tract infections produced conflicting results.(38,102) Most studies of vitamin D for influenza prevention were conducted on healthy populations with high baseline levels, rather than on the deficient populations who would benefit most.(16,41,119,123,145) Despite this, some found that higher 25(OH)D linearly enhance the innate immune response to acute winter respiratory infections, halving the incidence and significantly reducing the duration of illness.(42,146,147)

In 2017, 25 international researchers from 23 institutions performed a meta-analysis of individual participant data from 25 high-quality randomized controlled trials of vitamin D supplementation to prevent acute respiratory tract infections to determine why the results were inconsistent.(102) They found that bolus doses were not consistently protective, even in severely vitamin D deficient populations.(102) Removing bolus-dose data led to consistent findings of benefit, regardless of initial vitamin D status.(102) Daily or weekly vitamin D supplementation was most beneficial for participants with baseline 25(OH)D <10ng/ml (severe deficiency), providing more statistically significant ( $p < 0.001$ ) protection than the ( $p < 0.02$ ) protection vitamin D provided less deficient participants.(102) The authors found that response to vitamin D

supplementation is so variable that studies should base findings on changes in 25(OH)D levels, rather than relying upon the vitamin D dose given to each participant.(41,102) They also found that vitamin D supplementation is extremely safe: even large doses did not increase risk of serious adverse events, such as renal stones.(102)

Historical data provides modest support for the hypothesis that populations with high vitamin D deficiency rates allow new pandemic viral strains to propagate more freely. The only recorded time period void of new strains of pandemic influenza is 1920 – 1957, and vitamin D food supplementation was most prevalent during the middle of that time period: from 1930 to 1950.(61<sup>P</sup>) The COVID-19 pandemic began in Wuhan during a particularly dark January: 42% darker than their average January in 13 years (2007 - 2020).(148<sup>P</sup>)

Several studies have shown that vitamin D decreases the severity of dengue fever.(149) Oral vitamin D supplements of 4000IU for 10 days were significantly more effective than 1000IU in reducing dengue virus replication and controlling the damaging cytokine hyper-reaction.(149–152) Vitamin D supplementation also reduced rotavirus replication in pigs.(152) A recent review article by Sharma, et al., summarized biological plausibility arguments and found that vitamin D deficiency is associated with a wide range of viral illnesses, and that vitamin D supplementation was both preventative and decreased severity, limiting hyper-inflammatory complications.(105)

In the lungs, formation of the peptide LL37, an innate immune system component that, among other things, attacks enveloped viruses such as SARS-CoV-2 and modulates the immune system, requires sufficient vitamin D levels.(38,43,115,153) LL37 is inhibited by carbon and other nanoparticles in air pollution.(43) Therefore, vitamin D deficient individuals can be expected to be at increased risk of both developing COVID-19 and experiencing the “cytokine storm” if they become infected, particularly in areas of the world with high levels of air pollution.(43,115)

### **3.2.2 How vitamin D may decrease serious COVID-19-associated complications**

During the “Spanish flu” pandemic of 1918-1919, deaths were substantially reduced when patients were treated in “open air” hospitals with access to sunlight, perhaps due to vitamin D’s “cytokine storm” suppression.(61<sup>P</sup>,148<sup>P</sup>,154) In the deep south, dramatically increased incidence of pneumonia led to much higher Spanish flu case fatality rates for African Americans than for whites.(16) COVID-19 usually produces mild symptoms in the seemingly-vulnerable homeless, who are disproportionately outdoors, despite the population skew towards older males and African Americans.(155<sup>P</sup>,156,157) Prior to antibiotics, cod liver oil, UVB phototherapy, and sunshine, all of which are vitamin D sources, were considered successful treatments for tuberculosis.(97<sup>P</sup>)

Vitamin D enhances the innate immune response while, paradoxically, protecting against excessive inflammation by suppressing TNF $\alpha$  and the cytokines (e.g. IL-6, IL-17) implicated in severe COVID-19, and elevating anti-inflammatory IL-10.(16,29,38,42,44<sup>P</sup>,89<sup>P</sup>,147,158–167) Many of the articles referenced here include detailed descriptions of the role of vitamin D in preventing a “cytokine storm” and several authors, including Meftahi, et al., and Biesalski, added a series of cartoons to their papers to simplify the concept.(165,166)

Given that vitamin D decreases pro-inflammatory IL-6 and that IL-6 is implicated in the COVID-19 “cytokine storm,”(168) and finding that mean IL-6 levels are higher in males and African Americans and increase with age and obesity (groups with increased risk for COVID-19 mortality), Silberstein went on to evaluate the possibility that vitamin D deficiency causes upregulation of IL-6 in high risk individuals prior to exposure to COVID-19, increasing their likelihood of developing fatal COVID-19 complications.(169<sup>P</sup>) Using detailed IL-6 data from Tuscany, Italy, Silberstein found a strong correlation between age stratified COVID-19 deaths in Italy and mean IL-6 levels ( $r(6)=0.9837$ ,  $p=0.00025$ ). (169<sup>P</sup>) Data for a similarly detailed analysis for sex, obesity, and ethnicity was not available.(169<sup>P</sup>) The authors note that IL-6 is generally low in children, but it is high for a brief time in early childhood, which could explain the Kawasaki-like COVID-19 sequela in some children.(169<sup>P</sup>)

Vitamin D also helps prevent viral infections from progressing to pneumonia by tightening cell junctions.(16,38,163,170) And, vitamin D’s influence on the coagulation pathway decreases risk of acute respiratory distress syndrome as it decreases thrombosis risks.(112,125,155<sup>P</sup>,160,167,171) Therefore, correcting vitamin D deficiency might help prevent COVID-19 illness AND help limit complications when prevention is unsuccessful.(22,38,112,167)

Daneshkhan, et al., proposed that Vitamin D deficiency causes C-reactive protein (CRP) levels to rise, thus increasing the likelihood of a cytokine storm.(172<sup>P</sup>) The authors found that CRP and vitamin D status are inversely related in healthy individuals.(172<sup>P</sup>) CRP levels were increased in severe COVID-19 patients, but because CRP is a marker for inflammation, it was unclear if this was a cause or an effect.(172) The authors used population data from 10 countries to show a possible link between vitamin D status and the adaptive average case mortality ratio, and provided significant biological plausibility arguments in support of the hypothesis.(172<sup>P</sup>) The authors proposed further studies to determine if COVID-19 patients with high CRP are deficient in vitamin D.(172<sup>P</sup>)

### **3.3 Risk for Severe COVID-19 Parallels Risk for Vitamin D Deficiency**

Many authors, some with compelling statistical analyses, propose vitamin D deficiency from low sunlight levels (Nordic countries have high vitamin D intake) to explain the geographic distribution of severe COVID-19.(6,7,23,45,110<sup>P</sup>,120,148<sup>P</sup>) Italy and Spain have very high vitamin D deficiency rates.(9,37,45,61<sup>P</sup>,120,173) First-generation non-Western immigrants, even in countries with low overall rates, are often vitamin D deficient.(174–176) Vitamin D deficiency is especially common in the elderly, in part because synthesis from sunlight is muted in old age.(16,42,45,114,120,177-179,180<sup>P</sup>). Naturally melanin-rich skin increases vitamin D deficiency risks, particularly in high latitudes.(10,45,110<sup>P</sup>,114,120,146,179) It takes significantly more sunlight exposure for someone with dark skin to attain the benefits that someone with lighter skin receives.(11<sup>P</sup>) Lower 25(OH)D is associated with diabetes, hypertension, cardiovascular disease, and COPD risk.(16,22,42,45,181) Dialysis patients are often severely vitamin D deficient.(182) Up to 50% of US nursing home patients, and 75% of institutionalized people in general, are vitamin D deficient.(120,180<sup>P</sup>,183)

The UK's low sunlight have been posited in the public press as an explanation for why health workers with naturally melanin-rich skin (mostly nurses and physicians) are so disproportionately represented on the Telegraph's tribute wall.(184,185) The only postpartum COVID-19 fatality in the UK was a vitamin D deficient diabetic Pakistani woman who suffered a thrombotic complication.(186)

Current increased "stay at home" regulations and increased boredom and stress can be expected to result in eating patterns which increase obesity and the comorbidities with which it is associated.(187) In part because vitamin D is fat-soluble, obese individuals have increased daily vitamin D intake requirements and are often deficient.(22,45,55,89<sup>p</sup>,173,188,189) In addition, vitamin D deficiency causes the body to store more fat by increasing parathyroid hormone levels.(190) Obesity is a major risk factor for fatal COVID-19 complications, particularly in younger adults.(45,188) Ekiz, et al., found that increasing vitamin D levels makes it easier to lose excess weight, which could lower individual COVID-19 risk.(190)

Recent studies in Ireland and Switzerland both found that older males are at even higher risk of vitamin D deficiency than older females.(70,122) Vitamin D deficiency increases the X-chromosome linked 'Renin-Angiotensin' System (RAS) activity, making men more susceptible to ACE2 receptor dysregulation and theoretically, to increased COVID-19 morbidity.(22,44<sup>p</sup>,121,163,191,192) Although vitamin D deficiency is not universal in severe COVID-19, every deleterious symptom can be explained by RAS over-reaction, which would occur more easily in individuals without sufficient vitamin D to control the RAS.(22,124,163,181)

### **3.4 Evidence Informing the Hypotheses that Vitamin D Deficiency Influences COVID-19**

While data from randomized controlled trials is superior, the hypothesis that vitamin D deficiency is a major contributor in COVID-19 risk and severity is already supported by twenty population-data analyses, both causal inference modeling reports, four case studies/series, one prospective correlational study, one case control study, one cohort observational study, and ten retrospective chart reviews.. One population-data analysis and three retrospective chart reviews supported the dissenting view. One population-data analysis, one retrospective chart review, and the lone systematic review were neutral. Recognizing that truth is not exposed by the mere tallying of positions, but rather, by evaluating the specifics of the data and the strength of the study designs, all 47 studies are summarized here and in Table 2.

#### **3.4.1 Analyses of population data**

Bäcker asked whether temperature or radiance could explain the speed and level of geographic spread of COVID-19.(148<sup>p</sup>) Every location with over 2000 cases by March 15, 2020 had an average temperature of 10°C or lower.(148<sup>p</sup>) And, over a longer time period, locations with 4-week temperature averages under 14°C when they reached 100 cases all had faster growth than any of the warmer locations (p=0.0001).(148<sup>p</sup>) The same analysis using deaths instead of cases yielded a similar negative correlation (p<0.02).(148<sup>p</sup>) However, Finland, Norway, and Russia, all reaching 2000 cases after March 15, did not conform to the pattern, leading to a study of sunlight.(148<sup>p</sup>) Indeed, irradiance and cloudopacity better accounted for all the of data

( $p < 0.01$ ).<sup>(148<sup>P</sup>)</sup> Bäcker suggests (with data to back up his hypothesis) that increased cloudiness and air pollution can explain why in Korea, Daegu had ten times as many cases of COVID-19 as more internationally-connected Seoul, and in Italy, Lombardy had over ten times as many cases as more internationally-connected Lazio (Rome).<sup>(148<sup>P</sup>)</sup> Multivariate regression found that the best independent predictor of COVID-19 case ( $p < 0.001$ ) and death ( $p < 0.001$ ) growth rates was the average zenith (most direct sun rays) when the location reached its 100<sup>th</sup> case.<sup>(148<sup>P</sup>)</sup> Zenith, correlated with both irradiation ( $p < 0.01$ ) and temperature ( $p < 0.001$ ), explained the lower growth rate in Finland, Norway, and Russia, and fully accounted for the variance from both.<sup>(148<sup>P</sup>)</sup> No association with increased travel or visiting was found.<sup>(148<sup>P</sup>)</sup> Bäcker concluded that sunlight leads to less COVID-19 transmission, likely due to both the direct result of irradiation and increased vitamin D, and thus, advising people to stay indoors rather than opening up outdoor recreation areas during the COVID-19 pandemic appears to be a poor choice.<sup>(148<sup>P</sup>)</sup>

In contrast, Yao, et al., found no association between COVID-19 transmission rates and temperature or UV radiation across the 62 cities (of 224) in China with at least 50 cases at the peak of the outbreak (10 Feb) and at least 10 cases remaining on 9 March.<sup>(193)</sup> However, the authors note that their study examined data from early January to early March, 2020, a time during which strict travel restrictions were put into place to prevent COVID-19 transmission in China.<sup>(193,194)</sup> It is possible that many of the cities in which UV light or temperature effectively reduced transmission were eliminated from the study because they no longer had the minimum of 10 cases by 9 March.

Two statistical analyses of geographical areas in the USA addressed the question of whether high COVID-19 fatality rates in African Americans could be explained by income levels.<sup>(11<sup>P</sup>,195)</sup> Bäcker's initial statistical analysis from 8 cities and states that provide a racial breakdown of COVID-19 victims found that race-based fatality rate differences diminish in direct proportion to available sunlight, with COVID-19 deaths among blacks in Detroit at 193% higher than the percent-black area population, but only 7% higher in Florida (Pearson  $-0.76$ ,  $p < 0.05$ ).<sup>(11<sup>P</sup>)</sup> African Americans comprise 26% of Milwaukee County's population, half of their COVID-19 cases, and 81% of its deaths.<sup>(11<sup>P</sup>)</sup> This led to an exploration of the hypothesis that rather than socioeconomics (lower incomes, jobs that do not permit social distancing) being solely responsible, irradiance may play a large role in the disproportionate COVID-19 morbidity and mortality rates among African Americans in the USA.<sup>(11<sup>P</sup>)</sup> In Michigan, the state with the highest racial disparity in COVID-19 deaths, a county-by-county analysis showed that percent African American, but not percent over 65 years, median income, median age, or number of people per household, significantly ( $p < 0.05$ ) correlated with COVID-19 morbidity.<sup>(11<sup>P</sup>)</sup>

Similarly, Li, et al., focused on US counties with at least 50 COVID-19 cases (661 counties) and those with at least 10 deaths (221 counties), grouping them into quartiles and comparing highest to lowest.<sup>(195)</sup> Multivariate analysis demonstrated that "percent black" predicted county cases and fatalities, even after controlling for other demographics, socioeconomics, and comorbidities.<sup>(195)</sup> Higher daily temperatures decreased county case numbers, but not mortality rate.<sup>(195)</sup> They proposed vitamin D deficiency among black Americans as a "unifying theory" to explain their results.<sup>(195)</sup>



Laird, et al., plotted COVID-19 mortality/million against mean 25(OH)D levels for twelve European countries, finding a significant correlation ( $p=0.046$ ).<sup>(37)</sup> Panarese and Shahini ranked the 108 countries with at least 100 COVID-19 cases on 2 April 2020 by latitude, demonstrating visually that, overall, deaths per million were higher in the northern-most countries, whose citizens would be the most likely to be vitamin D deficient from the dark winter.<sup>(196)</sup> Following up on Panarese and Shahini's work, Rhodes, et al., compared the 120 countries with more than 150 COVID-19 cases by 15 April 2020, finding that COVID-19 mortality rates were significantly correlated with latitude ( $r=0.53$ ,  $p<0.0001$ ).<sup>(23)</sup> Rhodes, et al., used a simple scatter-graph to illustrate that the COVID-19 mortality rates per million population were dramatically lower in countries with capitals south of  $35^{\circ}\text{N}$ , where sunshine in the time immediately preceding the pandemic made maintaining vitamin D levels possible.<sup>(23)</sup>

Ilie, et al., reported a significant correlation between low mean vitamin D levels across 20 European countries and both COVID-19 fatalities/million population ( $p=0.05$ ) and COVID-19 cases/million population ( $p=0.050$ ).<sup>(120)</sup> Kumar and Srivastava objected to the study by Ilie, et al., stating that the correlation was being stretched by the media to claim that vitamin D supplements may reduce COVID-19 mortality rates by 50%.<sup>(197<sup>P</sup>)</sup> Purportedly concerned that this exaggerated claim would lead to fatal overdoses, the authors conducted a statistical analysis of COVID-19 case and death rates and life expectancy using the data from Ilie, et al.<sup>(120,197<sup>P</sup>)</sup> The authors asserted that because vitamin D deficiency increases with age, controlling for life expectancy would reveal the true relationship between vitamin D and COVID-19 infection and fatality rates.<sup>(197<sup>P</sup>)</sup> The researchers found that life expectancy was a better predictor of both COVID-19 mortality and case rates than vitamin D.<sup>(197)</sup> The authors called for clinical trials of vitamin D supplements.<sup>(197<sup>P</sup>)</sup>

Citing Ilie, et al., Singh, et al., compared mean 25(OH)D levels and COVID-19 cases and deaths per million population for 20 European countries on 8 April and again on 12 May.<sup>(198<sup>P</sup>)</sup> The significance of the inverse correlation between vitamin D and case rates from ( $r(20): -0.4435$ ;  $R^2 = 0.1967$  ( $p=0.0501$ ) in April to ( $r(20): -0.5504$ ;  $R^2 = 0.3029$  ( $p=0.0119$ ) in May.<sup>(198<sup>P</sup>)</sup> However, the inverse correlation for death rates decreased from ( $r(20): -0.4378$ ;  $R^2 = 0.1917$  ( $p=0.0535$ ) to ( $r(20): -0.3935$ ;  $R^2 = 0.1549$  ( $p=0.0860$ ).<sup>(198<sup>P</sup>)</sup> Singh, et al., did not discuss the possibility that vitamin D levels increased between April and May as sunshine increased, potentially protecting patients from the "cytokine storm."<sup>(198<sup>P</sup>)</sup>

Notari and Torrieri's much larger, more detailed, comprehensive 126 country data review found that most of the 24 identified potential risk factors for COVID-19 propagation, including blood type, life expectancy, and even greeting habits, were significantly correlated with one another.<sup>(199)</sup> Prevalence of Type-I diabetes, BCG vaccination, and vitamin D levels were the only "almost independent factors."<sup>(199)</sup> In the 42-country subset with vitamin D data and high GDP, lower mean annual levels of vitamin D were linearly related to increased COVID-19 risk ( $p=0.006$ ), with seasonal values (March) demonstrating even more significance ( $p=0.002$ ).<sup>(199)</sup>

Kara, et al., mapped the population prevalence of vitamin D deficiency ( $<20\text{ng/ml}$ ) and severe deficiency ( $<10\text{ng/ml}$ ) against COVID-19 total fatalities for the 40 most affected countries, worldwide, finding a clear relationship.<sup>(45)</sup> Regression analyses demonstrated a quadratic relationship between prevalence of vitamin D deficiency and insufficiency and COVID-19

cases.(45) A histogram with regression lines illustrated the relationship between latitude, population vitamin D status, and country rank (by number of cases).(45) Finding vitamin D deficiency and COVID-19 to be related pandemics, they agreed with Grant, et al., in recommending vitamin D (without high calcium) supplementation, as well as encouraging fortified food intake and increased sun (UVB) exposure.(45)

Braiman noted that as of March 22, 2020, although 10% of the COVID-19 cases lived south of the Tropic of Cancer, they represented only 1% of the fatalities.(144<sup>P</sup>) The three exceptions could all be explained by mean population vitamin D levels.(144<sup>P</sup>) Nordic countries have vitamin D deficiency rates below 1% (due to diet or supplementation) and impressively low COVID-19 fatality rates, except Sweden.(144<sup>P</sup>,200) In Stockholm, severe vitamin D deficiency is common among displaced Somalis, who with less than 1% of the population have suffered 40% of the COVID-19 fatalities.(7,176,201) Indonesia straddles the equator, but its predominately Muslim women have vitamin D levels that are only half that of notoriously low Italy.(144) Sunscreen use is popular in the Philippines, which may account for the high levels of vitamin D deficiency there.(144<sup>P</sup>) Braiman recommended ethical testing of the hypothesis that vitamin D and COVID-19 outcomes are related by encouraging supplementation in deficient populations and evaluating death rate changes.(144<sup>P</sup>)

Although Latinos and African Americans were found to be at higher risk of COVID-19 mortality in New York City, it is difficult to determine the influence of Latino ethnicity versus race (over 75% of Latinos identify as non-white), because New York City does not provide sufficiently detailed data.(202<sup>P</sup>) In contrast, Georgia does break down COVID-19 data by both ethnicity and race.(202<sup>P</sup>) Black Latino COVID-19 morbidity was 123% higher than white Latino morbidity ( $p<0.001$ ), supporting researcher Bäcker's hypothesis that a darker complexion decreases sun exposure benefits.(202<sup>P</sup>) COVID-19 morbidity is 37% higher for white non-Latinos than for white Latinos ( $p<0.0001$ ), 689% higher for Native American non-Latinos than for their Latino counterparts ( $p<0.01$ ), and there were no cases of COVID-19 among Latino Asians.(202<sup>P</sup>) Latinos spend more time outdoors than any other racial group (85% more than African Americans) which could explain why Latinos defied externally-imposed racial disparity explanations.(202<sup>P</sup>)The author concluded that irradiance exposure seems to help prevent COVID-19.(202<sup>P</sup>)

Countries with higher rates of vitamin D-rich sea fish consumption or food supplementation have lower COVID-19 mortality rates than adjacent countries.(180<sup>P</sup>) The elderly, especially in nursing homes, where 84 - 93% of residents in the US are vitamin D insufficient, are at highest risk for severe COVID-19.(180<sup>P</sup>) Bäcker and Mageswaran evaluated vitamin D deficiency rates among elderly females and COVID-19 deaths prior to May 31<sup>st</sup> in 32 countries, finding that case fatality rates were up to twice as high in countries with high vitamin D deficiency rates ( $p<0.04$ ). (180<sup>P</sup>) They also found that case fatality rates were significantly higher ( $p<0.026$ ) in countries with a high percentage of black inhabitants.(180<sup>P</sup>) Noting many biological plausibility arguments and vitamin D deficiency and insufficiency race disparities, the authors recommend COVID-19 prevention and treatment studies.(180<sup>P</sup>)

Li, et al., used machine learning to produce logistic models to predict case rates, death rates, and case fatality rates in all 50 US states and 154 countries listed on the Johns Hopkins COVID-19 dashboard on 15 May 2020, assessing the interdependence of the 57 factors LASSO identified as potentially influencing COVID-19 outcomes.(203<sup>P</sup>) Among their many findings, Lee, et al., determined that higher population vitamin D intake is an independent factor in reduced COVID-19 cases.(203<sup>P</sup>)

Kohlmieir performed a Mendelian randomization to test the effect of latitude (a proxy for vitamin D) on rates of African American COVID-19 deaths in the 22 reporting states with more than 15 African American deaths as of 16 April 2020, finding a strong relationship ( $r=0.427$ ). (204) A correlational analysis found that excess mortality rates were significantly higher ( $r=0.435$ ,  $p=0.02$ ) in states with higher latitudes.(204) The highest excess mortality rates were all in states near or above  $40^{\circ}$  N, where UVB intensity in winter and spring is too low to provide vitamin D.(204) The African American fatality over-representation was 5.6-fold in Wisconsin compared with 1.3-fold in Florida.(204)

Adding the proposed relationship between latitude and COVID-19 to knowledge that ozone filters the ultraviolet-B radiation the body requires to produce vitamin D, Alipio evaluated data from all 34 countries with April 2019 ozone data available on an open-access database.(205<sup>P</sup>) Kendall rank correlation test found that ozone concentration significantly ( $P<0.001$ ) positive correlated with COVID-19 cases, but latitude and COVID-19 cases appeared to have no relationship.(205<sup>P</sup>)

Recognizing the advantages of comparing cities within a single country with varied UV radiation, altitude, and weather patterns, such as consistent policies, culture, and genetic factors, Skutsch, et al., conducted a multiple regression analysis of data from 45 cities in Mexico, comparing the rate of increase in cumulative COVID-19 cases and fatalities.(206<sup>P</sup>) Data from January was included because, while UV light's sterilization effect would be immediate, physiologic vitamin D formation precedes its impact on infection and mortality rates.(206<sup>P</sup>) Skutsch, et al., found a negative relationship between rate of transmission and altitude ( $r=-0.354$ ,  $p=0.014$ ), but temperature, relative humidity, and latitude were insignificant.(206<sup>P</sup>) UV levels in January correlated a bit more strongly with transmission rates ( $r=-0.369$ ,  $p=0.014$ ) than UV levels during the transmission period ( $r=-0.32$ ,  $p=0.032$ ), supporting the hypothesis that the influence of UV is due to vitamin D rather than sterilization.(206<sup>P</sup>) In contrast, UV was only marginally associated with rates of mortalities.(206<sup>P</sup>) Mexico City's air pollution may have explained this.(206) Surprisingly, altitude and UV levels were not significantly interrelated, but their combined effect accounted for 18% of transmission rate variation ( $p=0.0062$ ). (206<sup>P</sup>) Data for 834 individuals scattered across 561 municipalities showed that lower altitude is a highly significant ( $r=-0.35$ ,  $p=0.0005$ ) predictor of vitamin D levels, perhaps influenced by the high levels of UV light in coastal cities and the cooler climate of higher altitude cities leading to more clothing coverage.(206<sup>P</sup>)

Noting that all five US states with fatalities greater than 5000 and four of the five states with cases over 90,000 are in latitudes above  $37^{\circ}$ N, Li, et al., used latitude as an indicator to evaluate the relationship between sunlight, vitamin D, and COVID-19 case and death rates per 1000,000

population.(207<sup>P</sup>) Aggregate data (22 Jan – 23 May 2020) showed that states in latitudes above 37°N, when compared with states at lower latitudes, had significantly higher case rates (702 vs 255/100K) and death rates (43 vs 11 deaths/100K) ( $p < 0.001$ ).(207) The higher case rates were not attributable to higher test rates.(207<sup>P</sup>) The authors suggested sunlight and vitamin D as the explanation, calling for studies to evaluate the impact of vitamin D on the prevention of COVID-19.(207<sup>P</sup>)

In a less detailed study, Marik, et al., evaluated the case fatality rates for all 50 US States, mapping the results to illustrate that, with the exception of states with very low population densities and Louisiana, case fatality rates increased with increasing latitude.(208) The cumulative summary case fatality rate for states over 40°N was significantly higher than for states below 40°N (6.0% vs 3.5%,  $p < 0.001$ ).(208) Attributing the differences to vitamin D's dampening of excessive inflammation, they advocated for standard vitamin D supplement doses and further studies.(208)

Moozhipurath, et al., obtained UVB radiation data for 108 days (through 8 May 2020) in the 152 countries with more than 20 COVID-19 cases, beginning when the country had over 20 cases, analyzing the relationship between daily UV index (UVI - a surrogate for UVB), COVID-19 deaths, and COVID-19 cases, controlling for weather variables, including ozone levels.(21<sup>P</sup>) UVI increase was associated with a 1.2% decrease in the daily growth rate of cumulative COVID-19 deaths ( $p < 0.01$ ) and a 1.0% decrease in the daily growth rate of cumulative COVID-19 case fatality rates ( $p < 0.05$ ).(21<sup>P</sup>) The authors asserted that their methods led to very conservative estimates of the effect of UVB on COVID-19 deaths, and advocated for “sensible” increased exposure to sunlight, particularly for people at high risk of vitamin D deficiency.(21<sup>P</sup>)


A statistical analysis by Davies, et al., found that COVID-19 outbreaks with large fatality rates occurred exclusively above 30°N, with a 55:1 ratio between 30°N - 55°N and more southern latitudes.(61<sup>P</sup>) The Epidemic Severity Index was greater than 2.5 in nine of 239 locations, all above 30°N.(61<sup>P</sup>) Northern outliers all had higher vitamin D population levels, southern countries with the most severe outbreaks (Philippines and Brazil) have a high vitamin D deficiency prevalence, and fatality rates are doubled by naturally melanin-rich skin in the USA and UK.(61<sup>P</sup>) Iran, where religious full-body clothing is worn and vitamin D deficiency is common, fared far worse than Israel, whose vitamin D deficiency prevalence is relatively low.(61<sup>P</sup>)

### **3.4.2 Causal inference modeling reports**

Davies, et al., also analyzed three potential root causes for their influence on COVID-19 outcomes, categorizing factors as lowering vitamin D, negatively influenced by low vitamin D, or vitamin D neutral.(61<sup>P</sup>) Environmental conditions hostile to the virus and environmental measures (e.g., distancing) decrease COVID-19 spread, but do not influence case fatality rates.(61<sup>P</sup>) If vitamin D is a “bystander” variable (simply a marker of bad health), case fatality rates would correlate best with vitamin D-neutral comorbidities.(61<sup>P</sup>) The authors provide a detailed analysis of the known COVID-19 epidemiological, latitude, and environmental data.(61<sup>P</sup>) A table illustrates that 16 predictions of the causal model accurately match the known

facts, while 14 predictions of the bystander model strongly contradict the data and two more are not supported [Figure 3].(61<sup>P</sup>)

Figure 3: Davies, et al., Causal Model Results(61<sup>P</sup>)



Annweiler, et al., used Hill’s methodology for determining causality, which states that the more of the seven criteria are met, the stronger the claim, to evaluate the hypothesis that vitamin D is causally linked to COVID-19 outcomes.(209) Vitamin D met six of the criteria, failing only on specificity (because vitamin D deficiency is high in the general population).(209) Concluding that vitamin D deficiency is highly likely to be a cause of poor COVID-19 outcomes, the authors suggest that these results, coupled with the excellent safety profile of vitamin D and lack of other treatments, support testing vitamin D as an adjuvant treatment and prophylaxis for the general population.(209)

### 3.4.3 Case Studies and Case Series in which vitamin D is mentioned

Ahmed, et al., reported that a COVID-19 positive maternity patient with diabetic ketoacidosis, vitamin D deficiency, and a history of asthma developed a fatal thrombosis four days post extubation.(186) Horowitz, et al., reported on two COVID-19 pneumonia patients with histories of immunosuppression from Lyme disease who responded to repeated doses of glutathione, along with a multitude of other drugs and remedies.(210) One had a history of low vitamin D.(210) Bossoni, et al., reported on a 72-year-old thyroidectomized COVID-19 positive patient who experienced sudden onset severe hypocalcemia.(211) Her parathyroid level was low, and she was extremely vitamin D deficient (8ng/ml).(211) Bossoni, et al., noted that home confinement can worsen vitamin D deficiency, increasing the risk of systemic infections and potentially life-threatening hypocalcemia.(211)

Vitamin D deficiency is common in Indonesia, affecting 35.1% of elderly institutionalized women and 23% of the general population.(212<sup>P</sup>) Pinzon, et al., tested ten PCR-positive COVID-

19 patients in Indonesia, finding that nine were vitamin D deficient (25(OH)D <10ng/ml) and the remaining patient was insufficient (25(OH)D=20.5).(212<sup>P</sup>) Finding no clinical evidence to inform the decision to provide vitamin D supplements to prevent or treat COVID-19 in their review of the literature, they called for randomized controlled trials and now prescribe all patients 2000IU/day.(212<sup>P</sup>)

#### **3.4.4 Prospective correlational study, case-controlled survey, and cohort observational study**

Vitamin D deficiency is common among Irish males (median 25(OH)D of 18.8 ng/ml for ages 40 – 60).(213) Faul, et al., drew 25(OH)D in 33 COVID-19 positive Caucasian males over the age of 40 who were admitted to the hospital in respiratory failure without cancer, diabetes, cardiovascular disease, or chronic immunosuppressant intake in Ireland in March of 2020.(213) The twelve requiring mechanical ventilation (including all four fatalities) had mean serum 25(OH)D levels of 10.8 ng/ml, compared with 16.4 ng/ml for those requiring only oxygen (p=0.03).(213). Patients with 25(OH)D <12 ng/ml had a hazard ratio for requiring ventilator care of 3.19 (p=0.03).(213) The authors concluded that low vitamin D is either a marker for poor health, or it permits pro-inflammatory changes that lead to severe COVID-19: “a thought worthy of further study.”(213)

Concerned about the effects of COVID-19 on their community-dwelling Parkinson’s Disease patients in Lombardy, Italy, Fasano, et al., conducted telephone interviews with 1486 patients and 1207 family-member case controls.(214) The 105 Parkinson’s patients with COVID-19 and 92 family members with COVID-19 differed only in decreased shortness of breath (p=0.004) and decreased hospitalization rates (p=0.018) for the Parkinson’s patients.(214) The authors adjusted the data for the age differences between groups, thought to be due to aggressive protective measures for the elderly in the area.(214) Parkinson’s, hypertension, and COPD medications did not influence the likelihood of developing COVID-19, while Vitamin D supplementation was protective (p=0.048).


Tan, et al., compared the 26 consecutive patients 50 years or older not requiring oxygen on admission who were hospitalized immediately prior to initiation of a daily oral combination of 1000IU vitamin D<sub>3</sub>, 150mg magnesium, and 500mcg vitamin B<sub>12</sub> with the next 17 consecutive patients meeting the same criteria to determine if these supplements altered support needs.(215<sup>P</sup>) Of the 9 patients supplemented within a week of symptom onset, only one required oxygen, and that was within 24 hours of supplement initiation.(215<sup>P</sup>) Of the 8 patients supplemented over a week after symptom onset, one required ICU care within 24 hours of supplement initiation, and one required oxygen three days later.(215<sup>P</sup>) Supplemented patients were less likely to need any oxygen (17.6% vs 61.5%, p=0.006) or ICU care (5.9% vs 30.8%).(215<sup>P</sup>)

#### **3.4.5 Retrospective chart reviews favoring the hypothesis**

Alipio performed a chart review using de-identified data from 212 COVID-19 patients with recorded pre-COVID-19 25(OH)D levels from three hospitals in Southern Asia in which 25(OH)D was tested initially and weekly.(40<sup>P</sup>) Individuals’ 25(OH)D levels did not vary significantly during hospitalization, confirming that battling COVID-19 does not, in and of itself,

deplete vitamin D.(40<sup>p</sup>) Vitamin D status (3 categories: >30ng/ml, 21-29ng/ml, or <20ng/ml) correlated significantly and linearly with more critical COVID-19 illness (4 levels clearly defined by previous researchers).(40) For each standard deviation increase in serum 25(OH)D, the odds of having a mild, rather than a critical, case of COVID-19 were almost 20 times as great (OR=0.051, p<0.001).(40<sup>p</sup>) [Figure 4]


Figure 4: Some of the Results of the Retrospective Chart Review by Alipio, et al.(40<sup>p</sup>)



Legend: Of the 212 hospitalized COVID-19 patients, 96% of those with mild COVID-19 had normal vitamin D levels (above 30 ng/ml). In contrast, over 50% of the patients with severe or critical COVID-19 were vitamin D deficient (level 20ng/ml or lower).

A statistical analysis by D’Avolio, et al., of records in a Swiss clinic’s database for 107 symptomatic individuals obtaining a SARS-CoV-2 PCR test found that the 27 PCR-positive patients had significantly lower (p = 0.004) 25(OH)D (11.1ng/ml) when compared with test-negative subjects (24.6 ng/ml).(122) PCR-positive patients were 70.4% male, while PCR-negative patients were only 48.8% male, with similar ages.(122) Differences between 2019 median 25(OH)D versus PCR-positive 2020 median 25(OH)D were significant for both women (25.6ng/ml vs. 9.3ng/ml, p = 0.019) and men (22.9ng/ml vs. 11.4ng/ml, p = 0.005), but not for PCR-negative for either gender [Figure 5].(122)


Figure 5: Some of the Results of the Retrospective Chart Review by D’Avolio, et al.(122)



Legend: In the 107 individuals tested for both 25(OH)D and COVID-19, vitamin D predicted infection

Lau, et al., found that among all 20 COVID-19 patients with recorded 25(OH)D at a New Orleans hospital, every ICU patient under age 75 had vitamin D insufficiency.(155<sup>P</sup>) Eleven of 13 ICU patients had vitamin D insufficiency versus 4 of 7 patients with milder COVID-19.(155<sup>P</sup>) Seven ICU patients had critically low 25(OH)D (<20ng/mL) and three had levels below 10ng/mL.(155<sup>P</sup>) Patients with the lowest 25(OH)D levels were African American [Figure 6].(155<sup>P</sup>)

Figure 6: Some of the Results of the Retrospective Chart Review by Lau, et al.



Legend: In the 20 hospitalized individuals tested for both 25(OH)D and COVID-19, vitamin D predicted severity of infection



In Jakarta, Indonesia, hospitals are designed to provide patients sunlight and home patients exercise outdoors.(216) In this setting, daily minutes of sunshine were compared with patient recovery, death rates, and incidence.(216) Asary and Veruswati found sunshine was not related to prevention, but COVID-19 patient recovery briskness was significantly (Spearmen's  $\alpha = 0.05$ ) correlated with sunnier days.(216)


Sun, et al., conducted a 241 patient retrospective chart review in a hospital in Wuhan, China, using standardized definitions of mild, moderate, severe, and critical COVID-19. (217<sup>P</sup>) On admission, 74.7% of patients were hypocalcemic.(217<sup>P</sup>) Noting that vitamin D deficiency can cause hypocalcemia, the researchers found a median 25(OH)D of 10.20ng/ml (severe deficiency) among the 26 patients tested; none were vitamin D sufficient.(217<sup>P</sup>) These 26 patients had worse CRP ( $p < 0.001$ ), D-dimer ( $p < 0.001$ ), and parathyroid hormone ( $p = 0.048$ ) levels.(217<sup>P</sup>) Calcium levels positively correlated with 25(OH)D levels ( $p = 0.004$ ), and lower calcium levels correlated linearly with lower SpO2 levels ( $p < 0.001$ ), higher complication rates ( $p < 0.001$ ), and higher 28-day mortality rates ( $p < 0.001$ ). (217<sup>P</sup>) Vitamin D deficiency and hypoproteinemia were associated with increased mortality in critically ill patients.(217<sup>P</sup>)

Cuñat, et al., found that although recommended for all ICU patients, vitamin D was tested in only 17 of the 226 consecutive COVID-19 patients admitted to their hospital in Spain.(218<sup>P</sup>) All 17 were vitamin D deficient (25(OH)D  $< 20$ ng/ml), 13 had  $< 12.5$ ng/ml, and three had  $< 5$ ng/ml.(218<sup>P</sup>) Of these 17 patients, 35.2% had hypocalcemia and 64.7% had hypophosphatemia.(218<sup>P</sup>) The incidence of nosocomial infections was very high (76.5%).(218<sup>P</sup>) The authors stated that vitamin D deficiency is especially problematic for COVID-19 ICU patients because vitamin D reduces pro-inflammatory and increases anti-inflammatory cytokines.(218<sup>P</sup>)

Raharusuna, et al., conducted a retrospective chart review of 780 hospitalized test-confirmed COVID-19 patients in Indonesia.<sup>P</sup>) After controlling for age, sex, and comorbidity, both insufficient (odds ratio 7.63) and deficient vitamin D (odds ratio 10.12) were significantly associated with COVID-19 mortality ( $p < 0.001$  for each).(46<sup>P</sup>) Fatalities were 4.1% in patients with normal 25(OH)D, 87.8% with insufficiency, and 98.9% with deficiency.(46<sup>P</sup>)

In India, Glicio, et al., performed a statistical analysis on the data from the 176 COVID-19 patients 60 years or older in two tertiary medical centers whose medical records included body mass index (BMI), sex, comorbidities, clinical characteristics, and pre-hospitalization 25(OH)D.(47<sup>P</sup>) Over 80% were vitamin D insufficient or deficient, and of those, 72% were male.(47) Inadequate 25(OH)D was strongly associated with chronic kidney disease, hypertension, and diabetes.(47<sup>P</sup>) Vitamin D levels were lower, with a linear distribution, in older patients (oldest age was 85).(47<sup>P</sup>)

Figure 7: Some of the Results of the Retrospective Chart Review by Glicio, et al.(47<sup>p</sup>)



Legend: In the 176 elderly (over age 60) hospitalized individuals tested for both 25(OH)D and COVID-19, vitamin D predicted severity of infection

Insufficient 25(OH)D was found in 45% of the 24 patients with mild COVID-19 versus 86% of the 131 patients with severe outcomes.(47<sup>p</sup>) As age increased, vitamin D levels correlated linearly with outcomes, with patients over 70 suffering severe COVID-19 only if they were vitamin D insufficient.(47<sup>p</sup>) In contrast with obese patients, those with healthy BMIs tended to have severe COVID-19 only if they were vitamin D deficient (also a linear correlation) [Figure 7].(47<sup>p</sup>)

De Smet, et al., found endemic vitamin D deficiency in their area of Belgium, with lower mean levels in men than women except in summer ( $p < 0.05$ ). (219<sup>p</sup>) Children under age 18 had lower deficiency rates ( $p < 0.05$ ). (219<sup>p</sup>) Comparing 186 consecutive test-positive COVID-19 patients (109 male) with the 2717 consecutive age-matched controls whose 25(OH)D was tested during the same season in 2019, they found that vitamin D deficiency was prevalent in controls (45.2%), but significantly ( $p < 0.05$ ) more common in the hospitalized COVID-19 patients (58.6%). (219<sup>p</sup>) The median 25(OH)D for COVID-19 patients was 18.6 ng/ml, compared with 21.5 ng/ml for controls ( $p = 0.0016$ ). (219<sup>p</sup>) Male patients were more likely than their control counterparts to be deficient (67.0% vs 49.2%,  $p = 0.0006$ ). (219<sup>p</sup>) Vitamin D deficiency was strongly associated with more severe COVID-19 pneumonia in males (55.2% with stage 1, 66.7% with stage 2, and 74% with stage 3,  $p = 0.001$ ), but not in females. (219<sup>p</sup>) Vitamin D was stable across all stages of COVID-19 for females, suggesting that the illness itself does not deplete vitamin D. (219) The authors argue that as a whole, their data supports a causal role for vitamin D deficiency in COVID-19. (219<sup>p</sup>)

Meltzer, et al., analyzed data from their US facility's COVID-19 positive patients with documented 25(OH)D levels in EPIC within the previous two years to determine if deficiency

increases COVID-19 incidence.(220<sup>P</sup>) Data for the most recent 25(OH)D and treatment (dose and time span) led to four categories 1) likely still deficient, 2) likely sufficient, 3) likely deficient but improved since testing, and 4) uncertain status.(220) Known risk factors and factors that influence vitamin D activation were evaluated.(220<sup>P</sup>) A multivariate analysis found that, of patients with 25(OH)D levels within the previous year, those likely to still be deficient (category 1) were more likely (RR=1.77, p<0.02) to test positive for COVID-19 than those likely to be vitamin D sufficient (category 2).(220<sup>P</sup>) Older age, non-white race, and immunosuppression were the only other factors associated with testing positive for COVID-19.(220<sup>P</sup>) Hypertension, obesity, and diabetes were not covariates with vitamin D.(220<sup>P</sup>) Vitamin D deficiency was associated with supplement type and dose (p<0.01), unless the relatively few patients receiving 2000IU or more of vitamin D<sub>3</sub> were omitted (indicating that lower doses, D<sub>2</sub>, and calcitrol did not improve deficiency).(220<sup>P</sup>) The authors concluded that the relatively low doses of vitamin D usually given to correct deficiency in their institution decreased the apparent benefit of supplementation on COVID-19 rates, and that 4000-5000IU/day may be indicated for COVID-19 prevention.(220<sup>P</sup>)

### **3.4.6 Retrospective chart reviews that are neutral or strongly oppose the hypothesis**

Fox and Sizemore evaluated the Electronic Health Records of over 15,000,000 patients in EPIC across 26 US states, finding 28,185 patients with documented 25(OH)D (of which, 86% were deficient) and a documented COVID-19 test.(83) No association was found between vitamin D deficiency (defined by each lab) and COVID-19 rates, hospitalizations, or fatalities.(83) In contrast with the study by Meltzer, et al., no date limits were placed on the testing; the authors noted that vitamin D levels are usually drawn to confirm suspected deficiency.(83) The authors recognized this limitation and recommended future studies including patients with normal vitamin D levels, along with studies to assess the effect of vitamin D supplementation on prevention or treatment of COVID-19.(83)

Hastie, et al., evaluated data from the 1474 participants in the UK Biobank study whose COVID-19 test results were available to them.(10) Rather than comparing the 1025 PCR-negative participants to the 449 PCR-positive patients, every person in the 348,598 database without a PCR-positive test result was assumed negative.(10) The 25(OH)D levels obtained 10-14 years prior were significantly lower in blacks and South Asians.(10) Black or South Asian ethnicity was also strongly associated (p < 0.001) with confirmed COVID-19 infection.(10) Median 25(OH)D was significantly lower (p = 0.013) for those with confirmed COVID-19 infection, and 25(OH)D predicted infection univariably.(10) In contrast, the multivariate analysis did not find 25(OH)D was significant.(10) Unlike most other studies of COVID-19, the authors found no association between diabetes or hypertension and COVID-19 risk, raising concerns that important variables were factored out in their analysis.(10,82)

Another review using 2006 – 2010 data from the UK Biobank was conducted by Darling, et al., who compared the vitamin D status, BMI, ethnicity, and lifestyle factors of 580 COVID-19 positive cases (including outpatients) with 723 negative controls of similar age.(74<sup>P</sup>) 25(OH)D levels were 3.6ng/ml lower (p<0.001) in patients who were obese and 6.4ng/ml lower for those whose ethnicity was not white (p<0.001).(74<sup>P</sup>) COVID-19 risk was increased for non-smokers,

London dwellers, males, and non-whites.(74<sup>P</sup>) After factoring out overweight and obesity (the factor with the highest odds ratio), and after grouping participant data into quartiles rather than using individual data, 25(OH)D did not independently predict COVID-19 risk.(74<sup>P</sup>)

Raisi-Estabragh, et al., conducted a third multivariate analysis on the UK Biobank participants, including all 4510 who had positive (1326) or negative (3184) COVID-19 tests from 16 March to 18 May 2020, almost all of whom were hospitalized.(75) The researchers used the baseline data from 10-14 years ago for age, sex, deprivation, BMI, and 25(OH)D levels, adjusting the 25(OH)D levels for seasonality and ethnicity.(75) Compared with the 497,996 untested participants of the UK Biobank study, men and non-white ethnicities were over-represented in the test group, with black ethnicity being 3.5 times more likely to be test positive than the untested cohort.(75) Men and whites had higher average 25(OH)D levels than women and non-white ethnicities.(75) Evaluating data from males and females independently, statistical significance was reached for males only for non-white ethnicity, more deprivation, and higher BMI.(75) For women, in addition to these three factors, lower 25(OH)D, more overcrowding, and greater risk-taking were all statistically significantly related to testing COVID-19 positive.(75) Rather than conducting a multivariate analysis on all potential influencers of COVID-19 positivity, Raisi-Estabragh, et al., grouped exposures, testing each group against sex, age, and ethnicity, finding no significant association between these three factors, seasonally and ethnically adjusted 25(OH)D levels, and positive COVID-19 status.(75) The researchers found that 25(OH)D and COVID-19 status are confounded by ethnicity and BMI.(75) Mean 25(OH)D levels for both COVID-19 negative (14.18ng/ml) and COVID-19 positive (13.55ng/ml) primarily hospitalized patients were extremely low.(75)

### **3.4.7 Rapid systematic review and meta-analysis with an ecological approach**

Ghasemian, et al., conducted a formal systematic review of nine studies, with six studies entering into a meta-analysis, and added their own evaluation of the correlation between global vitamin D status and COVID-19 recovery and mortality.(68<sup>P</sup>) The meta-analysis revealed that 46.5% of COVID-19 patients were vitamin D deficient and an additional 43.3% were vitamin D insufficient.(68<sup>P</sup>) Although their basic evaluation of 51 countries did not find a significant correlation between population vitamin D status and recovery or mortality rates, when latitude was factored in, both mortality rates and recovery rates weakly supported the vitamin D hypothesis.(68<sup>P</sup>) The researchers recommended large randomized clinical trials of vitamin D during the “Age of COVID-19.”(68<sup>P</sup>)

## **4 COVID-19-Specific Recommendations of Experts**

Although a few recommended only sunshine or 400IU/day, none of the authors strongly opposed vitamin D supplements during the pandemic. At the extremes, some researchers recommend large bolus doses of vitamin D, or correction of deficiency, primarily for patients who are diagnosed with COVID-19, and others recommended only the dose of vitamin D needed to maintain bone health (200-400IU/day).(28,29,71,89<sup>P</sup>,106,158,167,221,222<sup>P</sup>,223–228) Additional authors recommend vitamin D supplements to boost the immune systems of patients diagnosed with COVID-19.(2<sup>P</sup>,48,67,69,72,73,116,118,123,229–232) However, most authors recommend



Prior to modern times, individuals living in high latitudes had a much larger food supply from April – October, leading to weight gain.(251) Excess vitamin D from sunshine was stored in accumulated fat.(21<sup>P</sup>,204) Weight loss during relatively dark, food-scarce winters, released this excess vitamin D, preserving immune function.(21<sup>P</sup>,204) Now, food is plentiful year-round, leading to weight gain from decreased activity in winter.(251) Without weight-loss related vitamin D release, dangerously low 25(OH)D can develop by spring, and the obese, the elderly, those with naturally melanin-rich skin living outside the tropics, and anyone not spending time in the sun are at risk year-round.(204)

Sunscreen with a rating of only 15 SPF decreases vitamin D production in the skin by 99%.(204) Studies show that non-burning sun exposure increases vitamin D levels and may be melanoma-protective.(50) In tropical areas with wealthier populations, sun exposure may decrease in the summer due to a preference for air conditioning.(252) Encouraging uninfected people, including the homeless, to stay indoors could cause an increase in COVID-19 fatalities by increasing vitamin D deficiency rates. In contrast, encouraging weight loss through increased activity and structured programs can serve to improve vitamin D levels.(204) Studies show that exercise increases serum vitamin D levels, even when indoors, perhaps by triggering release of vitamin D stored in fat.(253<sup>P</sup>)

## 6 Conclusion

The 141 articles [**Table 3**] presenting primarily biological plausibility evidence overwhelmingly support the assertions that vitamin D sufficiency increases resistance to viral infections and helps prevent every symptom of severe COVID-19 that results in fatalities. They show that vitamin D deficiency can also explain every major risk factor, including the mystery of why children seem relatively protected and why males, the elderly, and people with naturally melanin-rich skin are especially vulnerable.

The 47 studies [**Table 2**] summarized here demonstrate that vitamin D deficiency explains the geographical differences in COVID-19 case and fatality rates. They provide overwhelming correlational evidence for the hypothesis, and causal evidence as well. COVID-19 mortality was predicted by vitamin D in sixteen studies(11<sup>P</sup>,21<sup>P</sup>,37,46<sup>P</sup>,61<sup>P</sup>,120,125,144<sup>P</sup>,148<sup>P</sup>,180<sup>P</sup>,195,204,207<sup>P</sup>,208,209,217<sup>P</sup>) and vitamin D levels or sunlight predicted contracting COVID-19 in seventeen.(45,120,122,148<sup>P</sup>,195,196,198<sup>P</sup>,199,202<sup>P</sup>,203<sup>P</sup>,205<sup>P</sup>,206,207<sup>P</sup>,209,214,219<sup>P</sup>,220<sup>P</sup>). Both causal modeling studies and eight chart reviews demonstrated that lower 25(OH)D was linearly associated with more severe COVID-19 outcomes.(40<sup>P</sup>,46<sup>P</sup>,47<sup>P</sup>,61<sup>P</sup>,155<sup>P</sup>,209,213,215<sup>P</sup>,217<sup>P</sup>,219<sup>P</sup>)

None of the four objections to recommending universal vitamin D supplements are supported by the evidence. The exhaustive literature search found no vitamin D proponent who suggested that COVID-19 could be completely eliminated with supplementation. Rather than overstating the case, they present compelling evidence that vitamin D deficiency is one factor which increases risk for COVID-19 infection and progression. Although overdoses are theoretically possible, they are highly improbable. The recommended dose by consensus, 2000IU/day for adults, is 1/20<sup>th</sup> the amount that must be taken for many months to risk toxicity.(38,86,89<sup>P</sup>,90) The

evidence strongly suggests that vitamin D deficiency is an easily modifiable risk factor and correcting it is potentially life-saving. Suppressing this evidence out of fear that the public might believe supplements will make them “immune” to COVID-19 is not only elitist, but it is inconsistent with existing public policy approaches. Many mitigation strategies are publicized. None are seen as conferring immunity.

This succinct but comprehensive review of the evidence found that despite almost complete absence of official government guidelines favoring vitamin D supplements to potentially decrease COVID-19 risk and severity, support among clinicians and other researchers for correcting and preventing vitamin D deficiency with modest daily vitamin D supplementation during the COVID-19 pandemic is very strong, worldwide. The evidence supports recommending 2000IU (50 mcg) vitamin D daily for at-risk teens and adults, which is well within safe limits and might dramatically reduce COVID-19 fatalities.

## **7 Limitations**

Many of the articles and studies included in this review were preprints, or were published in haste. The study descriptions were often too brief for a critical appraisal of the designs. Definitions of variables, such as race and ethnicity, were often omitted. Many researchers did not make their data public, although some emailed corrections or clarifications. Although the author is familiar with inflammation and cytokines from her work with chronic wounds, and she is familiar with epidemiology from her health education work in developing countries, she is not an endocrinologist or an epidemiologist. Single authorship could also be considered a limitation.

## **8 Afterward**

### **8.1 Conflict of Interest**

*The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.*

### **8.2 Author Contributions**

This article is the sole work of the author, Linda L Benskin.

### **8.3 Funding**

No special funding was provided for the writing of this article.

### **8.4 Acknowledgments**

The author wishes to thank her research associate, Richard Benskin, who provided invaluable editing help, John Newton whose encouragement led to this review, and her employer, Ferris Mfg. Corp., makers of PolyMem, who generously provided her with the time, as their charity liaison, to conduct the literature search and write this review.

### **8.5 Author Brief Biography**

Linda Benskin is an independent nurse researcher working to improve the evidence base for village health workers in remote and conflict areas of tropical developing countries, where health care professionals are absent. Her research into how pain and inflammation impact wound healing has provided her with a basic familiarity with cytokine pathophysiology. Dr. Benskin's improvised wound dressings clinical research study has been sidelined by the travel restrictions of COVID-19. Dr. Benskin is also the Clinical Research, Education, & Charity Liaison for Ferris Mfg. Corp. (makers of PolyMem dressings).

## 8.6 Abbreviations:

RCT = Randomized Controlled Trial

25(OH)D = serum vitamin D level

RAS = Renin-Angiotensin System

UV = ultra violet (light)

UVI = ultra violet light index

## 9 References

1. Wang L, Wang Y, Ye D, Liu Q. Review of the 2019 novel coronavirus (SARS-CoV-2) based on current evidence. *International Journal of Antimicrobial Agents* (2020)105948. doi:10.1016/j.ijantimicag.2020.105948
2. Bloukh SH, Shaikh AA, Pathan HM, Edis Z. Prevalence of COVID-19: A Look behind the Scenes. (2020) doi:10.20944/preprints202004.0179.v2 (Preprint)
3. Chinese CDC. The Epidemiological Characteristics of an Outbreak of 2019 Novel Coronavirus Diseases (COVID-19) — China, 2020. Available at: <http://weekly.chinacdc.cn/en/article/id/e53946e2-c6c4-41e9-9a9b-fea8db1a8f51> [Accessed April 23, 2020]
4. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet* (2020) **395**:497–506. doi:10.1016/S0140-6736(20)30183-5
5. Lighter J, Phillips M, Hochman S, Sterling S, Johnson D, Francois F, Stachel A. Obesity in patients younger than 60 years is a risk factor for Covid-19 hospital admission. *Clin Infect Dis* (2020) doi:10.1093/cid/ciaa415
6. Simonnet A, Chetboun M, Poissy J, Raverdy V, Noulette J, Duhamel A, Labreuche J, Mathieu D, Pattou F, Jourdain M, et al. High prevalence of obesity in severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) requiring invasive mechanical ventilation. *Obesity (Silver Spring)* (2020) doi:10.1002/oby.22831
7. Brown RA. Re: Preventing a covid-19 pandemic - COVID-19: Vitamin D deficiency; and, death rates; are both disproportionately higher in elderly Italians, Spanish, Swedish Somali, and African Americans? A connection? Research urgently required! *The BMJ* (2020) Available at: <https://www.bmj.com/content/368/bmj.m810/rr-46> [Accessed April 23, 2020]



8. Docherty AB, Harrison EM, Green CA, Hardwick HE, Pius R, Norman L, Holden KA, Read JM, Dondelinger F, Carson G, et al. Features of 10 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: prospective observational cohort study. *medRxiv* (2020)2020.04.23.20076042. doi:10.1101/2020.04.23.20076042
9. Ebadi M, Montano-Loza AJ. Perspective: improving vitamin D status in the management of COVID-19. *Eur J Clin Nutr* (2020) doi:10.1038/s41430-020-0661-0
10. Hastie CE, Mackay DF, Ho F, Celis-Morales CA, Katikireddi SV, Niedzwiedz CL, Jani BD, Welsh P, Mair FS, Gray SR, et al. Vitamin D concentrations and COVID-19 infection in UK Biobank. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews* (2020) **14**:561–565. doi:10.1016/j.dsx.2020.04.050
11. Bäcker A. Why COVID-19 May Be Disproportionately Killing African Americans: Black Overrepresentation among COVID-19 Mortality Increases with Lower Irradiance, Where Ethnicity Is More Predictive of COVID-19 Infection and Mortality Than Median Income. Social Science Research Network (2020). doi:10.2139/ssrn.3571699 (Preprint)
12. Kass DA, Duggal P, Cingolani O. Obesity could shift severe COVID-19 disease to younger ages. *Lancet* (2020) **395**:1544–1545. doi:10.1016/S0140-6736(20)31024-2
13. Cook T, Kursumovic E, Lennane S. Exclusive: deaths of NHS staff from covid-19 analysed. *Health Service Journal* Available at: <https://www.hsj.co.uk/exclusive-deaths-of-nhs-staff-from-covid-19-analysed/7027471.article> [Accessed June 24, 2020]
14. Gudbjartsson DF, Helgason A, Jonsson H, Magnusson OT, Melsted P, Norddahl GL, Saemundsdottir J, Sigurdsson A, Sulem P, Agustsdottir AB, et al. Spread of SARS-CoV-2 in the Icelandic Population. *New England Journal of Medicine* (2020) **0**:null. doi:10.1056/NEJMoa2006100
15. National Centre for Immunisation Research and Surveillance. COVID-19 in schools – the experience in NSW | NCIRS. National Centre for Immunisation Research and Surveillance (2020). Available at: <http://www.ncirs.org.au/covid-19-in-schools> [Accessed May 21, 2020]
16. Grant WB, Lahore H, McDonnell SL, Baggerly CA, French CB, Aliano JL, Bhattoa HP. Evidence that Vitamin D Supplementation Could Reduce Risk of Influenza and COVID-19 Infections and Deaths. *Nutrients* (2020) **12**:988. doi:10.3390/nu12040988
17. Conticini E, Frediani B, Caro D. Can atmospheric pollution be considered a co-factor in extremely high level of SARS-CoV-2 lethality in Northern Italy? *Environ Pollut* (2020)114465. doi:10.1016/j.envpol.2020.114465
18. Martelletti L, Martelletti P. Air Pollution and the Novel Covid-19 Disease: a Putative Disease Risk Factor. *SN Compr Clin Med* (2020)1–5. doi:10.1007/s42399-020-00274-4
19. Zhu Y, Xie J, Huang F, Cao L. Association between short-term exposure to air pollution and COVID-19 infection: Evidence from China. *Sci Total Environ* (2020) **727**:138704. doi:10.1016/j.scitotenv.2020.138704

20. Carleton T, Meng KC. Causal empirical estimates suggest COVID-19 transmission rates are highly seasonal. *medRxiv* (2020)2020.03.26.20044420. doi:10.1101/2020.03.26.20044420 (Preprint)
21. Moozhipurath RK, Kraft L, Skiera B. Evidence of Protective Role of Ultraviolet-B (UVB) Radiation in Reducing COVID-19 Deaths. *medRxiv* (2020)2020.05.06.20093419. doi:10.1101/2020.05.06.20093419 (Preprint)
22. Cao Z, Wu Y, Faucon E, Sabatier J-M. SARS-CoV-2 & Covid-19: Key-Roles of the “Renin-Angiotensin” System / Vitamin D Impacting Drug and Vaccine Developments. *Infect Disord Drug Targets* (2020) doi:10.2174/1871526520999200505174704
23. Rhodes JM, Subramanian S, Laird E, Anne Kenny R. Editorial: low population mortality from COVID-19 in countries south of latitude 35 degrees North - supports vitamin D as a factor determining severity. *Aliment Pharmacol Ther* (2020) doi:10.1111/apt.15777
24. Perry T. Tyler Perry Encouraging Blacks To Increase Vitamin D During COVID-19. *Black Enterprise* (2020) Available at: <https://www.blackenterprise.com/tyler-perry-is-encouraging-black-people-to-increase-their-vitamin-d-during-the-covid-19-pandemic/> [Accessed May 4, 2020]
25. Illmatical C. African Americans High COVID-19 Mortality Rates: Is This Due To A Lack Of Vitamin D? | Afro. Available at: <https://www.afro.com/african-americans-high-covid-19-mortality-rates-is-this-due-to-a-lack-of-vitamin-d/> [Accessed May 4, 2020]
26. Pinnock D. Vitamin D and COVID-19. The Evidence Warrants Discussion! *The Medicinal Chef making health simple* (2020) Available at: <https://www.themedicinalchef.co.uk/vitamin-d-and-covid-19-the-evidence-warrants-discussion/> [Accessed June 25, 2020]
27. Trinity College Dublin. Vitamin D determines severity in COVID-19 so government advice needs to change. *EurekaAlert!* Available at: [https://www.eurekaalert.org/pub\\_releases/2020-05/tcd-vdd051220.php](https://www.eurekaalert.org/pub_releases/2020-05/tcd-vdd051220.php) [Accessed May 14, 2020]
28. Garg M, Al-Ani A, Mitchell H, Hendy P, Christensen B. Editorial: low population mortality from COVID-19 in countries south of latitude 35 degrees North - supports vitamin D as a factor determining severity. Authors’ reply. *Aliment Pharmacol Ther* (2020) doi:10.1111/apt.15796
29. Lee J, van Hecke O, Roberts N. Vitamin D: A rapid review of the evidence for treatment or prevention in COVID-19. *CEBM* Available at: <https://www.cebm.net/covid-19/vitamin-d-a-rapid-review-of-the-evidence-for-treatment-or-prevention-in-covid-19/> [Accessed May 4, 2020]
30. Should people take vitamin D to ward off the new coronavirus? Available at: <https://www.medicalnewstoday.com/articles/should-people-take-vitamin-d-to-ward-off-the-new-coronavirus> [Accessed May 8, 2020]
31. McRae M. COVID-19 Deaths Are Being Linked to Vitamin D Deficiency. Here’s What That Means. *ScienceAlert* Available at: <https://www.sciencealert.com/covid-deaths-are-being-linked-with-vitamin-d-deficiency-here-s-what-that-means> [Accessed May 8, 2020]

32. Lee BY. Can Vitamin D Help With Covid-19 Coronavirus? Here Is The Science. *Forbes* Available at: <https://www.forbes.com/sites/brucelee/2020/05/17/can-vitamin-d-help-with-covid-19-coronavirus-here-is-the-science/> [Accessed May 19, 2020]
33. Can Vitamin D Prevent COVID-19? *Health Essentials from Cleveland Clinic* (2020) Available at: <https://health.clevelandclinic.org/can-vitamin-d-prevent-covid-19/> [Accessed May 19, 2020]
34. Barron A, Patel N. Vitamin D: A rapid review of the evidence for treatment or prevention of COVID-19 in adults (excluding pregnant & breastfeeding women). London, UK: NIH (2020). Available at: <https://www.ncl-mon.nhs.uk/faq/guidelines/>
35. Auwaerter PG. UPDATE 5/20/2020 - COVID-19: Keeping Up with a Moving Target. Johns Hopkins: DKBMed (2020). Available at: <https://www.youtube.com/watch?v=LaAQ8rX4x9g> [Accessed May 24, 2020]
36. Wan DM. In Conversation – Nobel Laureate Prof. Peter Doherty. *SR Vol57(06) [June 2020]* (2020) Available at: <http://nopr.niscair.res.in/handle/123456789/54393> [Accessed May 28, 2020]
37. Laird E, Rhodes J, Kenny RA. Vitamin D and Inflammation – Potential Implications for Severity of Covid-19 – *Irish Medical Journal*. *Irish Medical Journal* (2020) **113**:81. Available at: <http://imj.ie/vitamin-d-and-inflammation-potential-implications-for-severity-of-covid-19/> [Accessed May 14, 2020]
38. Hribar CA, Cobbold PH, Church FC. Potential Role of Vitamin D in the Elderly to Resist COVID-19 and to Slow Progression of Parkinson’s Disease. *Brain Sci* (2020) **10**: doi:10.3390/brainsci10050284
39. Holick MF. VITAMIN D STATUS: MEASUREMENT, INTERPRETATION AND CLINICAL APPLICATION. *Ann Epidemiol* (2009) **19**:73–78. doi:10.1016/j.annepidem.2007.12.001
40. Alipio M. Vitamin D Supplementation Could Possibly Improve Clinical Outcomes of Patients Infected with Coronavirus-2019 (COVID-2019). Rochester, NY: Social Science Research Network (2020). doi:10.2139/ssrn.3571484 (Preprint)
41. Fabbri A, Infante M, Ricordi C. Editorial - Vitamin D status: a key modulator of innate immunity and natural defense from acute viral respiratory infections. *Eur Rev Med Pharmacol Sci* (2020) **24**:4048–4052. doi:10.26355/eurrev\_202004\_20876
42. Boucher BJ. The Problems of Vitamin D Insufficiency in Older People. *Aging Dis* (2012) **3**:313–329. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3501367/> [Accessed April 23, 2020]
43. Crane-Godreau MA, Clem KJ, Payne P, Fiering S. Vitamin D Deficiency and Air Pollution Exacerbate COVID-19 Through Suppression of Antiviral Peptide LL37. *Front Public Health* (2020) **8**: doi:10.3389/fpubh.2020.00232
44. Ghavideldarestani M, Honardoost M, Khamseh ME. Role of Vitamin D in Pathogenesis and Severity of COVID-19 Infection. (2020) doi:10.20944/preprints202004.0355.v1 (Preprint)

45. Kara M, Ekiz T, Ricci V, Kara Ö, Chang K-V, Özçakar L. ‘Scientific Strabismus’ or two related pandemics: coronavirus disease and vitamin D deficiency. *Br J Nutr* 1–6. doi:10.1017/S0007114520001749
46. Raharusun P, Priambada S, Budiarti C, Agung E, Budi C. Patterns of COVID-19 Mortality and Vitamin D: An Indonesian Study. *Emerginnova* (2020). Available at: <https://emergingnova.com/patterns-of-covid19-mortality-and-vitamin-d-an-indonesian-study/> [Accessed May 20, 2020] (Preprint)
47. Glicio EJ. Vitamin D Level of Mild and Severe Elderly Cases of COVID-19: A Preliminary Report. Rochester, NY: Social Science Research Network (2020). Available at: <https://papers.ssrn.com/abstract=3593258> [Accessed May 20, 2020] (Preprint)
48. Caccialanza R, Laviano A, Lobascio F, Montagna E, Bruno R, Ludovisi S, Corsico AG, Di Sabatino A, Belliato M, Calvi M, et al. Early nutritional supplementation in non-critically ill patients hospitalized for the 2019 novel coronavirus disease (COVID-19): Rationale and feasibility of a shared pragmatic protocol. *Nutrition* (2020)110835. doi:10.1016/j.nut.2020.110835
49. Calder PC, Carr AC, Gombart AF, Eggersdorfer M. Optimal Nutritional Status for a Well-Functioning Immune System Is an Important Factor to Protect against Viral Infections. *Nutrients* (2020) **12**: doi:10.3390/nu12041181
50. Hoel DG, Berwick M, Grujil FR de, Holick MF. The risks and benefits of sun exposure 2016. *Dermato-Endocrinology* (2016) **8**:e1248325. doi:10.1080/19381980.2016.1248325
51. La Vignera S, Cannarella R, Condorelli RA, Torre F, Aversa A, Calogero AE. Sex-Specific SARS-CoV-2 Mortality: Among Hormone-Modulated ACE2 Expression, Risk of Venous Thromboembolism and Hypovitaminosis D. *Int J Mol Sci* (2020) **21**: doi:10.3390/ijms21082948
52. Chhetri JK, Chan P, Arai H, Chul Park S, Sriyani Gunaratne P, Setiati S, Assantachai P. Prevention of COVID-19 in Older Adults: A Brief Guidance from the International Association for Gerontology and Geriatrics (IAGG) Asia/Oceania region. *J Nutr Health Aging* (2020) **24**:471–472. doi:10.1007/s12603-020-1359-7
53. Raymond M, Ching-A-Sue G, Oliver VH. Mast cell stabilisers, leukotriene antagonists and antihistamines: A rapid review of effectiveness in COVID-19. *Centre for Evidence-Based Medicine* (2020) Available at: <https://covid19-evidence.paho.org/handle/20.500.12663/1569> [Accessed June 16, 2020]
54. Isaia G, Medico E. Possible prevention and therapeutic role of vitamin D in the management of the COVID-19 2020 pandemics. (2020) Available at: <https://minapim.com/en/possible-preventive-and-therapeutic-role-of-vitamin-d-in-the-management-of-the-covid-19-pandemic/>
55. Ribeiro H, Santana KV de S de, Oliver SL, Rondó PH de C, Mendes MM, Charlton K, Lanham-New S. Does Vitamin D play a role in the management of Covid-19 in Brazil? *Revista de Saúde Pública* **54**: doi:10.11606/s1518-8787.2020054002545

56. Nota de Esclarecimento: Vitamina D e Covid-19. *Sociedade Brasileira de Endocrinologia e Metabologia* Available at: <https://www.endocrino.org.br/nota-de-esclarecimento-vitamina-d-e-covid-19/> [Accessed June 26, 2020]
57. Thebault R, Tran AB, Williams V. The coronavirus is infecting and killing black Americans at an alarmingly high rate. *Washington Post* (2020) Available at: <https://www.washingtonpost.com/nation/2020/04/07/coronavirus-is-infecting-killing-black-americans-an-alarmingly-high-rate-post-analysis-shows/> [Accessed May 7, 2020]
58. Eligon J, Burch ADS, Searcey D, Jr RAO. Black Americans Face Alarming Rates of Coronavirus Infection in Some States. *The New York Times* (2020) Available at: <https://www.nytimes.com/2020/04/07/us/coronavirus-race.html> [Accessed June 28, 2020]
59. CDC. COVID-19 in Racial and Ethnic Minority Groups. *Centers for Disease Control and Prevention* (2020) Available at: <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/racial-ethnic-minorities.html> [Accessed June 28, 2020]
60. Wadhera RK, Wadhera P, Gaba P, Figueroa JF, Maddox KEJ, Yeh RW, Shen C. Variation in COVID-19 Hospitalizations and Deaths Across New York City Boroughs. *JAMA* (2020) **323**:2192–2195. doi:10.1001/jama.2020.7197
61. Davies G, Garami AR, Byers JC. Evidence Supports a Causal Model for Vitamin D in COVID-19 Outcomes. *medRxiv* (2020)2020.05.01.20087965. doi:10.1101/2020.05.01.20087965 (Preprint)
62. Tom Frieden MD. Former CDC Chief Dr. Tom Frieden: Coronavirus infection risk may be reduced by Vitamin D. *Fox News* (2020) Available at: <https://www.foxnews.com/opinion/former-cdc-chief-tom-frieden-coronavirus-risk-may-be-reduced-with-vitamin-d> [Accessed May 8, 2020]
63. Manson JE. Does Vitamin D Protect Against COVID-19? *Medscape* (2020) Available at: <http://www.medscape.com/viewarticle/930152> [Accessed May 12, 2020]
64. Doheny K. More Vitamin D, Lower Risk of Severe COVID-19? *WebMD* Available at: <https://www.webmd.com/lung/news/20200518/more-vitamin-d-lower-risk-of-severe-covid-19> [Accessed May 19, 2020]
65. British Dietetic Association. COVID-19 / Coronavirus - Advice for the General Public. (2020) Available at: <https://www.bda.uk.com/resource/covid-19-corona-virus-advice-for-the-general-public.html> [Accessed May 19, 2020]
66. McCall B. Vitamin D: A Low-Hanging Fruit in COVID-19? *Medscape* Available at: <http://www.medscape.com/viewarticle/930660> [Accessed June 15, 2020]
67. Mitchell F. Vitamin-D and COVID-19: do deficient risk a poorer outcome? *The Lancet Diabetes & Endocrinology* (2020) **0**: doi:10.1016/S2213-8587(20)30183-2
68. Ghasemian R, Shamshirian A, Heydari K, Malekan M, Alizadeh-Navaei R, Ebrahimzadeh MA, Jafarpour H, Shahmirzadi AR, Khodabandeh M, Seyfari B. The Role of Vitamin D in The Age of COVID-19: A Systematic Review and Meta-Analysis Along with an Ecological Approach. *medRxiv* (2020) (Preprint)

69. Taheri S, Chagoury O, Tourette M, Skaroni I, Othman M, Bashir M, Dukhan K, Mohannadi DA, Hassan D, Ghadban W, et al. Managing diabetes in Qatar during the COVID-19 pandemic. *The Lancet Diabetes & Endocrinology* (2020) **0**: doi:10.1016/S2213-8587(20)30154-6
70. Laird E, Kenny RA. Vitamin D deficiency in Ireland – implications for COVID-19. Results from the Irish Longitudinal Study on Ageing (TILDA). *The Irish Longitudinal Study on Ageing* (2020). doi:10.38018/TildaRe.2020-05
71. McKenna MJ, Flynn MAT. Covid-19 Cocooning and Vitamin D Intake Requirements – Irish Medical Journal. *Irish Medical Journal* (2020) **113**:79. Available at: <http://imj.ie/covid-19-cocooning-and-vitamin-d-intake-requirements/> [Accessed May 14, 2020]
72. Molloy EJ, Murphy N. Vitamin D, Covid-19 and Children. *Ir Med J* (2020) **113**:64.
73. McCartney DM, Byrne DG. Optimisation of Vitamin D Status for Enhanced Immuno-protection Against Covid-19. *Ir Med J* (2020) **113**:58.
74. Darling AL, Ahmadi KR, Ward KA, Harvey NC, Alves AC, Dunn-Waters DK, Lanham-New SA, Cooper C, Blackbourn DJ. Vitamin D status, body mass index, ethnicity and COVID-19: Initial analysis of the first-reported UK Biobank COVID-19 positive cases (n 580) compared with negative controls (n 723). *medRxiv* (2020)2020.04.29.20084277. doi:10.1101/2020.04.29.20084277 (Preprint)
75. Raisi-Estabragh Z, McCracken C, Bethell MS, Cooper J, Cooper C, Caulfield MJ, Munroe PB, Harvey NC, Petersen SE. Greater risk of severe COVID-19 in Black, Asian and Minority Ethnic populations is not explained by cardiometabolic, socioeconomic or behavioural factors, or by 25(OH)-vitamin D status: study of 1326 cases from the UK Biobank. *J of Public Health (Oxf)* (2020) <https://academic.oup.com/jpubhealth/article/doi/10.1093/pubmed/fdaa095/5859581>
76. Meng JE, Hovey KM, Wactawski-Wende J, Andrews CA, LaMonte MJ, Horst RL, Genco RJ, Millen AE. Intraindividual Variation in Plasma 25-Hydroxyvitamin D Measures 5 Years Apart among Postmenopausal Women. *Cancer Epidemiol Biomarkers Prev* (2012) **21**:916–924. doi:10.1158/1055-9965.EPI-12-0026
77. Roy AS, Matson M, Herlekar R. Response to ‘Vitamin D concentrations and COVID-19 infection in UK Biobank.’ *Diabetes & Metabolic Syndrome: Clinical Research & Reviews* (2020) **14**:777. doi:10.1016/j.dsx.2020.05.049
78. Etsy E. The Possible Link Between Severe COVID-19 and Low Vitamin D Levels. Available at: [https://www.youtube.com/watch?v=\\_ZJkDvVFbRE](https://www.youtube.com/watch?v=_ZJkDvVFbRE) [Accessed June 22, 2020]
79. Jorde R, Sneve M, Hutchinson M, Emaus N, Figenschau Y, Grimnes G. Tracking of Serum 25-Hydroxyvitamin D Levels During 14 Years in a Population-based Study and During 12 Months in an Intervention Study. *Am J Epidemiol* (2010) **171**:903–908. doi:10.1093/aje/kwq005
80. Yong SJ. Population Studies Confirm Risk Factors for Catching Covid-19. *Medium* (2020) Available at: <https://medium.com/microbial-instincts/population-studies-confirm-risk-factors-for-catching-covid-19-853779e7a3c6> [Accessed June 21, 2020]

81. Boucher BJ. Adjustments in analyses of vitamin D status, allowing for vitamin D determinants, for Covid-19 risks. *Diabetes Metab Res Rev* (2020)e3375. doi:10.1002/dmrr.3375
82. Grant WB, McDonnell SL. Letter in response to the article: Vitamin D concentrations and COVID-19 infection in UK biobank (Hastie et al.). *Diabetes & Metabolic Syndrome: Clinical Research & Reviews* (2020) **14**:893–894. doi:10.1016/j.dsx.2020.05.046
83. Fox B, Sizemore JO. No Association Seen Between Vitamin D Deficiency and COVID-19 Infection, Hospitalization, or Mortality.
84. DeFilipps K. Comment on No Association Seen Between Vitamin D Deficiency and COVID-19 Infection, Hospitalization, or Mortality. *Epic Health Research Network* (2020) Available at: <http://ehrn.org/no-association-seen-between-vitamin-d-deficiency-and-covid-19-infection-hospitalization-or-mortality/> [Accessed June 23, 2020]
85. Christakos S, Li S, De La Cruz J, Bikle DD. New developments in our understanding of vitamin D metabolism, action and treatment. *Metabolism* (2019) **98**:112–120. doi:10.1016/j.metabol.2019.06.010
86. Marcinowska-Suchowierska E, Kupisz-Urbańska M, Łukasziewicz J, Płudowski P, Jones G. Vitamin D Toxicity—A Clinical Perspective. *Front Endocrinol* (2018) **9**: doi:10.3389/fendo.2018.00550
87. Solis M. A cloud of doubt over the sunshine vitamin. *Pharmaceutical Journal* (2020) **304**: doi:DOI: 10.1211/PJ.2020.20207521
88. Buttriss JL, Lanham-New SA. Is a vitamin D fortification strategy needed? *Nutr Bull* (2020) **45**:115–122. doi:10.1111/nbu.12430
89. Bakare TA, Soar JS. A cost-effective preventative approach to potentially save lives in the coronavirus pandemic, jointly using Vitamin D, Curcumin, and Vitamin C. IndiaRxiv Preprints (2020). Available at: <https://indiarxiv.org/cq8sa/> [Accessed May 8, 2020] (Preprint)
90. Meehan M, Penckofer S. The Role of Vitamin D in the Aging Adult. *J Aging Gerontol* (2014) **2**:60–71. doi:10.12974/2309-6128.2014.02.02.1
91. Kimball SM, Mirhosseini N, Holick MF. Evaluation of vitamin D3 intakes up to 15,000 international units/day and serum 25-hydroxyvitamin D concentrations up to 300 nmol/L on calcium metabolism in a community setting. *Dermatoendocrinol* (2017) **9**: doi:10.1080/19381980.2017.1300213
92. Lim K, Thadhani R, Lim K, Thadhani R. Vitamin D Toxicity. *Brazilian Journal of Nephrology* (2020) doi:10.1590/2175-8239-jbn-2019-0192
93. Spiller HA, Good TF, Spiller NE, Aleguas A. Vitamin D exposures reported to US poison centers 2000-2014: Temporal trends and outcomes. *Hum Exp Toxicol* (2016) **35**:457–461. doi:10.1177/0960327115595685
94. Veugelers P, Ekwaru J. A Statistical Error in the Estimation of the Recommended Dietary Allowance for Vitamin D. *Nutrients* (2014) **6**:4472–4475. doi:10.3390/nu6104472

95. Heaney R, Garland C, Baggerly C, French C, Gorham E. Letter to Veugelers, P.J. and Ekwaru, J.P., A statistical error in the estimation of the recommended dietary allowance for vitamin D. *Nutrients* 2014, 6, 4472-4475; doi:10.3390/nu6104472. *Nutrients* (2015) **7**:1688–1690. doi:10.3390/nu7031688
96. Burt LA, Billington EO, Rose MS, Raymond DA, Hanley DA, Boyd SK. Effect of High-Dose Vitamin D Supplementation on Volumetric Bone Density and Bone Strength: A Randomized Clinical Trial. *JAMA* (2019) **322**:736–745. doi:10.1001/jama.2019.11889
97. McCullough PJ, Amend J, McCullough WP, Repas SJ, Travers JB, Lehrer DS. The Essential Role of Vitamin D in the Biosynthesis of Endogenous Antimicrobial Peptides May Explain Why Deficiency Increases Mortality Risk in COVID-19 Infections. (2020) doi:10.20944/preprints202005.0265.v1 (Preprint)
98. Singer P, Blaser AR, Berger MM, Alhazzani W, Calder PC, Casaer MP, Hiesmayr M, Mayer K, Montejo JC, Pichard C, et al. ESPEN guideline on clinical nutrition in the intensive care unit. *Clinical Nutrition* (2019) **38**:48–79. doi:10.1016/j.clnu.2018.08.037
99. Early High-Dose Vitamin D3 for Critically Ill, Vitamin D–Deficient Patients. *New England Journal of Medicine* (2019) **381**:2529–2540. doi:10.1056/NEJMoa1911124
100. Han JE, Jones JL, Tangpricha V, Brown MA, Hao L, Hebbar G, Lee MJ, Liu S, Brown LAS, Ziegler TR, et al. High dose vitamin D administration in ventilated intensive care unit patients: A pilot double blind randomized controlled trial. *J Clin Transl Endocrinol* (2016) **4**:59–65. doi:10.1016/j.jcte.2016.04.004
101. Kow CS, Hadi MA, Hasan SS. Vitamin D Supplementation in Influenza and COVID-19 Infections Comment on: “Evidence that Vitamin D Supplementation Could Reduce Risk of Influenza and COVID-19 Infections and Deaths” *Nutrients* 2020, 12(4), 988. *Nutrients* (2020) **12**: doi:10.3390/nu12061626
102. Martineau AR, Jolliffe DA, Hooper RL, Greenberg L, Aloia JF, Bergman P, Dubnov-Raz G, Esposito S, Ganmaa D, Ginde AA, et al. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *BMJ* (2017) **356**:i6583. doi:10.1136/bmj.i6583
103. Grant WB, Baggerly CA, Lahore H. Reply: “Vitamin D Supplementation in Influenza and COVID-19 Infections. Comment on: Evidence That Vitamin D Supplementation Could Reduce Risk of Influenza and COVID-19 Infections and Deaths *Nutrients* 2020, 12(4), 988.” *Nutrients* (2020) **12**:1620. doi:10.3390/nu12061620
104. Aloia JF, Li-Ng M. Re: epidemic influenza and vitamin D. *Epidemiol Infect* (2007) **135**:1095–1096; author reply 1097-1098. doi:10.1017/S0950268807008308
105. Sharma SK, Mudgal SK, Pai VS, Chaturvedi J, Gaur R. Vitamin D: A cheap yet effective bullet against coronavirus disease-19—Are we convinced yet? *National Journal of Physiology, Pharmacy and Pharmacology* (2020) **10**:0–0.



106. Quesada-Gomez JM, Castillo ME, Bouillon R. Vitamin D Receptor stimulation to reduce Acute Respiratory Distress Syndrome (ARDS) in patients with Coronavirus SARS-CoV-2 infections: Revised Ms SBMB 2020\_166. *J Steroid Biochem Mol Biol* (2020)105719. doi:10.1016/j.jsbmb.2020.105719
107. Chakhtoura M, Napoli N, El Hajj Fuleihan G. Commentary: Myths and facts on vitamin D amidst the COVID-19 pandemic. *Metabolism* (2020) **109**:154276. doi:10.1016/j.metabol.2020.154276
108. Kakodkar P, Kaka N, Baig MN. A Comprehensive Literature Review on the Clinical Presentation, and Management of the Pandemic Coronavirus Disease 2019 (COVID-19). *Cureus* (2020) **12**:e7560. doi:10.7759/cureus.7560
109. Bokharee N, Khan YH, Wasim T, Mallhi TH, Alotaibi NH, Iqbal MS, Rehman K, Alzarea AI, Khokhar A. Daily versus stat vitamin D supplementation during pregnancy; A prospective cohort study. *PLoS One* (2020) **15**: doi:10.1371/journal.pone.0231590
110. Brown RA, Sarkar A. Vitamin D deficiency: a factor in COVID-19, progression, severity and mortality? – An urgent call for research. *MitoFit Preprint Arch* (2020) Available at: [https://www.mitofit.org/index.php/Brown\\_2020\\_MitoFit\\_Preprint\\_Arch](https://www.mitofit.org/index.php/Brown_2020_MitoFit_Preprint_Arch) [Accessed April 23, 2020] (Preprint)
111. Carter SJ, Baranauskas MN, Fly AD. Considerations for obesity, vitamin D, and physical activity amidst the COVID-19 pandemic. *Obesity (Silver Spring)* (2020) doi:10.1002/oby.22838
112. Tian Y, Rong L. Letter: Covid-19 and vitamin D-authors' reply. *Aliment Pharmacol Ther* (2020) doi:10.1111/apt.15764
113. Jakovac H. COVID-19 and vitamin D—Is there a link and an opportunity for intervention? *American Journal of Physiology-Endocrinology and Metabolism* (2020) **318**:E589–E589. doi:10.1152/ajpendo.00138.2020
114. Misra DP, Agarwal V, Gasparyan AY, Zimba O. Rheumatologists' perspective on coronavirus disease 19 (COVID-19) and potential therapeutic targets. *Clin Rheumatol* (2020) doi:10.1007/s10067-020-05073-9
115. Turashvili N, Javashvili L. Could vitamin D reduce the risk of COVID-19? *Translational and Clinical Medicine - Georgian Medical Journal* (2020) **5**:4–6. Available at: <http://tcm.tsu.ge/index.php/TCM-GMJ/article/view/204> [Accessed May 8, 2020]
116. Parvin F, Islam S, Urmay Z, Ahmed S. THE SYMPTOMS, CONTAGIOUS PROCESS, PREVENTION AND POST TREATMENT OF COVID-19. *European Journal of Physiotherapy and Rehabilitation Studies* (2020) **0**: Available at: <https://www.oapub.org/hlt/index.php/EJPRS/article/view/52> [Accessed May 7, 2020]
117. Islam MR, Fischer A. A Transcriptome Analysis Identifies Potential Preventive and Therapeutic Approaches Towards COVID-19. (2020) doi:10.20944/preprints202004.0399.v1 (Preprint)
118. Zhang L, Liu Y. Potential interventions for novel coronavirus in China: A systematic review. *J Med Virol* (2020) **92**:479–490. doi:10.1002/jmv.25707

119. Jayawardena R, Sooriyaarachchi P, Chourdakis M, Jeewandara C, Ranasinghe P. Enhancing immunity in viral infections, with special emphasis on COVID-19: A review. *Diabetes Metab Syndr* (2020) doi:10.1016/j.dsx.2020.04.015
120. Ilie PC, Stefanescu S, Smith L. The role of vitamin D in the prevention of coronavirus disease 2019 infection and mortality. *Aging Clin Exp Res* (2020) doi:10.1007/s40520-020-01570-8
121. Garami AR. Re: Preventing a covid-19 pandemic - Is there a magic bullet to save COVID-19 patients? We can give it a try! *BMJ (British Medical Journal)* (2020) **2020;368:m810**: Available at: <https://www.bmj.com/content/368/bmj.m810/rr-24> [Accessed May 10, 2020]
122. D'Avolio A, Avataneo V, Manca A, Cusato J, De Nicolò A, Lucchini R, Keller F, Cantù M. 25-Hydroxyvitamin D Concentrations Are Lower in Patients with Positive PCR for SARS-CoV-2. *Nutrients* (2020) **12**: doi:10.3390/nu12051359
123. Adams KK, Baker WL, Sobieraj DM. Myth Busters: Dietary Supplements and COVID-19. *Ann Pharmacother* (2020)1060028020928052. doi:10.1177/1060028020928052
124. Mansur J. Letter: low population mortality from COVID-19 in countries south of latitude 35 degrees North supports vitamin D as a factor determining severity. *Aliment Pharmacol Ther* (2020) doi:10.1111/apt.15820
125. Rhodes JM, Subramanian S, Laird E, Anne Kenny R. Letter: low population mortality from COVID-19 in countries south of latitude 35 degrees North supports vitamin D as a factor determining severity. Authors' reply. *Aliment Pharmacol Ther* (2020) doi:10.1111/apt.15823
126. Zhang J, Xie B, Hashimoto K. Current status of potential therapeutic candidates for the COVID-19 crisis. *Brain Behav Immun* (2020) doi:10.1016/j.bbi.2020.04.046
127. DeRobbio D. MUSC vitamin D research team to study connection to COVID-19 prevention and treatment. *WCIV* (2020) Available at: <https://abcnews4.com/news/coronavirus/musc-vitamin-d-research-team-to-study-connection-to-covid-19-prevention-and-treatment> [Accessed May 4, 2020]
128. Reporter S. Countries with Low Vitamin D Have Higher COVID-19 Infection and Mortality Rates: Here are They. *Science Times* (2020) Available at: <https://www.sciencetimes.com/articles/25567/20200504/coronavirus-countries-low-vitamin-d-high-risk-getting-disease.htm> [Accessed May 4, 2020]
129. Louisiana State University Health Sciences Center in New Orleans. The LEAD COVID-19 Trial: Low-risk, Early Aspirin and Vitamin D to Reduce COVID-19 Hospitalizations. *clinicaltrials.gov* (2020). Available at: <https://clinicaltrials.gov/ct2/show/NCT04363840> [Accessed May 18, 2020]
130. Tameside General Hospital. Investigating the Role of Vitamin D in the Morbidity of COVID-19 Patients. *clinicaltrials.gov* (2020). Available at: <https://clinicaltrials.gov/ct2/show/NCT04386044> [Accessed May 18, 2020]

131. VITACOV: Vitamin D Polymorphisms and Severity of COVID-19 Infection - Full Text View - ClinicalTrials.gov. Available at: <https://clinicaltrials.gov/ct2/show/NCT04370808> [Accessed May 19, 2020]
132. Montano-Loza A. Improving Vitamin D Status in the Management of COVID-19. clinicaltrials.gov (2020). Available at: <https://clinicaltrials.gov/ct2/show/NCT04385940> [Accessed May 18, 2020]
133. Garzón MC. Effect of Vitamin D Administration on Prevention and Treatment of Mild Forms of Suspected Covid-19. clinicaltrials.gov (2020). Available at: <https://clinicaltrials.gov/ct2/show/NCT04334005> [Accessed May 18, 2020]
134. Dandinoğlu T. Do Vitamin D Levels Really Correlated With Disease Severity in COVID-19 Patients? clinicaltrials.gov (2020). Available at: <https://clinicaltrials.gov/ct2/show/NCT04394390> [Accessed May 18, 2020]
135. COvid-19 and Vitamin D Supplementation: a Multicenter Randomized Controlled Trial of High Dose Versus Standard Dose Vitamin D3 in High-risk COVID-19 Patients (CoVitTrial) - Full Text View - ClinicalTrials.gov. Available at: <https://clinicaltrials.gov/ct2/show/NCT04344041> [Accessed May 19, 2020]
136. University Hospital, Lille. Impact of Zinc and Vitamin D3 Supplementation on the Survival of Institutionalized Aged Patients Infected With COVID-19. clinicaltrials.gov (2020). Available at: <https://clinicaltrials.gov/ct2/show/NCT04351490> [Accessed May 18, 2020]
137. Sanford Health. Randomized, Double-Blind, Controlled Trial of Hydroxychloroquine vs Placebo as Post-Exposure Prophylaxis Against COVID-19 Infection. clinicaltrials.gov (2020). Available at: <https://clinicaltrials.gov/ct2/show/NCT04372017> [Accessed May 18, 2020]
138. ProgenaBiome. A Randomized, Double-Blind, Placebo-Controlled Phase IIa Study of Quintuple Therapy to Treat COVID-19 Infection. clinicaltrials.gov (2020). Available at: <https://clinicaltrials.gov/ct2/show/NCT04334512> [Accessed May 18, 2020]
139. Oral 25-hydroxyvitamin D3 and COVID-19 - Full Text View - ClinicalTrials.gov. Available at: <https://clinicaltrials.gov/ct2/show/NCT04386850> [Accessed May 19, 2020]
140. dalili nooshin. Effects of Standard Protocol Therapy With or Without Colchicine in Covid-19 Infection: A Randomized Double Blind Clinical Trial. clinicaltrials.gov (2020). Available at: <https://clinicaltrials.gov/ct2/show/NCT04360980> [Accessed May 18, 2020]
141. Ricordi C. Umbilical Cord-derived Mesenchymal Stem Cells for COVID-19 Patients With Acute Respiratory Distress Syndrome (ARDS). clinicaltrials.gov (2020). Available at: <https://clinicaltrials.gov/ct2/show/NCT04355728> [Accessed May 18, 2020]
142. Urinary Incontinence and Sedentary Behaviour in Nursing Homes - Full Text View - ClinicalTrials.gov. Available at: <https://clinicaltrials.gov/ct2/show/NCT04297904> [Accessed May 19, 2020]

143. Prevention and Treatment With Calcifediol of COVID-19 Induced Acute Respiratory Syndrome - Full Text View - ClinicalTrials.gov. Available at: <https://clinicaltrials.gov/ct2/show/NCT04366908> [Accessed May 19, 2020]
144. Braiman M. Latitude Dependence of the COVID-19 Mortality Rate—A Possible Relationship to Vitamin D Deficiency? Rochester, NY: Social Science Research Network (2020). doi:10.2139/ssrn.3561958 (Preprint)
145. Gruber-Bzura BM. Vitamin D and Influenza—Prevention or Therapy? *Int J Mol Sci* (2018) **19**: doi:10.3390/ijms19082419
146. Sabetta JR, DePetrillo P, Cipriani RJ, Smardin J, Burns LA, Landry ML. Serum 25-Hydroxyvitamin D and the Incidence of Acute Viral Respiratory Tract Infections in Healthy Adults. *PLoS One* (2010) **5**: doi:10.1371/journal.pone.0011088
147. Aranow C. Vitamin D and the Immune System. *Journal of Investigative Medicine* (2011) **59**:881–886. doi:10.2310/JIM.0b013e31821b8755
148. Bäcker A. Slower COVID-19 Morbidity and Mortality Growth at Higher Solar Irradiance and Elevation. Rochester, NY: Social Science Research Network (2020). doi:10.2139/ssrn.3604729 (Preprint)
149. Arboleda J, Urcuqui-Inchima S. Vitamin D supplementation: a potential approach for COVID-19 therapeutics? Open Science Framework (2020). doi:10.31219/osf.io/cgd4t
150. Ardriaria M. PERAN VITAMIN D DALAM PENCEGAHAN INFLUENZA DAN COVID-19. *JNH (Journal of Nutrition and Health)* (2020) **8**:79–85. doi:10.14710/jnh.8.2.2020.79-85
151. Arya A, Dwivedi VD. Synergistic effect of Vitamin D and Remdesivir can fight COVID-19. *Journal of Biomolecular Structure and Dynamics* (2020) **0**:1–2. doi:10.1080/07391102.2020.1773929
152. Aslan MT, Aslan İÖ, Özdemir Ö. Is Vitamin D One of the Key Elements in COVID-19 Days? *J Nutr Health Aging* (2020) doi:10.1007/s12603-020-1413-5
153. Mohammed Hamad MN. Vitamin D Supplements Improve Efficacy of Minocycline, N-Acetylcysteine and Aspirin Triple Therapy to COVID-19 Infection. *SJBR* (2020) **05**:59–60. doi:10.36348/sjbr.2020.v05i04.005
154. Hobday RA, Cason JW. The Open-Air Treatment of PANDEMIC INFLUENZA. *Am J Public Health* (2009) **99**:S236–S242. doi:10.2105/AJPH.2008.134627
155. Lau FH, Majumder R, Torabi R, Saeg F, Hoffman R, Cirillo JD, Greiffenstein P. Vitamin D Insufficiency is Prevalent in Severe COVID-19. *medRxiv* (2020)2020.04.24.20075838. doi:10.1101/2020.04.24.20075838 (Preprint)
156. Tobolowsky FA. COVID-19 Outbreak Among Three Affiliated Homeless Service Sites — King County, Washington, 2020. *MMWR Morb Mortal Wkly Rep* (2020) **69**: doi:10.15585/mmwr.mm6917e2

157. Baggett TP, Keyes H, Sporn N, Gaeta JM. Prevalence of SARS-CoV-2 Infection in Residents of a Large Homeless Shelter in Boston. *JAMA* (2020) **323**:2191–2192. doi:10.1001/jama.2020.6887
158. de Lucena TMC, da Silva Santos AF, de Lima BR, de Albuquerque Borborema ME, de Azevêdo Silva J. Mechanism of inflammatory response in associated comorbidities in COVID-19. *Diabetes Metab Syndr* (2020) **14**:597–600. doi:10.1016/j.dsx.2020.05.025
159. Cantorna MT, Yu S, Bruce D. The paradoxical effects of vitamin D on Type 1 mediated immunity. *Mol Aspects Med* (2008) **29**:369–375. doi:10.1016/j.mam.2008.04.004
160. Mohammad S, Mishra A, Ashraf MZ. Emerging Role of Vitamin D and its Associated Molecules in Pathways Related to Pathogenesis of Thrombosis. *Biomolecules* (2019) **9**: doi:10.3390/biom9110649
161. Sundaram ME, Coleman LA. Vitamin D and Influenza. *Adv Nutr* (2012) **3**:517–525. doi:10.3945/an.112.002162
162. CANNELL JJ, VIETH R, UMHAU JC, HOLICK MF, GRANT WB, MADRONICH S, GARLAND CF, GIOVANNUCCI E. Epidemic influenza and vitamin D. *Epidemiol Infect* (2006) **134**:1129–1140. doi:10.1017/S0950268806007175
163. Martín Giménez VM, Inserra F, Tajer CD, Mariani J, Ferder L, Reiter RJ, Manucha W. Lungs as target of COVID-19 infection: Protective common molecular mechanisms of vitamin D and melatonin as a new potential synergistic treatment. *Life Sci* (2020)117808. doi:10.1016/j.lfs.2020.117808
164. Sparavigna, Amelia Carolina. Vitamin D for Covid-19? *Politecnico Di Torino* (2020) doi:10.5281/ZENODO.3822187
165. Meftahi GH, Jangravi Z, Sahraei H, Bahari Z. The possible pathophysiology mechanism of cytokine storm in elderly adults with COVID-19 infection: the contribution of “inflamm-aging.” *Inflamm Res* (2020) doi:10.1007/s00011-020-01372-8
166. Biesalski HK. Vitamin D deficiency and co-morbidities in COVID-19 patients – A fatal relationship? *NFS Journal* (2020) **20**:10–21. doi:10.1016/j.nfs.2020.06.001
167. Weir EK, Thenappan T, Bhargava M, Chen Y. Does vitamin D deficiency increase the severity of COVID-19? *Clinical Medicine* (2020) doi:10.7861/clinmed.2020-0301
168. Ulhaq ZS, Soraya GV. Interleukin-6 as a potential biomarker of COVID-19 progression. *Médecine et Maladies Infectieuses* (2020) **50**:382–383. doi:10.1016/j.medmal.2020.04.002
169. Silberstein M. Premorbid IL-6 levels may predict mortality from COVID-19. (2020) doi:10.21203/rs.3.rs-31814/v1 (Preprint)
170. Froum S. Vitamin D and COVID-19 disease severity: An old ally in a new war. *Perio-Implant Advisory* (2020) Available at: <https://www.perioimplantadvisory.com/clinical-tips/article/14176684/vitamin-d-and-covid19-disease-severity-an-old-ally-in-a-new-war> [Accessed June 16, 2020]

171. Speeckaert MM, Delanghe JR. Association between low vitamin D and COVID-19: don't forget the vitamin D binding protein. *Aging Clin Exp Res* (2020) doi:10.1007/s40520-020-01607-y
172. Daneshkhah A, Agrawal V, Eshein A, Subramanian H, Roy HK, Backman V. The Possible Role of Vitamin D in Suppressing Cytokine Storm and Associated Mortality in COVID-19 Patients. *medRxiv* (2020)2020.04.08.20058578. doi:10.1101/2020.04.08.20058578 (Preprint)
173. Muscogiuri G, Pugliese G, Barrea L, Savastano S, Colao A. Obesity: the "Achilles heel" for COVID-19? *Metabolism* (2020) doi:10.1016/j.metabol.2020.154251
174. Huibers MHW, Visser DH, Deckers MML, van Schoor NM, van Furth AM, Wolf BHM. Vitamin D deficiency among native Dutch and first- and second-generation non-Western immigrants. *Eur J Pediatr* (2014) **173**:583–588. doi:10.1007/s00431-013-2198-x
175. Yousef S, Elliott J, Manuel D, Colman I, Papadimitropoulos M, Hossain A, Leclair N, Wells GA. Study protocol: Worldwide comparison of vitamin D status of immigrants from different ethnic origins and native-born populations—a systematic review and meta-analysis. *Systematic Reviews* (2019) **8**:211. doi:10.1186/s13643-019-1123-4
176. Sääf M, Fernell E, Kristiansson F, Barnevik Olsson M, Gustafsson SA, Bågenholm G. Severe vitamin D deficiency in pregnant women of Somali origin living in Sweden: Vitamin D deficiency in pregnant Somali women. *Acta Paediatrica* (2011) **100**:612–614. doi:10.1111/j.1651-2227.2011.02134.x
177. Gasmi A, Noor S, Tippairote T, Dadar M, Menzel A, Björklund G. Individual risk management strategy and potential therapeutic options for the COVID-19 pandemic. *Clin Immunol* (2020) doi:10.1016/j.clim.2020.108409
178. Holick Michael F, Matsuoka Lois Y, Wortsman J. AGE, VITAMIN D, AND SOLAR ULTRAVIOLET. *The Lancet* (1989) **334**:1104–1105. doi:10.1016/S0140-6736(89)91124-0
179. Grant WB, Anouti FA, Moukayed M. Targeted 25-hydroxyvitamin D concentration measurements and vitamin D 3 supplementation can have important patient and public health benefits. *Eur J Clin Nutr* (2020) **74**:366–376. doi:10.1038/s41430-020-0564-0
180. Bäcker A, Mageswaran M. Double COVID-19 Confirmed Case Fatality Rate in Countries with High Elderly Female Vitamin D Deficiency Prevalence. Rochester, NY: Social Science Research Network (2020). doi:10.2139/ssrn.3623662 (Preprint)
181. Annweiler C, Cao Z, Wu Y, Faucon E, Mouhat S, Kovacic H, Sabatier J-M. Counter-regulatory "Renin-Angiotensin" System-based Candidate Drugs to Treat COVID-19 Diseases in SARS-CoV-2-infected patients. *Infect Disord Drug Targets* (2020) doi:10.2174/1871526520666200518073329
182. Walker J, Hiramoto J, Gasper W, Auyang P, Conte MS, Rapp J, Lovett D, Owens CD. Vitamin D deficiency is associated with mortality and adverse vascular access outcomes in patients with end stage renal disease. *J Vasc Surg* (2014) **60**:176–183. doi:10.1016/j.jvs.2014.01.037
183. Kennel KA, Drake MT, Hurley DL. Vitamin D Deficiency in Adults: When to Test and How to Treat. *Mayo Clin Proc* (2010) **85**:752–758. doi:10.4065/mcp.2010.0138

184. Lyons I, Ward V, Johnson J, Horton H. These are the health workers who have died from coronavirus. *The Telegraph* (2020) Available at: <https://www.telegraph.co.uk/news/0/nhs-workers-died-coronavirus-frontline-victims/> [Accessed April 29, 2020]
185. Swerling G, Penna D. Why are so many black and ethnic minority people dying from coronavirus? *The Telegraph* (2020) Available at: <https://www.telegraph.co.uk/news/0/bame-coronavirus-black-ethnic-minority/> [Accessed May 2, 2020]
186. Ahmed I, Azhar A, Eltaweel N, Tan BK. First Covid-19 maternal mortality in the UK associated with thrombotic complications. *Br J Haematol* (2020) doi:10.1111/bjh.16849
187. Muscogiuri G, Barrea L, Savastano S, Colao A. Nutritional recommendations for CoVID-19 quarantine. *European Journal of Clinical Nutrition* (2020) **74**:850–851. doi:10.1038/s41430-020-0635-2
188. Ekiz T, Pazarlı AC. Relationship between COVID-19 and obesity. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews* (2020) **14**:761–763. doi:10.1016/j.dsx.2020.05.047
189. Palaniveloo L, Yong HY, Mohd Shariff Z, Peng LS, Bindels J, Tee YYS, van der Beek EM. Vitamin D status is associated with high BMI, working status and gravidity among pregnant Malaysian women. *Mal J Nutr* (2020) **26**:129–139. doi:10.31246/mjn-2020-0010
190. Ekiz T, Kara M, Özçakar L. Revisiting vitamin D and home-based exercises for patients with sleep apnea facing the COVID-19 quarantine. *J Clin Sleep Med* (2020) doi:10.5664/jcsm.8586
191. Khunti K, Singh AK, Pareek M, Hanif W. Is ethnicity linked to incidence or outcomes of covid-19? *BMJ* (2020) **369**: doi:10.1136/bmj.m1548
192. Facchiano A, Facchiano A, Bartoli M, Ricci A, Facchiano F. Reply to Jakovac: About COVID-19 and vitamin D. *Am J Physiol Endocrinol Metab* (2020) **318**:E838. doi:10.1152/ajpendo.00185.2020
193. Yao Y, Pan J, Liu Z, Meng X, Wang W, Kan H, Wang W. No Association of COVID-19 transmission with temperature or UV radiation in Chinese cities. *Eur Respir J* (2020) doi:10.1183/13993003.00517-2020
194. Chinazzi M, Davis JT, Ajelli M, Gioannini C, Litvinova M, Merler S, Piontti AP y, Rossi L, Sun K, Viboud C, et al. The effect of travel restrictions on the spread of the 2019 novel coronavirus (2019-nCoV) outbreak. *medRxiv* (2020)2020.02.09.20021261. doi:10.1101/2020.02.09.20021261
195. Li AY, Hannah TC, Durbin J, Dreher N, McAuley FM, Marayati NF, Spiera Z, Ali M, Gometz A, Kostman JT, et al. Multivariate Analysis of Factors Affecting COVID-19 Case and Death Rate in U.S. Counties: The Significant Effects of Black Race and Temperature. *The American Journal of the Medical Sciences* (2020) <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7305735/>
196. Panarese A, Shahini E. Letter: Covid-19, and vitamin D. *Aliment Pharmacol Ther* (2020) doi:10.1111/apt.15752

197. Kumar V, Srivastava A. Spurious Correlation? A review of the relationship between Vitamin D and Covid-19 infection and mortality. *medRxiv* (2020)2020.05.25.20110338. doi:10.1101/2020.05.25.20110338 (Preprint)
198. Singh S, Kaur R, Singh RK. Revisiting the role of vitamin D levels in the prevention of COVID-19 infection and mortality in European countries post infections peak. (2020) doi:10.21203/rs.3.rs-32484/v1 (Preprint)
199. Notari A, Torrieri G. COVID-19 transmission risk factors. *arXiv:200503651 [physics, q-bio, stat]* (2020) Available at: <http://arxiv.org/abs/2005.03651> [Accessed May 20, 2020]
200. Coronavirus Update (Live): 4,005,655 Cases and 275,669 Deaths from COVID-19 Virus Pandemic - Worldometer. Available at: <https://www.worldometers.info/coronavirus/#countries> [Accessed May 8, 2020]
201. Osmanovic A, Demeke T, Gillstedt M, Angesjö E, Sinclair H, Abd El-Gawad G, Landin-Wilhelmsen K. Vitamin D treatment in Somali women living in Sweden - Two randomized, placebo-controlled studies. *Clin Endocrinol (Oxf)* (2016) **85**:535–543. doi:10.1111/cen.13097
202. Bäcker A. Non-black Hispanic/Latinos Spend More Time Outdoors and Have Lower COVID-19 Morbidity Than Non-Hispanic/Latinos. Rochester, NY: Social Science Research Network (2020). doi:10.2139/ssrn.3575217 (Preprint)
203. Li M, Zhang Z, Cao W, Liu Y, Du B, Chen C, Liu Q, Uddin MN, Jiang S, Chen C, et al. Identifying novel factors associated with COVID-19 transmission and fatality using the machine learning approach. *medRxiv* (2020)2020.06.10.20127472. doi:10.1101/2020.06.10.20127472 (Preprint)
204. Kohlmeier M. Avoidance of vitamin D deficiency to slow the COVID-19 pandemic. *BMJ Nutrition, Prevention & Health* (2020)bmjnph-2020-000096. doi:10.1136/bmjnph-2020-000096
205. Alipio MM. Do latitude and ozone concentration predict Covid-2019 cases in 34 countries? *medRxiv* (2020)2020.04.09.20060202. doi:10.1101/2020.04.09.20060202 (Preprint)
206. Skutsch M, Dobler C, McCall MBB, Ghilardi A, Salinas-Melgoza MA, McCall MK, Fenner-Sanchez G. The association of UV with rates of COVID-19 transmission and deaths in Mexico: the possible mediating role of vitamin D. *medRxiv* (2020)2020.05.25.20112805. doi:10.1101/2020.05.25.20112805 (Preprint)
207. Li Y, Li Q, Zhang N, Liu Z. Sunlight and vitamin D in the prevention of coronavirus disease (COVID-19) infection and mortality in the United States. (2020) (Preprint)
208. Marik PE, Kory P, Varon J. Does vitamin D status impact mortality from SARS-CoV-2 infection? *Med Drug Discov* (2020)100041. doi:10.1016/j.medidd.2020.100041
209. Annweiler C, Cao Z, Sabatier J-M. Point of view: Should COVID-19 patients be supplemented with vitamin D? *Maturitas* (2020) **140**:24–26. doi:10.1016/j.maturitas.2020.06.003



210. Horowitz RI, Freeman PR, Bruzzese J. Efficacy of glutathione therapy in relieving dyspnea associated with COVID-19 pneumonia: A report of 2 cases. *Respiratory Medicine Case Reports* (2020) **30**:101063. doi:10.1016/j.rmcr.2020.101063
211. Bossoni S, Chiesa L, Giustina A. Severe hypocalcemia in a thyroidectomized woman with Covid-19 infection. *Endocrine* (2020)1–2. doi:10.1007/s12020-020-02326-0
212. Pinzon RT, Angela A, Pradana AW. Vitamin D Deficiency Among Patients with COVID-19 : Case Series and Recent Literature Review. In Review (2020). doi:10.21203/rs.3.rs-29473/v1 (Preprint)
213. Faul JL, Kerley C, Love B, O-Nell E, Cody C, Tormey W, Hutchinson K, Cormican LJ, Burke CM. Vitamin D Deficiency and ARDS after SARS-CoV-2 Infection – Irish Medical Journal. *Irish Medical Journal* (2020) **113**:84. Available at: <http://imj.ie/vitamin-d-deficiency-and-ards-after-sars-cov-2-infection/> [Accessed May 14, 2020]
214. Fasano A, Cereda E, Barichella M, Cassani E, Ferri V, Zecchinelli AL, Pezzoli G. COVID-19 in Parkinson’s Disease Patients Living in Lombardy, Italy. *Mov Disord* (2020) doi:10.1002/mds.28176
215. Tan CW, Ho LP, Kalimuddin S, Cherng BPZ, Teh YE, Thien SY, Wong HM, Tern PJW, Chay JWM, Nagarajan C, et al. A cohort study to evaluate the effect of combination Vitamin D, Magnesium and Vitamin B12 (DMB) on progression to severe outcome in older COVID-19 patients. *medRxiv* (2020)2020.06.01.20112334. doi:10.1101/2020.06.01.20112334 (Preprint)
216. Asyary A, Veruswati M. Sunlight exposure increased Covid-19 recovery rates: A study in the central pandemic area of Indonesia. *Sci Total Environ* (2020) doi:10.1016/j.scitotenv.2020.139016
217. Sun J-K, Zhang W-H, Zou L, Liu Y, Li J-J, Kan X-H, Dai L, Shi Q-K, Yuan S-T, Yu W-K, et al. Serum calcium as a biomarker of clinical severity and prognosis in patients with coronavirus disease 2019: a retrospective cross-sectional study. In Review (2020). doi:10.21203/rs.3.rs-17575/v1 (Preprint)
218. Cuñat T, Ojeda A, Calvo A. Vitamin D deficiency in critically ill patients diagnosed with COVID -19. Are we doing enough? A retrospective analysis of 226 patients. In Review (2020). doi:10.21203/rs.3.rs-30390/v1 (Preprint)
219. De Smet D, De Smet K, Herroelen P, Gryspeerd S, Martens GA. Vitamin D deficiency as risk factor for severe COVID-19: a convergence of two pandemics. *medRxiv* (2020)2020.05.01.20079376. doi:10.1101/2020.05.01.20079376 (Preprint)
220. Meltzer DO, Best TJ, Zhang H, Vokes T, Arora V, Solway J. Association of Vitamin D Deficiency and Treatment with COVID-19 Incidence. *medRxiv* (2020)2020.05.08.20095893. doi:10.1101/2020.05.08.20095893 (Preprint)
221. Matos RI, Chung KK. DoD COVID-19 Practice Management Guide Clinical Management of COVID-19 | Technical Resources | ASPR TRACIE. (2020) Available at: <https://asprtracie.hhs.gov/technical-resources/resource/7899/dod-covid-19-practice-management-guide-clinical-management-of-covid-19> [Accessed May 20, 2020]

222. Dasgupta J, Sen U, Bakshi A, Dasgupta A, Manna K, Saha C, De RK, Mukhopadhyay S, Bhattacharyya NP. Nsp7 and Spike Glycoprotein of SARS-CoV-2 Are Envisaged as Potential Targets of Vitamin D and Ivermectin. (2020) doi:10.20944/preprints202005.0084.v1
223. Lanham-New SA, Webb AR, Cashman KD, Buttriss JL, Fallowfield JL, Masud T, Hewison M, Mathers JC, Kiely M, Welch AA, et al. Vitamin D and SARS-CoV-2 virus/COVID-19 disease. *BMJ Nutrition, Prevention & Health* (2020)bmjnph-2020-000089. doi:10.1136/bmjnph-2020-000089
224. Pal R, Bhadada SK. Managing common endocrine disorders amid COVID-19 pandemic. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews* (2020) **14**:767–771. doi:10.1016/j.dsx.2020.05.050
225. Nizami NS, Mujeebuddin CS. Strong Immunity - A Major Weapon to Fight against Covid-19. *IOSR Journal Of Pharmacy And Biological Sciences* (2020) **15**:22–29. doi:10.9790/1503032229
226. Parvin F, Islam S, Urmy Z, Ahmed S. The symptoms, contagious process, prevention and post treatment of Covid-19. *European Journal of Physiotherapy and Rehabilitation Studies* (2020) **0**: Available at: <https://oapub.org/hlt/index.php/EJPRS/article/view/52> [Accessed May 8, 2020]
227. Richards G, Mer M, Schleicher G, Stacey S. COVID-19 and the rationale for pharmacotherapy : a South African perspective. *Wits Journal of Clinical Medicine* (2020) **2**:11–18. doi:10.18772/26180197.2020.v2nSla2
228. Wimalawansa SJ. Global epidemic of coronavirus—Covid-19: What can we do to minimize risks. *European Journal of Biomedical and Pharmaceutical Sciences* (2020) **7**:7. Available at: [https://www.researchgate.net/profile/Sunil\\_Wimalawansa/publication/340102912\\_EJBPS-Prof\\_Wimalawansa-COVID-19-March\\_2020s/links/5e78eb434585158bd50069f3/EJBPS-Prof-Wimalawansa-COVID-19-March-2020s.pdf](https://www.researchgate.net/profile/Sunil_Wimalawansa/publication/340102912_EJBPS-Prof_Wimalawansa-COVID-19-March_2020s/links/5e78eb434585158bd50069f3/EJBPS-Prof-Wimalawansa-COVID-19-March-2020s.pdf)
229. Puig-Domingo M, Marazuela M, Giustina A. COVID-19 and endocrine diseases. A statement from the European Society of Endocrinology. *Endocrine* (2020)1–4. doi:10.1007/s12020-020-02294-5
230. Koenig HG. Ways of Protecting Religious Older Adults from the Consequences of COVID-19. *Am J Geriatr Psychiatry* (2020) doi:10.1016/j.jagp.2020.04.004
231. Uçar D, Tayfun K, Müslümanoğlu AY, Kalayci MZ. Coronavirus ve Fitoterapi. *Bütünleyici ve Anadolu Tıbbı Dergisi* (2020) **1**:49–57. Available at: <https://dergipark.org.tr/en/pub/batd/711108> [Accessed May 20, 2020]
232. Zicarelli V. Nutrition Therapy for Severe Viral Infections (COVID-19): Recommendations and Considerations for Integrative Medical Treatments. *Journal of Orthomolecular Medicine* (2020) **35**: Available at: <https://isom.ca/article/nutrition-therapy-for-severe-viral-infections-covid-19/> [Accessed June 26, 2020]
233. Suresh PS. Hypovitaminosis D and COVID-19: Matter of Concern in India? *Ind J Clin Biochem* (2020) doi:10.1007/s12291-020-00894-6
234. Queiroz NSF, Barros LL, de Azevedo MFC, Oba J, Sobrado CW, de Sousa Carlos A, Milani LR, Sipahi AM, Damião AOMC. Management of inflammatory bowel disease patients in the COVID-19

- pandemic era: a Brazilian tertiary referral center guidance. *Clinics (Sao Paulo)* (2020) **75**: doi:10.6061/clinics/2020/e1909
235. Nihra V. Identifying Patterns in Covid-19: Morbidity, Recovery, and the Aftermath. (2020) doi:10.13140/RG.2.2.30040.93449
236. Docea AO, Tsatsakis A, Albulescu D, Cristea O, Zlatian O, Vinceti M, Moschos SA, Tsoukalas D, Goumenou M, Drakoulis N, et al. A new threat from an old enemy: Re-emergence of coronavirus (Review). *Int J Mol Med* (2020) **45**:1631–1643. doi:10.3892/ijmm.2020.4555
237. Gee S, Gaughran F, MacCabe J, Shergill S, Whiskey E, Taylor D. Management of clozapine treatment during the COVID-19 pandemic: *Therapeutic Advances in Psychopharmacology* (2020) doi:10.1177/2045125320928167
238. Kurthkoti K, Das G. Potential benefits of Vitamin D supplementation in COVID-19 patients: a hypothesis. *Open Science Framework* (2020). doi:10.31219/osf.io/dnjav
239. Ramos EM, Abreu AC de, Freitas SLF de, Lima MD de, Reis FJM dos, Ramos HV, Souza ID de, Bochenek L de MS, Fonseca AC da, Nascimento VA do. COVID-19, Rate of Case Factors and Nutritional Characteristics of Patients Dying in Italy and Brazil: A Critical Analyze. *Global Journal of Health Science* (2020) **12**:p133. doi:10.5539/gjhs.v12n7p133
240. Razdan K, Singh K, Singh D. Vitamin D Levels and COVID-19 Susceptibility: Is there any Correlation? *Medicine in Drug Discovery* (2020) **7**:100051. doi:10.1016/j.medidd.2020.100051
241. Saul AW. Nutritional Treatment of Coronavirus. *Orthomolecular Medicine News Service* (2020) Available at: <http://orthomolecular.org/resources/omns/v16n06.shtml>
242. Siuka D, Pfeifer M, Pinter B. Vitamin D supplementation in the COVID-19 pandemic. *Mayo Clinic Proceedings* (2020) **0**: doi:10.1016/j.mayocp.2020.05.036
243. Tan SHS, Hong CC, Saha S, Murphy D, Hui JH. Medications in COVID-19 patients: summarizing the current literature from an orthopaedic perspective. *Int Orthop* (2020) doi:10.1007/s00264-020-04643-5
244. Tian Y, Rong L. Letter: does vitamin D have a potential role against COVID-19? Authors' reply. *Aliment Pharmacol Ther* (2020) doi:10.1111/apt.15817
245. Yousfi N, Bragazzi NL, Briki W, Zmijewski P, Chamari K. The COVID-19 pandemic: how to maintain a healthy immune system during the quarantine - a multidisciplinary approach with special focus on athletes. *Biology of Sport* (2020) doi:10.5114/biol sport.2020.95125
246. Zabetakis I, Lordan R, Norton C, Tsoupras A. COVID-19: The Inflammation Link and the Role of Nutrition in Potential Mitigation. *Nutrients* (2020) **12**: doi:10.3390/nu12051466
247. Zemb P, Bergman P, Camargo CA, Cavalier E, Cormier C, Courbebaisse M, Hollis B, Minisola S, Pilz S, Pludowski P, et al. Vitamin D deficiency and COVID-19 pandemic. *J Glob Antimicrob Resist* (2020) doi:10.1016/j.jgar.2020.05.006

248. Piller C. ‘This is insane!’ Many scientists lament Trump’s embrace of risky malaria drugs for coronavirus. *Science / AAAS* (2020) Available at: <https://www.sciencemag.org/news/2020/03/insane-many-scientists-lament-trump-s-embrace-risky-malaria-drugs-coronavirus> [Accessed April 23, 2020]
249. NIH. Office of Dietary Supplements - Vitamin D. Available at: <https://ods.od.nih.gov/factsheets/VitaminD-HealthProfessional/> [Accessed April 23, 2020]
250. McCartney DM, Byrne DG. McCartney et al comment on ‘Vitamin D and Covid-19 – A Note of Caution’ response letter – Irish Medical Journal. *Irish Medical Journal* **113**:79. Available at: <http://imj.ie/mccartney-et-al-comment-on-vitamin-d-and-covid-19-a-note-of-caution-response-letter/> [Accessed May 14, 2020]
251. Cronise RJ, Sinclair DA, Bremer AA. The “Metabolic Winter” Hypothesis: A Cause of the Current Epidemics of Obesity and Cardiometabolic Disease. *Metab Syndr Relat Disord* (2014) **12**:355–361. doi:10.1089/met.2014.0027
252. Siddiqui R, Ahmed Khan N. Centralized air-conditioning and transmission of novel coronavirus. *Pathog Glob Health* (2020)1–2. doi:10.1080/20477724.2020.1765653
253. Nigro E, Polito R, Alfieri A, Mancini A, Imperlini E, Elce A, Krustrup P, Orrù S, Buono P, Daniele A. Molecular Mechanisms Involved in the Positive Effects of Physical Activity on Coping with COVID-19. (2020) (Preprint)
254. Ilie PC, Stefanescu S, Smith L. The role of Vitamin D in the prevention of Coronavirus Disease 2019 infection and mortality. In Review (2020). doi:10.21203/rs.3.rs-21211/v1
255. Kara M, Ekiz T, Ricci V, Kara Ö, Chang K-V, Özçakar L. “Scientific Strabismus” or Two Related Pandemics: COVID-19 & Vitamin D Deficiency. *Br J Nutr* (2020)1–20. doi:10.1017/S0007114520001749
256. Rhodes JM, Subramanian S, Laird E, Kenny RA. Editorial: low population mortality from COVID-19 in countries south of latitude 35 degrees North – supports vitamin D as a factor determining severity. *Alimentary Pharmacology & Therapeutics n/a*: doi:10.1111/apt.15777
257. Alabada HKM, Saleh WMM. Vitamin D effectiveness and pathology in humans and domestic animals. *Multi* (2020) **3**:15.
258. Álvarez-López JA, García-Contreras AI. Vitamina D y la pandemia por COVID-19. *RME* (2020) **7**:4329. doi:10.24875/RME.20000036
259. Anderson G, Reiter RJ. COVID-19 pathophysiology: interactions of gut microbiome, melatonin, vitamin D, stress, kynurenine and the alpha 7 nicotinic receptor: Treatment implications. *Melatonin Research* (2020) **3**:322–345. doi:10.32794/mr11250066
260. Alpalhão M, Filipe P. SARS-CoV-2 pandemic and Vitamin D deficiency—A double trouble. *Photodermatology, Photoimmunology & Photomedicine n/a*: doi:10.1111/phpp.12579

261. Aygun H. Vitamin D can prevent COVID-19 infection-induced multiple organ damage. *Naunyn Schmiedebergs Arch Pharmacol* (2020) doi:10.1007/s00210-020-01911-4
262. Barazzoni R, Bischoff SC, Breda J, Wickramasinghe K, Krznaric Z, Nitzan D, Pirlich M, Singer P. ESPEN expert statements and practical guidance for nutritional management of individuals with SARS-CoV-2 infection. *Clin Nutr* (2020) doi:10.1016/j.clnu.2020.03.022
263. Bauer SR, Kapoor A, Rath M, Thomas SA. What is the role of supplementation with ascorbic acid, zinc, vitamin D, or N-acetylcysteine for prevention or treatment of COVID-19? *CCJM* (2020) doi:10.3949/ccjm.87a.ccc046
264. Belančić A, Kresović A, Rački V. Potential pathophysiological mechanisms leading to increased COVID-19 susceptibility and severity in obesity. *Obes Med* (2020) **19**:100259. doi:10.1016/j.obmed.2020.100259
265. Brown RA, Sarkar A. Vitamin D deficiency: a factor in COVID-19, progression, severity and mortality? – An urgent call for research. *Modern Ghana* (2020) Available at: <https://www.modernghana.com/news/992403/vitamin-d-deficiency-a-factor-in-covid-19-progre.html> [Accessed April 23, 2020]
266. Brown RA, Rhein H, Alipio MM, Annweiler C, Gnaiger E, Holick MF, Boucher BJ, Duque G, Féron F, Kenny RA, Montero-Odasso M, McCarrison Society, La Route de Mont Cochon, St Lawrence, Jersey, Grimes D, de Voil CWB,, Minisola S, Rhodes J, Haq A, Bejerot S, Reiss LAJ, Zgaga L, Crawford MA, Fricker RA, Cobbold P, Lahore HW, Humble MB, Sakar A, Karras S, Iglesias-Gonzalez J, Gezen-Ak D, Dursun E, Cooper I,. COVID-19 'ICU' risk – 20-fold greater in the Vitamin D Deficient. BAME, African Americans, the Older, Institutionalised and Obese, are at greatest risk. Sun and 'D'-supplementation – Game-changers? Research urgently required. *BMJ* (2020) **2020;369:m1548/rr-6**: Available at: <https://www.bmj.com/content/369/bmj.m1548/rr-6> [Accessed June 24, 2020]
267. Chan B, Wong C, Leung PC. What can we do for the Personal Protection against the CoVID-19 Infection? Immuno-Boosting Specific Supplement could be the Answer. *J Emerg Med Trauma Surg Care* (2020) **2**:
268. Dhillon P, Breuer M, Hirst N. COVID-19 breakthroughs: separating fact from fiction. *The FEBS Journal* (2020)
269. Farias E dos S, Silva DF, Conceição VS da, Rocha AM. Vitamin D as an Immunological Factor in Combating COVID-19. *International Journal of Advanced Engineering Research and Science* (2020) **7**: Available at: <http://journal-repository.com/index.php/ijaers/article/view/2067> [Accessed June 16, 2020]
270. Gasmi A, Noor S, Tippairote T, Dadar M, Menzel A, Bjørklund G. Individual risk management strategy and potential therapeutic options for the COVID-19 pandemic. *Clinical Immunology* (2020) **215**:108409. doi:10.1016/j.clim.2020.108409
271. Glinsky GV. Tripartite Combination of Candidate Pandemic Mitigation Agents: Vitamin D, Quercetin, and Estradiol Manifest Properties of Medicinal Agents for Targeted Mitigation of the

- COVID-19 Pandemic Defined by Genomics-Guided Tracing of SARS-CoV-2 Targets in Human Cells. *Biomedicines* (2020) **8**: doi:10.3390/biomedicines8050129
272. Vázquez--Tanus JB. COVID-19 Protocol. Puerto Rico: Vascular Nutrition PR Website (4/12/2020). Available at: <https://www.vascularnutritionpr.com/covid-19.html> [Accessed June 22, 2020]
273. Grant WB, Lahore H, McDonnell SL, Baggerly CA, French CB, Aliano JL, Bhattoa HP. Evidence that Vitamin D Supplementation Could Reduce Risk of Influenza and COVID-19 Infections and Deaths. *Nutrients* (2020) **12**:988. doi:10.3390/nu12040988
274. Guan H, Okely AD, Aguilar-Farias N, del Pozo Cruz B, Draper CE, El Hamdouchi A, Florindo AA, Jáuregui A, Katzmarzyk PT, Kontsevaya A, et al. Promoting healthy movement behaviours among children during the COVID-19 pandemic. *Lancet Child Adolesc Health* (2020) **4**:416–418. doi:10.1016/S2352-4642(20)30131-0
275. Gupta A. Is Immuno-modulation the Key to COVID-19 Pandemic? *Indian J Orthop* (2020)1–4. doi:10.1007/s43465-020-00121-7
276. Hasan A, Paray BA, Hussain A, Qadir FA, Attar F, Aziz FM, Sharifi M, Derakhshankhah H, Rasti B, Mehrabi M, et al. A review on the cleavage priming of the spike protein on coronavirus by angiotensin-converting enzyme-2 and furin. *J Biomol Struct Dyn* (2020) **0**:1–9. doi:10.1080/07391102.2020.1754293
277. Heiser K, McLean PF, Davis CT, Fogelson B, Gordon HB, Jacobson P, Hurst B, Miller B, Alfa RW, Earnshaw BA, et al. Identification of potential treatments for COVID-19 through artificial intelligence-enabled phenomic analysis of human cells infected with SARS-CoV-2. *bioRxiv* (2020)2020.04.21.054387. doi:10.1101/2020.04.21.054387 (Preprint)
278. Hossain Z, Uddin MdN. Natural Polyphenol Engineering Field Crops based Diet to Promote Innate Immunity to Combat Covid-19 Disease. *AgriXiv* (2020). doi:10.31220/osf.io/da85c (Preprint)
279. Jukic I, Calleja-González J, Cos F, Cuzzolin F, Olmo J, Terrados N, Njaradi N, Sassi R, Requena B, Milanovic L, et al. Strategies and Solutions for Team Sports Athletes in Isolation due to COVID-19. *Sports (Basel)* (2020) **8**: doi:10.3390/sports8040056
280. Kalantar-Zadeh K, Moore LW. Impact of Nutrition and Diet on COVID-19 Infection and Implications for Kidney Health and Kidney Disease Management. *J Ren Nutr* (2020) **30**:179–181. doi:10.1053/j.jrn.2020.03.006
281. Kamani E. Effect of Low Level Yellow Laser Light 589nm on Virus Prevention Corona Virus (COVID-19). *OSP Journal of Case Reports* (2020) **2**:1–4. Available at: [https://www.ospublishers.com/Effect-of-Low-Level-Yellow-Laser-Light-589nm-on-Virus-Prevention-Corona-Virus-\(COVID-19\).html](https://www.ospublishers.com/Effect-of-Low-Level-Yellow-Laser-Light-589nm-on-Virus-Prevention-Corona-Virus-(COVID-19).html) [Accessed June 16, 2020]
282. Khalil I, Barma P. Sub-continental Atmosphere and Inherent Immune System may have Impact on Novel Corona Virus' 2019 (nCovid-19) Prevalence in South East Asia. *Mymensingh Med J* (2020) **29**:473–480.

283. Kumar D, Gupta P, Banerjee D. Letter: does vitamin D have a potential role against COVID-19? *Aliment Pharmacol Ther* (2020) doi:10.1111/apt.15801
284. Lawton G. Will warmer spring weather slow down the rate of spread? *New Sci* (2020) **245**:8. doi:10.1016/S0262-4079(20)30659-X
285. Martín Giménez VM, Inserra F, Tajer CD, Mariani J, Ferder L, Reiter RJ, Manucha W. Lungs as target of COVID-19 infection: Protective common molecular mechanisms of vitamin D and melatonin as a new potential synergistic treatment. *Life Sci* (2020) **254**:117808. doi:10.1016/j.lfs.2020.117808
286. Maruta H, He H. PAK1-blockers: Potential Therapeutics against COVID-19. *Med Drug Discov* (2020) **6**:100039. doi:10.1016/j.medidd.2020.100039
287. McCartney DM, Byrne DG. McCartney et al comment on ‘Covid-19, Cocooning and Vitamin D Requirements’ response report – Irish Medical Journal. *Irish Medical Journal* (2020) **113**:80. Available at: <http://imj.ie/mccartney-et-al-comment-on-covid-19-cocooning-and-vitamin-d-requirements-response-report/> [Accessed May 14, 2020]
288. Merow C, Urban MC. Seasonality and uncertainty in COVID-19 growth rates. *medRxiv* (2020)2020.04.19.20071951. doi:10.1101/2020.04.19.20071951 (Preprint)
289. Musselwhite CBA. JTH editorial v17 – The importance of psychosocial factors in transport and health. *J Transp Health* (2020) doi:10.1016/j.jth.2020.100876
290. Palmer K, Monaco A, Kivipelto M, Onder G, Maggi S, Michel J-P, Prieto R, Sykara G, Donde S. The potential long-term impact of the COVID-19 outbreak on patients with non-communicable diseases in Europe: consequences for healthy ageing. *Aging Clin Exp Res* (2020) doi:10.1007/s40520-020-01601-4
291. Rabbitt L, Slattery E. Vitamin D and Covid-19 – A Note of Caution – Irish Medical Journal. *Irish Medical Journal* (2020) **113**:82. Available at: <http://imj.ie/vitamin-d-and-covid-19-a-note-of-caution/> [Accessed May 14, 2020]
292. Stuart Richer OD. How to build a lifestyle and nutritional firewall against viruses like COVID-19. *Optometry Times* (2020) **12**:1–7. Available at: <https://www.optometrytimes.com/article/how-build-lifestyle-and-nutritional-firewall-against-viruses-covid-19> [Accessed June 16, 2020]
293. Rocha JC, Calhau C, MacDonald A. Reply to Jakovac; Severity of COVID-19 infection in patients with phenylketonuria: is vitamin D status protective? *American Journal of Physiology-Endocrinology and Metabolism* (2020) **318**:E890–E891. doi:10.1152/ajpendo.00195.2020
294. Romano L, Bilotta F, Dauri M, Macheda S, Pujia A, De Santis GL, Tarsitano MG, Merra G, Di Renzo L, Esposito E, et al. Short Report - Medical nutrition therapy for critically ill patients with COVID-19. *Eur Rev Med Pharmacol Sci* (2020) **24**:4035–4039. doi:10.26355/eurrev\_202004\_20874
295. Rusciano D, Bagnoli P, Galeazzi R. The Fight Against COVID-19: The Role of Drugs and Food Supplements. *J Pharmacol Pharm Res* (2020) **3**:5–15. Available at:

<https://researchopenworld.com/the-fight-against-covid-19-the-role-of-drugs-and-food-supplements/> [Accessed June 17, 2020]

296. Silberstein M. Vitamin D: A simpler alternative to tocilizumab for trial in COVID-19? *Med Hypotheses* (2020) **140**:109767. doi:10.1016/j.mehy.2020.109767
297. Thomson B. The COVID-19 Pandemic: A Global Natural Experiment. *Circulation* (2020) doi:10.1161/CIRCULATIONAHA.120.047538
298. Yong SJ. Population Studies Confirm Risk Factors for Catching Covid-19. *Medium* (2020) Available at: <https://medium.com/microbial-instincts/population-studies-confirm-risk-factors-for-catching-covid-19-853779e7a3c6> [Accessed June 26, 2020]
299. Yong SJ. Vitamin D as an Independent Risk Factor for COVID-19 Death. *Medium* (2020) Available at: <https://medium.com/microbial-instincts/lack-of-vitamin-d-as-an-independent-risk-factor-for-covid-19-death-82365d0520fa> [Accessed June 28, 2020]
300. Zicarelli vince. Nutrition Therapy for Severe Viral Infections (COVID-19): Recommendations and Considerations for Integrative Medical Treatments. *Journal of Orthomolecular Medicine* (2020) **35**: Available at: <https://isom.ca/article/nutrition-therapy-for-severe-viral-infections-covid-19/> [Accessed June 16, 2020]



## **10 Appendices**

### **10.1 Table 2**

<b>Table 2: Brief Summaries of Original Research (in vitro research is included in Table 3) (47 entries)</b>					
1st Author /Location	Title (Truncated)	Number of Participants /Type*/ #pages	Supporting Information **	Results/ Recommendation Regarding COVID-19	Support for Supplementati on Now
Ahmed/ UK	First COVID-19 maternal mortality ...thrombotic complications(186)	1/CS 4 pages	T	Thrombosis kills/ Vigilance is needed	5
Alipio/ Philippines	Vitamin D ... possibly improve clinical outcomes of patients infected with ... (Covid-2019)(40)	212/R 6 pages	R,C	Sig. D-outcome link /D suppl. could improve outcomes	8
Alipio/ Philippines	Do latitude and ozone concentration predict Covid-2019 cases in 34 countries?(205)	34 countries /A 9 pages	R,	Sig COVID-19-UV (D) link, no latitude link /D suppl. indicated	7
Annweiler/France & China	Point of view: Should COVID-19 patients be supplemented with vitamin D?(209)	6 of 7 causal criteria/C 25 pages	G,P,R,F,C,J,S	Strong biological plausibility case - supplement D	9
Asyary/ Indonesia	Sunlight exposure increased COVID-19 recovery rates: A study in ... Indonesia(216)	9 hospitals /R 4 pages	G,I,R,F,	Patient exposure to sunshine helped Encourage it	7
Bäcker/ USA (see note)	Why COVID-19 May Be Disproportionately Killing African Americans...Irradiance... Income(11)	8 locations in the USA/ A 6 pages	G,I	Irradiation for disinfection, sun for protection	6
Bäcker/ USA (see note)	Slower COVID-19 morbidity and mortality growth at higher solar	159 locations/ A 58 21 pages	G,I,	Encourage people to go where the sun is overhead	6

	irradiance and elevation(148)				
ann/ USA (see note)	Non-black Hispanic/Latinos Spend More Time Outdoors ... Lower ... Morbidity Than Non-Hispanic...(202)	All of Georgia /A 7 pages	G,I,P	Urgently test sunlight exposure, especially for African Americans	6
Bäcker/ USA	Double COVID-19 ... Fatality Rate in Countries with High Elderly Female Vitamin D Deficiency Prevalence(180)	32 countries /A 11 pages	G,I,P,C,S	Test Vitamin D for prevention and treatment	6
Bossoni/ Italy	Severe hypocalcemia in a thyroidectomized woman with COVID-19 infection(211)	1/CS 2 pages	R	Warn thyroid patients due to home confinement	8
Braiman/ USA	Latitude ... COVID-19 mortality rate—a possible relationship to vitamin D deficiency?(144)	all nations (WHO data) P/ 14 pages	G	Test for link by supplementing and looking for changes	7
Cuñat/ Spain	Vitamin D deficiency in critically ill patients ... COVID -19. Are we doing enough? ... 226 patients.(218)	226 patients (subset: 17) /R/7 pages	P,R,C,S	Should assess D more often – all 17 were deficient	8
Daneshkhah/USA	The Possible Role of Vitamin D in Suppressing Cytokine Storm and ... Mortality in COVID-19 Patients(172)	10 countries /A/ 23 pages	C	Study if Vitamin D deficiency speeds mortality	6
Darling /UK	Vitamin D status, body mass index, ethnicity and COVID-19: ... UK	580 Covid+ vs 580 negative /R	None mentioned	All identified risk factors are uncontrollable - No	0

	Biobank COVID-19 ... (n 580) ... negative controls (n 723)(74)	3 pages		recommendations	
Davies/UK Switzerland	Evidence Supports a Causal Model for Vitamin D in COVID-19 Outcomes(61) (two studies)	Global data /A and C/ 30 pages	G,P,R	D deficiency is a cause of COVID-19 Give up to 4000IU	9
D'Avolio/ Switzerland, Italy	25-Hydroxyvitamin D ..Are Lower in Patients with Positive PCR for SARS-CoV-2(122)	107 Covid+ vs 1377 /R/ 7 pages	R	D correlated to + Supplements urged as per Grant, et al.	9
De Smet/ Belgium	Vitamin D deficiency as risk factor for severe COVID-19: a convergence of two pandemics(219)	186 Covid+ vs 2717 /R/ 23 pages	D,P,R,C,	Deficiency seems to cause severe, supplements urged	9
Fasano/ Italy, Canada	COVID-19 in Parkinson's Disease Patients Living in ... Italy(214)	105 Covid+ vs 1381/S/ 12 pages	G,C	Parkinson's doesn't matter; D does. Conduct D studies	7
Faul/ Ireland	Vitamin D Deficiency and ARDS after SARS-CoV-2 Infection(213)	33 Covid+ S/ 2 pages	D,C	Worse outcomes in men with lower D	7
Fox/ USA	No Association ...Between Vitamin D Deficiency and COVID-19 Infection, Hospitalization, or Mortality(83)	28,185 patients 3 pages	None mentioned	No difference with deficiency, but authors call for better studies	0
Ghasemian/ Iran, Australia, Russia,	The Role of Vitamin D in The Age of COVID-19: A Systematic Review and Meta-Analysis Along with an Ecological Approach(68)	9 studies (subset:6) 51 countries 20 pages	G,P,R,C,	Patients with COVID-19 lacked D, slight country link Need RCTs	6
Glicio/ India	Vitamin D Level of	176 Covid+	P,R,C,	Link is	6

	Mild and Severe Elderly Cases of COVID-19...(47)	18 pages		suggested Supplement elderly	
Hastie/ UK	Vitamin D concentrations and COVID-19 infection in UK Biobank(10)	449 Covid+ vs 348,149 R/5 pages	G,P,	Link attributed to confounds – D has no role in COVID-19	0
Horowitz/ USA	Efficacy of glutathione ..dyspnea ... COVID-19 pneumonia ... 2 cases(210)	2 (1 D) 8 pages	None mentioned	One patient had D deficiency	5
Ilie/UK	The role of vitamin D in the prevention of coronavirus disease 2019 infection and mortality(254)	20 countries /P/ 4 pages	D,G,P,R,C,S	Recommend study comparing D levels & COVID-19 severity	6
Kara/Italy,Turkey, Taiwan	‘Scientific Strabismus’ or Two Related Pandemics COVID-19 & Vitamin D Deficiency(255)	40 countries /P/ 6 pages	D,G,P,R,S	Goal 40-60ng/ml 10,000IU/day x1mo then 5000IU/day	10
Kohlmeier/ USA	Avoidance of vitamin D deficiency to slow the COVID-19 pandemic(204)	22 states/A/ 7 pages	D,G,P,R,C	Precision nutrition RDA - <4000IU/Day	8
Kumar/ India	Spurious Correlation? A review of the relationship ... Vitamin D and COVID-19 infection & mortality(197)	20 countries /P/ 7 pages	None mentioned	Ilie is wrong – no role for vitamin D Overdose is likely	0
Laird/ UK, Ireland	Vitamin D and Inflammation: Potential Implications for Severity of COVID-19(37)	12 countries /P/ 7 pages	D,P,G,R,C,S	Correlation exists Publicize D recommendations	6
Lau/USA	Vitamin D insufficiency is prevalent in severe COVID-19(155)	20 Covid+ R/14 pages	G,P,R,C,T	Worse with low D Study D further	6

Li/ China, USA	Identifying novel factors ... COVID-19 using ...machine learning(203)	A (machine) 32 pages	Not discussed	D is independent risk factor for cases	7
Li/USA	Multivariate Analysis of Factors Affecting COVID-19 ...U.S. Counties: ...Black Race and Temperature(195)	661 & 217 counties A/ 21 pages	D,P,R,C	D is a likely factor Conducting studies Supplement all	7
Li/USA, China	Sunlight and vitamin D in ... (COVID-19) infection and mortality ... (207)	49 states A/ 15 pages	D,G,P,R,C,	Appears related Need more studies	6
Marik/ USA	Does vitamin D status impact mortality from SARS-CoV-2 ...? (208)	50 states A/ 2 pages	G,P,R,C	Appears related Standard dosages	6
Meltzer/ USA	Association of Vitamin D Deficiency ... Treatment with COVID-19 ... (220)	499 patients R/ 22 pages	D,G,P,R,C	D decreases risk 4000-5000IU/day	10
Moozhipurath/ Germany	Evidence of Protective Role of Ultraviolet-B (UVB) Radiation in Reducing COVID-19 Deaths(21)	152 affected countries/ A/ 42 pages	R,C,F	D decreases deaths Encourage time in sun for deficient	6
Notari/ Spain, Brazil	COVID-19 transmission risk factors(199)	126 countries, 24 factors A/ 42 pages	R	Independent risk: Type 1 diabetes, BCG vaccination, and vitamin D	7
Panarese/ Italy	Letter: COVID-19, and vitamin D (response to Tian, et al. Gastro)(196)	108countries A/3 pages	G,P,R,C,S	Immune control 2000IU/day	8
Pinzon/ Indonesia	Vitamin D Deficiency ... Patients with COVID-19 : Case Series and Recent	10 patients CS 9 pages	D,G,P,R,C	Immune benefits They give all patients 2000IU	8

	Literature Review(212)				
Raharusuna/ Indonesia	Patterns of COVID-19 Mortality and Vitamin D: An Indonesian Study(46)	780 patients R 14 pages	R,C	D is strongly associated with death – do RCTs	8
Raisi-Estabragh/ UK	Greater risk of severe COVID-19 in non-White ethnicities is not explained by ...vitamin D status: study of 1,326 ... UK Biobank(75)	Hospitalized 1326 + 3184 - R/ 21 pages	Not discussed	Ruled out cardio-metabolic, social, behavior, and D – leaving genetic?	0
Rhodes/ UK	Editorial: low population mortality ...south of latitude 35 degrees North supports vitamin D ...(256)	120 countries A/ 4 pages	D,P,C,	Could protect from cytokine storm 1000IU/day	7
Singh/ India	Revisiting the role of vitamin D ... prevention and mortality in European ...post peak(198)	20 countries 2 dates A/ 8 pages	R,C,S	D influenced cases more than deaths study ? of giving D	3
Skutsch/Netherlands, Mexico, Germany	The association of UV with rates of COVID-19 transmission and deaths in Mexico: the possible mediating role of vitamin D(206)	45 cities in Mexico A/ 29 pages	D,P,R,C,	Further study of link between UV (vitamin D) and transmission rates	6
Sun/ China	Serum calcium as a biomarker of clinical severity and prognosis...(217)	241 patients P/18 pages	not discussed	Deficiency suggests larger studies	7
Tan/ Singapore	A cohort study to evaluate the effect of combination Vitamin D, Magnesium and	43 patients 26 controls 17	R,C	Give combination earlier to all cases and to high risk	9

	Vitamin B12 (DMB) on progression to severe outcome in older COVID-19 patients(215)	supplement R/ 6 pages		contacts	
Yao/ China	No Association of COVID-19 transmission with temperature or UV radiation in Chinese cities(193)	224 cities in China/ G/ 9 pages	G,R	UV (therefore D) not related to transmission	0

**\*Type of Study**

A = Analyses of population data or latitude data (geographic)

C = Causal inference modeling report, Hill’s methodology for exploring causality, etc.

P = Prospective correlational study

R = Retrospective chart (or data) review

CS = Single or Multiple Case Study

**\*\* Supporting Information provided by these authors:**

G = geographical observations (not research) of relationship between low vitamin D and high COVID-19

I = Irradiance in the geographic area influences vitamin D deficiency

Biological plausibility:

R = vitamin D enhances resistance to respiratory viruses, decreasing incidence of infection

F = vitamin D decreases overall fatalities from respiratory viruses

C = vitamin D suppresses cytokines that are implicated in severe COVID-19

J = vitamin D tightens junctions, helping prevent viral infections from progressing to pneumonia

T = vitamin D decreases the risk of thrombosis

S = vitamin D suppresses the ‘Renin-Angiotensin’ System activity, which is more of a problem for males

Note: Content from an earlier preprint by Bäcker, Follow the Sun: Slower COVID-19 morbidity and mortality growth at higher irradiances, <https://ssrn.com/abstract=3567587>, last revised 13 April 2020, appears to have been divided into the two much more complete and more recently revised articles included here in Table 3. Similarly, Bäcker’s preprint, Non-black Latinos Spend More Time Outdoors and Have Lower COVID-19 Morbidity and Case Fatality Rates Than Non-Latinos, last revised April 16, appears to have been completely incorporated into a similar work, Non-black Hispanic/Latinos Spend More Time Outdoors and Have Lower COVID-19 Morbidity Than Non-Hispanic/Latinos, Last revised April 20, 2020.



## 10.2 Table 3

<b>Table 3: In Vitro Studies and Brief Perspectives with Summaries of Pertinent Previously Published Data (141 entries)</b>			
1st Author /Location	Title (Truncated)	Supporting Information* /# of pages	Recommendation for COVID-19 & D (dose-who-why)
Adams/ USA	Myth Busters: Dietary Supplements and COVID-19(123)	R,J,C 7 pages	600-4000IU/day – all adults - R
Alabada/ Iraq	Vitamin D effectiveness and pathology in humans and domestic animals(257)	C 15 pages	None - COVID-19 barely mentioned
Álvarez-López /Mexico	Vitamin D y la pandemia por COVID-19 ( Vitamin D and the pandemic of COVID-19)(258)	P,R,F,J,C 6 pages	10- 25 mcg/day - high risk – D, Goal = 30 ng/ml
Anderson/ UK & USA	COVID-19 Pathophysiology: Interactions of Gut Microbiome, Melatonin, Vitamin D, Stress, Kynurenine... Nicotinic ... Treatment Implications(259)	P,R,F,J,C,S 24 pages	Vitamin D supplements - no details provided
Annweiler /France & China	Counter-regulatory ‘Renin-Angiotensin’ System-... Drugs to Treat COVID-19 ...infected patients(181)	F,C,S 2 pages	RAS regulators to treat COVID-19 - no vitamin D mention
Alpalhão/ Portugal	SARS-CoV-2 pandemic and Vitamin D deficiency—A double trouble(260)	D,R,F 2 pages	20-40 min sun/day; 10-50mcg/day; test (more if deficient)
Arboleda-Alzate/ Columbia	Vitamin D supplementation: a potential approach for COVID-19 therapeutics(149)	G,R,F,C,S 9 pages	“Conventional” oral vitamin D supplementation
Ardiaria/ Indonesia	Peran vitamin d dalam pencegahan influenza dan COVID-19(150)	P,R,C,J 7 pages	Clinical trials are needed urgently
Arya/ India	Synergistic effect of Vitamin D and Remdesivir can fight COVID-19(151)	R,C, 4 pages	D plus Remdesivir should be trialed
Aslan/ Turkey	Is vitamin D one of the key elements in COVID-19 days?(152)	D,G,P,R,C,S 2 pages	Supplement entire population
Aygun/ Turkey	Vitamin D can prevent COVID-19 infection-induced multiple organ damage(261)	G,R,C,S 4 pages	Supplement due to R,F
Bakare/ UK	A cost-effective preventative approach to ...save lives ...using Vitamin D, Curcumin, ... Vitamin C(89)	R,C, 7 pages	Supplement with no added calcium
Barazzoni/ Italy, Israel, Germany, Denmark, Russia, Croatia	ESPEN expert statements and practical guidance for nutritional management of individuals with SARS-CoV-2 infection(262)	R 9 pages	Screen at-risk populations and patients for malnutrition, including vitamin D, and supplement accordingly
Barron/ UK	Vitamin D: A rapid review of the evidence for treatment or prevention of COVID-19 in adults	P,R 10 pages	Those at risk of deficiency should

	(excluding pregnant & breastfeeding women)(34)		take 400IU/day
Bauer/ USA	What is the role of supplementation with ascorbic acid, zinc, vitamin D, or N-acetylcysteine for prevention or treatment of COVID-19?(263)	D,P,R,C, 3 pages	No known risks, but recommend D only after RCTs
Belančić /Croatia	Potential pathophysiological mechanisms leading to increased COVID-19 ... in obesity(264)	D,P,R,C, 3 pages	Measure D and supplement obese
Biesalski/ Germany	Vitamin D deficiency and co-morbidities in COVID-19 patients – A fatal relationship?(166)	D,G,P,R,F,C,J, T,S 13 pages	D prevents storm Supplement
Bloukh/ UAE, India	Prevalence of COVID-19: A Look behind the Scenes(2)	D,P,R,C 18 pages	Prophylactically supplement if low
Brown/ UK, India	Vitamin D deficiency: a factor in COVID-19 progression severity and mortality? An urgent call for research(110,265)	D,G,P,R,F,C 29 pages	Urgent call for research, possible supplementation
Brown / UK	Rapid Response: Re: Preventing a COVID-19 pandemic - COVID-19: Vitamin D deficiency; and, death rates; ... connection? Research urgently...(7)	D,G,P 4 pages	Urgent call for research, possible supplementation
Brown/ UK, India, Philippine s, Austria, France, Australia, Ireland, Canada, Italy, USA, Sweden, Greece, Turkey	COVID-19 ICU risk – 20-fold greater in the Vitamin D Deficient... Sun and D-supplementation – Game-changers? Research urgently required(266)	P,R,C,J,S 6 pages	Urgent call for research, correct deficiencies
Buttriss/ UK	Is a vitamin D fortification strategy needed?(88)	D,R 8 pages	400IU/day due to staying indoors
Caccialanza/ Italy	Early nutritional supplementation in non-critically ill patients hospitalized for ...(COVID-19): protocol(48)	R,C 5 pages	Correct deficiency aggressively
Calder/UK, USA, New Zealand, The Netherlands	Optimal Nutritional Status for a Well-Functioning Immune System Is an Important Factor to Protect against Viral Infections(49)	D,R, 10 pages	All adults should take 2000 IU of vitamin D/day, US National Academy recommendations
Cao/China , France	SARS-CoV-2 & COVID-19: Key-Roles of the ‘Renin-Angiotensin’ System / Vitamin D Impacting ...(22)	P,R,F,C,S 2 pages	Take note: vitamin D is crucial
Carter /USA	Considerations for Obesity, Vitamin D, and Physical Activity Amid the COVID-19 Pandemic(111)	P,R,C,S 2 pages	Encourage outdoor activities, give D
Chakhtour a/Italy, USA,	Commentary: Myths and facts on vitamin D amidst the COVID-19 pandemic(107)	G,D,R,C, 3 pages	Supplement with 1000-4000IU/day to achieve

Lebanon			>30ng/ml 25(OH)D
Chan/ Hong Kong	What can we do for the Personal Protection against the COVID-19 infection? Immuno-boosting(267)	R,C 6 pages	Supplement with vitamin D, herbals
Chhetri/ China, Indonesia, Japan, Korea, Thailand, Sri Lanka	Prevention of COVID-19 in older adults: A brief guidance from The International Association for Gerontology and Geriatrics (IAGG) Asia/Oceania Region(52)	R 2 pages	Get enough sunlight “in the morning” for vitamin D production
Crane- Godreau/ USA	Vitamin D Deficiency and Air Pollution Exacerbate COVID-19 Through Suppression of Antiviral Peptide LL37(43)	D,G,P,R,F,C, 5 pages	Deficiency + air pollution is trouble Daily dose is best
Dasgupta / India	Nsp7 and Spike Glycoprotein of SARS-CoV-2 ...Potential Targets of Vitamin D and Ivermectin(222)	R,C, 13 pages	D may dock onto virus, be curative
De Lucena /Brazil	Mechanism of inflammatory response in associated comorbidities in COVID-19(158)	R,C,S 4 pages	D can decrease complications
Dhillon/ UK	COVID-19 breakthroughs: separating fact from fiction(268)	P,R, 37 pages	Evidence is weak But, supplement
Docea/ USA, Greece, Romania, Russia	A new threat from an old enemy: Re-emergence of coronavirus (Review)(236)	R,C 13 pages	Evidence is weak, but biologically plausible Try D for prevention and treatment
Ebadi/ Canada	Perspective: improving vitamin D status in the management of COVID-19(9)	G,P,R,C,S 4 pages	Very plausible High doses for pts.
Ekiz/ Turkey	Relationship between COVID-19 and obesity(188)	P,R,C 3 pages	Plausible link Supplement
Ekiz/ Turkey	Revisiting vitamin D and ... exercises for patients with sleep apnea ... the COVID-19 quarantine(190)	P,R 6 pages	10,000IU/day x1mo then 5000IU/day
Fabbri/US A, Italy	Editorial – Vitamin D status: a key modulator of innate immunity and natural defense ...(41)	D,R,C 5 pages	Up to 6000IU/day Goal: 30 - 60ng/ml
Facchiano /Italy, USA	Reply to Jakovac: About COVID-19 and vitamin D (In support of Jakovac)(192)	R,C,S 1 page	No doubt Conduct trials, give
Farias/ Brazil	Vitamin D as an Immunological Factor in Combating COVID-19(269)	D,P,R,C 10 pages	Immune benefits Eat fish and eggs
Froum/ USA	Vitamin D and COVID 19 disease severity: An old ally in a new war(170)	R,C,J, 5 pages	Dentists should test, supplement
Garami/S witzerland	Re: Preventing a COVID-19 pandemic - Is there a magic bullet to save patients?... give it a try!(121)	D,G,C,S 3 pages	Biological plausibility, give D!
Garg/UK, Australia	Editorial: low population mortality from COVID-19 ... latitude 35 degrees North—supports vitamin D as a factor determining severity. Authors' reply(28)	D – denies others 2 pages	Premature, but for musculoskeletal health, give D

Gasmi/ Iran, Pakistan, Norway, Thailand, Luxembou rg, France	Individual risk management strategy and potential therapeutic options for the COVID-19 pandemic(270)	D,P,R 9 pages	vitamin D prevents infections when given prior to exposure Goal 40-60ng/ml 10,000IU/day x1mo then 5000IU/day
Gee/UK	Management of clozapine treatment during the COVID-19 pandemic(237)	D,P,R,C 10 pages	D decreases risk Supplement
Ghavidel- darestani/ Iran	Role of vitamin D in pathogenesis and severity of COVID-19 infection(44)	D,P,R,C,S 16 pages	Biological plausibility D for all deficient
Glinsky/ USA	Tripartite Combination of ... Pandemic Mitigation Agents: Vitamin D, Quercetin, and Estradiol(271)	D,R,C,S 26 pages	The 3 listed agents should be given
González/ Puerto Rico	2020 COVID-19 Protocol(272)	R,F 9 pages	Support immune system: 10,000IU/d Infected: 25,000IU
Grant/ USA, Hungary	Evidence that Vitamin D Supplementation Could Reduce Risk of Influenza and COVID-19 Infections And Deaths(273)	R,F,C,S (full review) 19 pages	Evidence is strong Goal:40-60ng/ml 2000-5000IU+ daily
Grant/ USA, Hastie/UK	Letter in response to the article: Vitamin D concentrations and COVID-19 infection in UK biobank (Hastie et al.) and Response to letter(82)	Debate - see section 1.1 2 pages	Researchers of differing views fail to communicate
Grant/USA	Reply: "Vitamin D Supplementation in Influenza and COVID-19 Infections. Comment on: Evidence ..."(103)	See section 1.2, 5 pages	Response to Kow softened D dose
Guan/USA , China, Australia, Brazil, UK, Korea, South Africa, Sweden, Morocco, Russia, Mexico, Canada, Netherlan ds, Chile	Promoting healthy movement behaviours among children during the COVID-19 pandemic(274)	Vitamin D mentioned in passing only, assumed to be beneficial 3 pages	Children need to play outdoors
Gupta/ India	Is Immuno-modulation the Key to COVID-19 Pandemic?(275)	R,C 4 pages	D helps immunity BCG vaccine
Hamad/ Sudan	Vitamin D Supplements Improve Efficacy of Minocycline...Triple Therapy to COVID-19 ...(153)	R,C 2 pages	Improves immune Add D to treatment

Hasan/Saudi Arabia, Qatar, UAE, Iran, Iraq	A review on the cleavage priming of the spike protein on coronavirus by angiotensin-converting enzyme-2 and furin (276)	S 9 pages	D can inhibit ACE-2 therefore could be therapeutic
Heiser/USA	Identification of potential treatments for COVID-19 through artificial intelligence-enabled ...analysis of human cells infected with SARS-CoV-2(277)	P,C 15 pages	Weak effect in D analogues – study for new drugs
Hossain/Bangladesh	Natural Polyphenol ... Crops based Diet to Promote Innate Immunity to Combat COVID-19 Disease(278)	R,C 22 pages	Polyphenols + D enhance immunity
Hribar/UK, USA	Potential Role of Vitamin D in the Elderly to Resist COVID-19 and to Slow ... Parkinson's Disease(38)	D,P,R,F,C,J,S? 8 pages	Enhances immunity 2000-5000IU/day
Isaia / Italy	Possible prevention and therapeutic role of vitamin D in ... the COVID-19 2020 pandemics(54)	D,G,P,R,C,J,	Enhances immunity Lit. review only
Islam /Germany	A transcriptome analysis identifies potential preventive 1 and therapeutic 2 approaches ...(117)	C, 21 Pages	Improves immune Give D + ketogenic
Jakovac /Croatia	COVID-19 and vitamin D—Is there a link and an opportunity for intervention?(113)	R,C 1 page	D link is intriguing D now – no harm
Jayawardena/Greece Sri Lanka, Australia	Enhancing Immunity in Viral Infections, with special emphasis on COVID-19: A Review(119)	D,R,C, 59 pages	supplementation of vitamins for COVID-19 is appropriate, especially D 5000IU
Jukic/ Croatia, Spain,	Strategies and Solutions for Team Sports Athletes in Isolation due to COVID-19(279)	No details 9 pages	Educate about nutrition Give D
Kakodkar/ Ireland	A Comprehensive Literature Review on the Clinical Presentation, and Management ... (COVID-19)(108)	R, 18 pages	Prophylaxis and treatment
Kalantar-Zadeh/ USA	Impact of Nutrition and Diet on COVID-19 Infection ..Implications for Kidney Health ...(280)	No details 4 pages	Improves overall health
Kamani/ Iran	Effect of Low Level Yellow Laser Light 589nm on Virus Prevention Corona Virus (COVID-19)(281)	R 4 pages	Use their device to increase D
Khalil/Bangladesh	Sub-continental Atmosphere and Inherent Immune System ... Impact ...COVID-19... South East Asia(282)	R 9 pages	Influences immune Get vitamin D (sun)
Khunti/UK, India	Is ethnicity linked to incidence or outcomes of COVID-19?(191)	P 3 pages	D one of many possible reasons
Koenig/ USA, Iran, China, Saudi Arabia	Ways of Protecting Religious Older Adults from the Consequences of COVID-19(230)	P,R 9 pages	D is often deficient in elderly, needed for immunity Take supplements
Kow/ UK, Malaysia	Vitamin D ...in Influenza and COVID-19 Infections Comment on: “Evidence that Vitamin D ...”(101)	Rationale OK 2 pages	High dose lacks evidence – trials 1 <sup>st</sup>
Kumar/	Letter: does vitamin D have a potential role against	R,C,S	D has two roles

India	COVID-19? (Comment on Tian, et al.)(283)	2 pages	
Kurthkoti/ India	Potential benefits of Vitamin D supplementation in COVID-19 patients. A Hypothesis(238)	G,R,C 4 pages	Influences immune Supplement all
La Vignera / Italy	Sex-Specific SARS-CoV-2 Mortality: Among Hormone-Modulated ACE2 Expression, Risk of ... Thromboembolism and Hypovitaminosis D(51)	D,P,C,T,S 6 pages	D explains high male mortality & thrombosis
Lanham-New/ USA, UK	Vitamin D and SARS-CoV-2 virus/COVID-19 disease(223)	D,P,R, 5 pages	D is important, but risks are great RDA or max 4000IU
Lawton/ USA	Will warmer spring weather slow down the rate of spread?(284)	G,R 1 page	D helps immunity
Lee /UK	Vitamin D:A rapid review of the evidence for treatment or prevention in COVID-19(29)	D,G,P,R 10 pages	No evidence found Take RDA
Manson/ USA	Does Vitamin D Protect Against COVID-19?(63)	R,P,C, 1 page	Strong evidence 1000-2000IU
Mansur/ USA	Letter: low ...mortality from COVID-19 in countries south of latitude...vitamin D ...(reply to Rhodes) (124)	R,C,S 1 page	Additional info Supports vitamin D
Martín Giménez/ Argentina	Lungs as target of COVID-19 infection: Protective ... molecular mechanisms of vitamin D and melatonin ... new potential synergistic treatment(285)	D,P,R,J,C,S 8 pages	Detailed biological plausibility data D + melatonin
Maruta/ Australia	PAK1-blockers: Potential Therapeutics against COVID-19(286)	S 6 pages	Use MART-10, a derivative with D3
McCartney / Ireland	Optimisation of Vitamin D Status for Enhanced Immuno-protection Against COVID-19(73)	D,P,R,C 4 pages	800-2000IU/day - all adults - R
McCartney / Ireland	Authors of Article 'Optimisation of Vitamin D Status ... comment on response ... by McKenna et al(287)	D,P, 5 pages	800-2000IU/day - all adults - R
McCartney / Ireland	Authors of Article 'Optimisation of Vitamin D Status ... comment on response ... by Rabbitt et al(250)	D,R, 3 pages	Biological Plausibility
McCullough/ USA	The Essential Role of Vitamin D in the Biosynthesis ...Antimicrobial Peptides May Explain ...Deficiency Increases Mortality Risk in COVID-19 Infections(97)	G,P,R 23 pages	A plethora of data from other illness 10,000IU+/day
McKenna/ Ireland	COVID-19, Cocooning and Vitamin D Intake Requirements(71)	D,P,R 6 pages	NOT 800-2000IU!!! Goal: 10ng/ml
Meftahi/ Iran	The possible pathophysiology mechanism of cytokine storm in elderly adults with COVID-19 infection: the contribution of "inflamm-aging"(165)	P,R,F,C,S 15 pages	Deficiency is a likely cause of cytokine storm
Merow/ USA	Seasonality and uncertainty in COVID-19 growth rates(288)	R 27 pages	Increases resistance
Misra/ UK, India, Ukraine	Rheumatologists' perspective on coronavirus disease 19 (COVID-19) and potential therapeutic targets(114)	R,F,C 8 pages	Prevent & mitigate High dose to correct if <50ng/ml
Mitchell/ UK	Vitamin-D and COVID-19: do deficient risk a poorer outcome?(67)	R,C,S 1 page	Influences immune Correct deficiency

Molloy/ Ireland	Vitamin D, COVID-19 and Children(72)	D,G,P,R,C 3 pages	Influences immune Correct deficiency
Muscogiuri / Italy	Nutritional recommendations for COVID-19 quarantine(187)	D,R,J,C 2 pages	Change diet and eat D-rich foods
Muscogiuri / Italy	Obesity: the “Achilles heel” for COVID-19?(173) (obesity research study, vitamin D in discussion)	D,P,R,C 12 pages	Obesity-related D deficiency fatal?
Musselwhite / UK	JTH editorial v17 – The importance of psychosocial factors in transport and health(289)	No mention 2 pages	Higher D with public transport
Nigro/ Italy, China, UK, Denmark	Molecular mechanisms involved in the positive effects of physical activity on coping with COVID-19(253)	P,R,J,C, 32 pages	Encourage physical activity, following guidelines. This will increase D
Nikhra / India	Identifying Patterns in COVID-19: Morbidity, Recovery, and the Aftermath(235)	P,R,F 9 pages	D supplementation may help prevent
Nizami/ India	Strong Immunity- A Major Weapon to Fight against COVID-19(225)	R, 8 pages	Correct deficiency for immune health
Pal / India	Managing common endocrine disorders amid COVID-19 pandemic(224)	G,P,R,C,F,S 5 pages	Speculation; Bone health: 800-1000IU
Palmer / Italy, UK, Sweden, Finland, Spain, Switzerland, France, Greece	The potential long-term impact of the COVID-19 outbreak on patients with non-communicable diseases in Europe: consequences for healthy ageing(290)	P,R, 6 pages	D deficiency is a concern; immune
Parvin/Bangladesh	The symptoms, contagious process, prevention and post treatment of COVID-19(226)	P 25 pages	Sunlight and daily supplements
Pinnock/ UK	Vitamin D and COVID-19. The Evidence Warrants Discussion!(26)	G,R,C 5 pages	Immune benefits 1000-2000IU
Queiroz / Brazil	Management of inflammatory bowel disease patients ...COVID-19 pandemic...: a Brazilian ... (234)	R 6 pages	Immune benefits Always supplement
Quesada-Gomez/ Spain, Belgium	Vitamin D Receptor stimulation to reduce Acute Respiratory Distress Syndrome (ARDS) in patients with Coronavirus SARS-CoV-2 infections(106)	D,P,R,C,T,S, 23 pages	Prevents infection Controls immune Calcifediol is best absorbed form
Rabbitt/ Ireland	Vitamin D and COVID-19: A Note of Caution (response to McCartney, Optimisation of ... D)(291)	(none) 2 pages	D is safe but need RCTs first
Ramos/ Brazil	COVID-19, Rate of Case Factors and Nutritional Characteristics of ...Dying in Italy and Brazil: ... (239)	D,P,R,C,F, 9 pages	Immune benefits 50,000IU/week
Raymond/ UK	Mast cell stabilisers, leukotriene antagonists and antihistamines: ... evidence ... use in COVID-19(53)	R,C 6 pages	No evidence of a D COVID-19 link
Razdan / India	Vitamin D Levels and COVID-19 Susceptibility: Is there any Correlation?(240)	D,R,C 3 pages	Clear COVID-19 link



			2000-4000IU/day
Rhodes/ UK	Letter: low population mortality from COVID-19 ...south of latitude 35° North supports vitamin D as a factor ...severity—authors' reply (to Mansur)(125)	S,T 2 pages	Circumstantial evidence is strong South: supplement
Ribeiro/ Brazil, UK, Australia	Does Vitamin D play a role in the management of COVID-19 in Brazil? (Building upon Isaia literature review)(55)	D,P,R, 6 pages	Enhances Immunity Get more sun. Do not take 10,000IU
Richards/ South Africa	COVID-19 and the Rationale for Pharmacotherapy: A South African Perspective(227)	S 7 pages	50,000IU stat for acute treatment 4000IU/day if mild
Richer/ USA	How to build a lifestyle and nutritional firewall against viruses like COVID 19(292)	D,P,R 5 pages	Enhances Immunity D in multivitamin
Rocha/UK, Portugal	Reply to Jakovac; Severity of COVID-19 infection in patients with phenylketonuria: is vitamin D status protective? (In support of Jakovac)(293)	P,R 2 pages	Enhances Immunity PKU supplement Supplement all
Romano / Italy	Short Report – Medical nutrition therapy for critically ill patients with COVID-19(294)	(none) 5 pages	Correct <12.5ng/ml (IV cholecalciferol)
Roy/ UK	Response to 'Vitamin D concentrations and COVID-19 infection in UK Biobank' (Hastie)(77)	R,F,J, 1 page	Enhances Immunity Better research!
Rusciano/ Italy	The Fight against COVID-19: The Role of Drugs and Food Supplements(295)	D,P,R,J,C 16 pages	Enhances Immunity eat well
Saul/ USA	Nutritional Treatment of Coronavirus(241)	(none) 9 pages	5000IU/day x 2 weeks then 2000IU
Sharma/ India	Vitamin D: A cheap yet effective bullet against coronavirus disease-19 – Are we convinced yet?(105)	D,G,P,R,F,C,J, S 8 pages	Evidence is strong 1000-2000IU/day
Siddiqui/ UAE	Centralized air-conditioning and transmission of novel coronavirus(252)	D,R 3 pages	Air conditioning limits outdoor time
Silberstein / Australia	Vitamin D: A simpler alternative to tocilizumab for trial in COVID-19?(296)	D,G,P,R,C 2 pages	Biological plausibility – study!
Silberstein / Australia	Premorbid IL-6 levels may predict mortality from COVID-19 (studied IL-6 levels, not vitamin D) (169)	P,C 9 pages	Correlation strong Supplement all
Siuka/ Slovenia	Vitamin D supplementation in the COVID-19 pandemic(242)	D,P,R,F,C 3 pages	Supplement high-risk, urgently
Sparavigna/ Italy	Vitamin D for COVID-19?(164)	D,R,C, 8 pages	Lists literature - no conclusion drawn
Speeckaert/ Belgium	Association between low vitamin D and COVID-19: don't forget the vitamin D binding protein(171)	C,T 2 pages	Decreases ARDs Avoid deficiency
Suresh/ India	Hypovitaminosis D and COVID-19: Matter of Concern in India?(233)	D,G,P,R,C,S 2 pages	Study, and correct D deficiency
Taheri/ Qatar	Managing diabetes in Qatar during the COVID-19 pandemic(69)	R 2 pages	D is delivered to diabetic patients
Tan/	Medications in COVID-19 patients: summarizing the	R,C,	Continue, but don't

Singapore	current literature ... orthopaedic perspective(243)	5 pages	add, B, C, and D
Thomson/ UK	The COVID-19 Pandemic: A Global Natural Experiment(297)	(none) 6 pages	Seasons allow study of D impact
Tian/ China	Letter: COVID-19, and vitamin D. Authors' reply (to Panarese)(112)	G,R,J,C,T,S 2 pages	Theory is sound Supplement all
Tian/ China	Letter: does vitamin D have a potential role against COVID-19? Authors' reply (to Kumar)(244)	R,C,T, 4 pages	Theory is sound Moderate doses
Trinity College Dublin/ Ireland	News Release 12-May-2020 Vitamin D determines severity in COVID-19 so government advice needs to change(27)	D,G,P,F,C 3 pages	Irish government should support supplements and food fortification
Turashvili/ Georgia	Could vitamin D reduce the risk of COVID-19?(115)	G,P,R,C,S 5 pages	Biological plausibility – study!
Uçar/ Turkey	Koronavirüs ve fitoterapi(231)	R,C, 9 pages	Boosts immunity Give supplements
Wang/ China	Review of the 2019 novel coronavirus (SARS-CoV-2) based on current evidence(1)	R 7 pages	Boosts immunity Give supplements
Weir/ USA	Does vitamin D deficiency increase the severity of COVID-19?(167)	D,P,R,C,T 2 pages	Boosts immunity Give modest doses
Wimalawansa/ USA	Global epidemic of coronavirus—COVID-19: What can we do to minimize risks(228)	D,R,C, 7 pages	Loading dose for all 200,000-300,000IU
Yong/ Malaysia	Population Studies Confirm Risk Factors for Catching COVID-19(298) and Vitamin D as an Independent Risk Factor for COVID-19 Death(299)	F 5 & 10 pages	Studies are compelling
Yousfi/ Tunisia, Italy, Poland, Canada, Qatar	The COVID-19 pandemic: how to maintain a healthy immune ... special focus on athletes.(245)	P,R,C,F 7 pages	10,000IU/day x1mo then 5000IU/day
Zabetakis/ Ireland, USA	COVID-19: The Inflammation Link and the Role of Nutrition in Potential Mitigation(246)	P,R 28 pages	Uncertain if high doses required, but supplement
Zemb/ France, Sweden, Belgium, Italy, USA, Austria, Poland, Romania	Vitamin D deficiency and COVID-19 pandemic(247)	D,P,R, 5 pages	Immune benefits for other infections are known, but not for COVID-19. No harm found with 2000-4000IU, so why not supplement?
Zhang/ Japan, China	Current status of potential therapeutic candidates for the COVID-19 crisis(126)	R,C	Studies including D are underway (not yet recommended)

Zhang/ China	Potential interventions for novel coronavirus in China: A systematic review(118)	D,G,P,R	Could work to prevent
Zicarelli/ Canada	Nutrition Therapy ... (COVID-19): Recommendations ... for Integrative Medical Treatments(300)	R,C, 9 pages	Benefits patients 2000IU/day

\* Supporting Information provided by these authors:

D = vitamin D deficiency is common

G = geographical observations (not data) of relationship between low vitamin D and high COVID-19

P = deficiency is more common in populations at higher risk for severe COVID-19 complications

Biological plausibility:

R = vitamin D enhances resistance to respiratory viruses, decreasing incidence of infection

F = vitamin D decreases overall fatalities from respiratory viruses; decreases COVID-19 severity

C = vitamin D suppresses pro-inflammatory cytokines implicated in severe COVID-19

J = vitamin D tightens junctions, helping prevent viral infections from progressing to pneumonia

T = vitamin D decreases the risk of thrombosis

S = vitamin D suppresses the 'Renin-Angiotensin' System activity, which is more of a problem for males

# of pages includes references, figures, and tables

### 10.3 The Irish Medical Journal Debate

McKenna and Flynn began the debate by objecting to McCartney et al.'s recommendation of 800IU-2000IU/day of vitamin D, stating that there is no randomized controlled trial (RCT) evidence to support such high doses, which can be harmful, particularly in children.(71,73) They cited early evidence from TILDA suggesting that 800IU/day is adequate for frail elderly and healthy elderly require only 400IU/day, and that only in the winter or when "cocooning."(71) They also cited Molloy, et al., in asserting that infants should not be given more than 200IU/day, expressing concern that such high doses will result in hypercalcemia in children, falls and fractures in the elderly, and the public may conclude that supplementation with 2000IU/day long-term is safe.(71) McKenna and Flynn advised against sunlight exposure due to skin cancer risks, and concluded by asserting that most adults obtain all the vitamin D they need (with 10ng/ml as their target serum level) from fortified foods, even with "cocooning," but conceded that supplementation with 200-800IU/day would be acceptable for those not eating fortified foods.(71) The authors provided a chart to guide dosages for three groups of adults.(71) McKenna and Flynn did not, however, dispute a link between vitamin D and COVID-19.(71)

McCartney and Byrne responded to McKenna and Flynn, pointing out that they submitted an early draft to the Irish Department of Health prior to publication, and the article was written to guide health care professionals, not the general public.(287) Also, the TILDA researchers now recommend 800 – 1000IU of vitamin D per day for adults over aged 70, and nursing home residents, who are the most frail of all (with a 50% COVID-19 case fatality rate in Ireland), were the target group of their article.(287) Supplementation for children was not discussed in the article.(70,287) They contended that concerns for vitamin D toxicity were overstated based upon the evidence, and pointed out that the 25(OH)D (serum vitamin D) levels required to maintain bone health are far lower than the amounts that have been shown to decrease risk of viral

respiratory infection.(287) They concluded by citing a publication, “from one of the world’s foremost authorities” recommending 2000IU/day for optimal protection against viral infection.(287)

Next, Laird, et al., presented a graph illustrating the correlation ( $p = 0.046$ ) between low population mean 25(OH)D levels and higher COVID-19 death rates (study details in Section 3.4.1).(37) They detailed several biological plausibility arguments for vitamin D’s role in COVID-19 outcomes, emphasizing limiting the potentially damaging cytokine response.(37) They found that the risk groups for severe COVID-19 mirror those for vitamin D deficiency.(37) They concluded by stating that RCTs of vitamin D supplementation would be difficult to conduct during the COVID-19 pandemic, because few appropriately informed people would agree to risk getting the placebo.(37)

Rabbitt and Slattery replied to McCartney, et al.’s April article, stating succinctly that immunonutrition is a field fraught with failures, and other vitamin supplements have proven to be harmful in large doses.(291) They agree that vitamin D is safe, but then describe calls for supplementation to help protect the general population from severe COVID-19 as “folly” due to lack of “clear evidence,” referring to Donald Trump.(291)

McCartney and Byrne replied to Rabbitt and Slattery, stating they fail to acknowledge the biological plausibility of vitamin D for COVID-19, which is supported by a very strong body of evidence.(250) They provided summaries of three additional studies, and pointed out that Rabbitt and Slattery conceded that vitamin D is safe; the fact that other choices are not safe is irrelevant.(250)

Faul, et al., concluded the debate with a brief letter supporting the April McCartney article.(213) The strong correlation between the risk factors for increased severity of COVID-19 and vitamin D deficiency led them to prospectively evaluate vitamin D levels in 33 hospitalized COVID-19 pneumonia patients who were all in respiratory failure (study details in section 3.4.4).(213)