

1
2
3
4
5
6 **Article Type: Original Article**

7
8 **Title: Low Vitamin D levels and prognosis in a COVID-19 paediatric population: A**
9
10 **systematic review**

11 Author: Komal Shah^{1*}, Varna VP², Apurvakumar Pandya³, Deepak Saxena⁴

12
13
14 ¹Assistant Professor, Indian Institute of Public Health Gandhinagar - 382042, Gujarat, India

15
16
17 ²MPH Student, Indian Institute of Public Health Gandhinagar - 382042, Gujarat, India

18
19
20
21 ³Economic Evaluation Specialist, Indian Institute of Public Health Gandhinagar - 382042,
22 Gujarat, India

23
24 ⁴Professor, Indian Institute of Public Health Gandhinagar - 382042, Gujarat, India
25
26
27
28
29

30
31 **Dr. Komal Shah (*Corresponding Author)**

32 Assistant Professor,
33 Indian Institute of Public Health - Gandhinagar
34 Opp. Air Force Head Quarters, Nr. Lekawada Bus Stop,
35 Gandhinagar-Chiloda Road, Gandhinagar - 382042
36 Mob: 9924264500
37 Email: kshah@iiphg.org
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56

1
2
3 **Title: Low Vitamin D levels and prognosis in a COVID-19 paediatric population: A**
4 **systematic review**
5
6

7 **Abstract**
8
9

10 **Aim:** We aim to study the relationship between vitamin D level, risk and severity of COVID-19
11 infection in pediatric population through systematic review.
12
13

14 **Methods:** We searched PubMed, CINAHL, EMBASE, Cochrane Library and Google Scholar
15 from December 2019 to June 2021 for retrieving articles studying association between vitamin D
16 deficiency with COVID-19. Qualitative details were synthesized in evidence table and quantitative
17 data was used for deriving pooled estimate through meta-analysis.
18
19

20 **Results:** After initial search of 2261 articles, eight eligible studies (two reviews) were included in
21 the systematic review. Meta-analysis of the quantitative data (six studies) showed pooled
22 prevalence of vitamin D deficiency as 45.91% (95% CI:25.148-67.450). In infected pediatric
23 patients, low levels of vitamin D increased the risk of severe disease (odds ratio - 5.5; 95%
24 CI:1.560- 19.515; p=0.008). It was also found that children and adolescents having vitamin D
25 deficiency had greater risk of COVID infection as compared to patients with normal vitamin D
26 levels. Improvement in disease severity with vitamin D supplementation was also noted.
27
28
29
30
31
32
33
34
35

36 **Conclusion:** The systematic review showed that almost half of the pediatric COVID patients suffer
37 from vitamin D deficiency. It is also clear that the low level of vitamin D is associated with greater
38 risk of infection and poorer outcome in pediatrics.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

The Coronavirus disease of 2019 (COVID-19) - a respiratory illness, was first reported in December 2019, in Wuhan, China and since then it has posed a threat for the health systems across the world. COVID-19, that affects everyone regardless of age, was found to be more severe in patients with low immunity and other underlying comorbidities such as hypertension and diabetes mellitus.^{1,2} The RNA virus with a unique characteristic of antigenic shift has resulted in the inability to find a proper cure. Moreover, the development of the COVID-19 vaccine which has proved to be efficacious would take a long time to cover the whole population. In the case of pediatrics, the vaccine is still under the developmental phase with very few countries initiating vaccination of the population below the age of 18 years. Given the dire circumstances, researchers have focused on other measures that can help prevent COVID-19 spread.

The role of macro and micronutrients in modulating the immune response in diseased and healthy individuals is known for a long. Vitamin D - a fat-soluble vitamin is known to exert a variety of biological effects including immunomodulatory effects and since the emergence of the pandemic, vitamin D is gathering attention from the scientific community.³ A recent systematic review of RCTs showed that Vitamin D supplementation significantly reduces the risk of acute respiratory infection.⁴ These studies laid the foundation of the theory that Vitamin D might have a role to play in COVID-19 susceptibility and disease severity. This was further strengthened with the emerging studies showing the poorer outcome of COVID-19 in patients with vitamin D deficiency.⁵ A significant association between vitamin D, sun exposure, COVID-19 outcomes and severity had also been reported in adults.⁶⁻⁸ One of the recently published meta-analysis showed that vitamin D supplementation reduces the risk of ICU admission in adult COVID patients and it also has a potential impact on reducing the risk of mortality.⁹ Although many studies have demonstrated vitamin D as a protective factor, controversies and heterogeneities exist in the literature showing the effect of vitamin D in the prevention and treatment of COVID-19, especially in the case of the younger population. With this systematic review, we aimed to estimate 1) prevalence of vitamin D deficiency in pediatric COVID patients 2) association between vitamin D deficiency and COVID-19 severity in pediatrics 3) the relationship between vitamin D supplementation and improvement in COVID-19 severity.

Methods

1
2
3 A systematic review as a methodological approach using the framework provided by Arksey &
4 Malley was adopted for the study.¹⁰ For reporting of the review, Preferred Reporting Items for
5 Systematic Review and Meta-Analysis (PRISMA) was used.
6
7

8 9 **Search Strategy**

10
11 We searched five databases – PubMed, CINAHL, EMBASE, Cochrane Library and Google
12 Scholar from December 2019 to June 2021. The search was conducted using index terms (e.g.,
13 MeSH and Entree terms): COVID-19, SARS-CoV-2, COVID-2019, 2019-nCoV, 2019 novel
14 coronavirus infection, coronavirus disease-19, coronavirus disease 2019, severe acute respiratory
15 syndrome coronavirus 2, novel coronavirus, vitamin D, vit D, vitamin D supplementation,
16 children, infant, neonate, young, newborn, baby, babies, pediatric, pediatric, juvenile. These search
17 terms were combined as search sets, using Boolean operators (AND, OR, NOT). Retrieved articles
18 were evaluated using title and abstract and selected articles were further screened using full-text
19 evaluation. Additionally, reference tracking of the searched articles was carried out to identify and
20 other relevant articles, which were missed during the initial search. We excluded duplicate studies
21 from the final search.
22
23

24
25 We included studies published in English in peer-reviewed journals that were conducted in
26 children (age ≤ 18), and those reporting vitamin D status and COVID-19 infection in pediatric
27 patients. Studies reporting the role of vitamin D supplementation in improving disease severity in
28 children and adolescents were also included. We excluded studies including only adult (age >18
29 years) patients and not published in English. The process of inclusion and exclusion is described
30 using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)
31 flowchart (Moher, Liberati, Tetzlaff, & Altman, 2009) (Figure 1).
32
33

34
35 The initial search yielded 2261 articles, of which, after elimination of duplicates and screening of
36 their titles and abstracts and full-text evaluation eight studies were considered relevant to this
37 review.
38
39

40 41 **Data Extraction**

42
43 Two independent reviewers screened the title and abstracts and assigned unique identification
44 numbers to all included articles. Key information from each screened article was extracted by two
45 independent researchers and categorized in an evidence table using the following heads: author(s),
46
47
48
49

1
2
3 title, country, population, study period, total cases, the status of vitamin D, the severity of COVID-
4 19, vitamin D supplementation, outcomes of the study and general findings. Any discrepancy in
5 the coding was resolved by consensus.
6
7

8 9 **Data Synthesis**

10
11 Data from the eligible articles were abstracted into evidence tables. Consensus or variations related
12 to relevant variables were reported and presented as descriptive summaries in the result section. A
13 meta-analysis was performed for assessing pooled deficiency of vitamin D in pediatric patients of
14 COVID-19. We used a random-effect/fixed-effect model (based on heterogeneity) to calculate
15 pooled weighted proportions with the 95% confidence interval. Heterogeneity or between-study
16 variance was assessed using Cochran's Q test ($p < 0.10$), and quantified using the I^2 statistics. $I^2 >$
17 50% and a $p < 0.05$ were considered substantial heterogeneity. All statistics of meta-analysis were
18 performed using MedCalc (Windows) version 15.0 (MedCalc Software, Ostend, Belgium).
19 Publication bias was assessed using Begg's test and Egger's test.
20
21
22
23
24
25
26
27

28 29 **Results**

30
31 The current systematic review synthesized evidence from eight articles having different study
32 designs.¹¹⁻¹⁸ This included two review articles published from Romania and Italy, rest of the
33 articles were either retrospective cohort or case series studies. Out of eight, six articles provided
34 details of vitamin D levels in pediatric COVID patients and were included for quantitative
35 assessment through meta-analysis. Two studies assessing the role of vitamin D levels in disease
36 severity were also included in meta-analysis to derive cumulative odds of developing poorer
37 outcomes in case of vitamin D deficiency.^{11,18}
38
39
40
41
42
43

44 Characteristic details of the included studies are presented in table 1. According to geographic
45 location, studies were reported from various parts of Europe and America. The sample size of the
46 population varied widely with the lowest reporting as 14 patients to the highest as 9,87,849
47 patients.^{14,13} Half of the studies included both pediatric and adult populations and four studies
48 presented data exclusively from patients ageing less than 18 years. Though our search included
49 details till June 2021, evidence synthesis included studies published from March 2020 to January
50 2021.
51
52
53
54
55
56
57
58
59
60

1
2
3 Key findings from each study were summarized and tabulated (table 2). Levels of vitamin D in
4 paediatric COVID patients were provided by six studies and a pooled prevalence of vitamin D
5 deficiency was analysed. Meta-analysis of the quantitative data (six studies) showed a pooled
6 prevalence of vitamin D deficiency as 45.91% (95% CI: 25.148 to 67.450; random effect model)
7 (figure 2). This meta-analysis showed higher heterogeneity ($p < 0.0001$) but a lack of significant
8 publication bias. This heterogeneity was imparted due to a wider range of vitamin D deficiency
9 reporting by various studies (range: 5.7 – 72.5%). Some of the studies have also pointed to an
10 inverse relationship between vitamin D levels and COVID-19 infection risk, indicating higher odds
11 of infection in individuals with vitamin D deficiency as compared to their counterparts having
12 normal vitamin D levels.^{13,17} Apart from increased risk of infection, meta-analysis also showed
13 that in infected paediatric patients, low levels of vitamin D increases the risk of severe course of
14 disease (figure 3; odds ratio: 5.5; 95% CI: 1.560- 19.515; $p = 0.008$). Though only two studies
15 included in the quantitative analysis, there was a complete absence of heterogeneity ($I^2 = 0\%$). We
16 further wanted to study the association in deficient and insufficient groups, however due to non-
17 availability of categorized data this cannot be explored.

30 Discussion

31
32
33 This global systematic review provides the most recent evidence synthesis addressing the
34 relationship between vitamin D and COVID-19 in pediatric patients. It provided not only the
35 quantitative estimate of vitamin D deficiency prevalence in children and adolescents infected with
36 COVID-19, but also provided cumulative odds of developing severe disease in patients with low
37 levels.

38
39
40
41
42 Vitamin D has been known for its protective role for ages, as it stimulates the production of
43 antimicrobials in the epithelium of the respiratory tract and provides protection against respiratory
44 tract infections.¹⁹ Vitamin D regulates immune responses by reducing plasma cells,
45 proinflammatory cytokine production and immunoglobulin release, thus increasing anti-
46 inflammatory cytokine production.²⁰⁻²³ Vitamin D also promotes the gene responsible for the
47 expression of angiotensin-converting enzyme 2, which has an anti-inflammatory action.²⁴

48
49
50
51
52
53
54 Our study noted that a considerable number of pediatric patients suffering from COVID-19 had
55 low vitamin D levels. A study by Molla et al assessed the importance of nutrition in the regulation
56 of the immune system, and it was found that most of the pediatric COVID-19 patients were

1
2
3 deficient in Vitamin D.¹⁵ The pathology of COVID-19 provokes release of inflammatory mediators
4 due to the complex interaction between the virus and the immune system. A greater emphasis is
5 placed on vitamin D, as it has become increasingly clear that hyper inflammation is an important
6 component of COVID-19.²⁵ Though evidence related to the effect of vitamin D on pediatric
7 COVID is limited, in adult patients' numerous studies have documented its role. Raharusun et al,
8 in their study conducted on adults, found deficient or insufficient levels of vitamin D in 47.3% of
9 780 patients diagnosed with COVID-19.²⁶ It was also found that vitamin D has a clear impact on
10 disease severity and hence COVID associated mortality. As a result, there is a policy implication
11 adopted by various regulatory bodies, where vitamin D supplementation has been recommended
12 in deficient adults.
13
14
15
16
17
18
19
20

21 Studies included in the current review also assessed the effect of vitamin D deficiency on
22 worsening of clinical outcomes such as increased days of stay in the intensive care unit and
23 mortality. Two studies revealed an association between vitamin D deficiency and clinical severity,
24 in addition to inflammation markers in COVID-19 pediatric patients. This association was
25 observed when there was a deficiency rather than an insufficiency in the study by Bayramoglu et
26 al.¹¹ A six times higher chance of having a moderate to severe clinical course was demonstrated in
27 Vitamin D deficient patients, using multivariate logistic regression analysis. A study by Yilmaz et
28 al recorded that 72.5% (n=29) of the cases were vitamin D deficient or insufficient out of which
29 two patients needed intensive care.¹⁸ A negative correlation was found between vitamin D levels
30 and fever symptoms. Patients with high fever had deficient or insufficient levels of Vitamin D.
31 Although adequate studies are not available assessing the role of vitamin D and disease severity in
32 pediatric patients, several studies have shown that children with insufficient or deficient levels of
33 vitamin D are more susceptible to respiratory infections.²⁷ Raharusun et al, observed mortality in
34 49.1% and 46.7% of the patients with insufficient and deficient vitamin D levels respectively in
35 contrast to only 4.1% mortality in patients with normal vitamin D levels.²⁶ Also previous studies
36 found an inverse correlation between vitamin D level and pneumonia severity, increased risk of
37 sepsis, CRP level and ARDS risk.²⁸⁻³¹
38
39
40
41
42
43
44
45
46
47
48
49
50

51 Vitamin D is known as 'sunshine vitamin', as sunlight helps in the conversion of 7-
52 dehydrocholesterol in the skin to cholecalciferol. It gets converted into 1,25(OH)D or calcitriol
53 through hydroxylation in the liver and kidney. Vitamin D receptors (VDR) are activated by 1,25
54 dihydroxy vitamin D.^{32,33} Since sunlight alone cannot provide sufficient amounts of vitamin D to
55
56
57
58
59
60

1
2
3 everyone, supplements are recommended. Vitamin D supplementation helps to decrease
4 proinflammatory cytokines in the lung by the modulation of T lymphocyte activity two concerns
5 remain regarding Vitamin D supplementation and disease outcome, firstly the appropriate levels
6 needed to produce an immunomodulatory effect is unknown. Secondly, the effects of
7 supplementation may not be observed by administration of a single dose of vitamin D. Therefore,
8 a study conducted in adults, they chose an arbitrary value of 25(OH)D levels >50ng/ml to produce
9 an immunomodulatory effect. It was observed, that a single dose administration of 5,40,000IU of
10 vitamin D3 increased the levels to >20ng/ml and more than 38.2±16.5ng/ml at one week.³⁴ A
11 recent meta-analysis, including 25 randomized controlled trial showed that the risk of developing
12 acute respiratory tract infection could be reduced by the supplementation of Vitamin D weekly.³⁵
13 Though the present study provides evidence that COVID-19 severity has an association with
14 vitamin D deficiency, more robust evidence through well-conducted randomized controlled trials
15 are required to substantiate this further. Till the time findings emerge from more prospective
16 studies, supplementation can be recommended in pediatric COVID patients especially in case of
17 deficiency based on results of the current review.
18
19
20
21
22
23
24
25
26
27
28
29

30 Though the review provides some of the most useful information, it suffers from few inherent
31 limitations. Due to limited and heterogeneously published literature, the effect of vitamin D
32 supplementation on COVID-19 infectivity and severity could not be explored in pediatrics.
33 Similarly, the influence of age, gender and other confounders on vitamin D deficiency and COVID
34 outcome were also not studied due to the unavailability of primary studies.
35
36
37
38
39

40 **Conclusion:**

41
42 The systematic review showed that almost half of the pediatric COVID patients suffer from
43 vitamin D deficiency. It is also clear that the deficiency is associated with greater risk of infection
44 and poor outcome in pediatrics. Vitamin D supplementation needs to be explored as a preventive
45 prophylactic, for the prevention and management of COVID-19 in children and adolescents.
46
47
48
49

50 **References**

- 51
52
53 1 Lim S, Bae JH, Kwon H-S, Nauck MA. COVID-19 and diabetes mellitus: from
54 pathophysiology to clinical management. *Nature Reviews Endocrinology*. 2020;17(1):11-30.
55 doi:10.1038/s41574-020-00435-4
56
57
58
59
60

- 1
2
3 **2** Sanyaolu A, Okorie C, Marinkovic A, Patidar R, Younis K, Desai P, et al. Comorbidity and its
4 Impact on Patients with COVID-19. *SN Comprehensive Clinical Medicine*. 2020;2(8):1069-
5 1076. doi:10.1007/s42399-020-00363-4
6
7
- 8 **3** Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases,
9 cancers, and cardiovascular disease. *The American Journal of Clinical Nutrition*. 2004;80(6).
10 doi:10.1093/ajcn/80.6.1678s
11
- 12 **4** Jolliffe D, Camargo C, Sluyter J, Martineau A. S100 Vitamin D supplementation to prevent
13 acute respiratory infections: systematic review and meta-analysis of aggregate data from
14 randomised controlled trials. *Baby and bathwater: not all lung infections are COVID-19*.
15 2021:276-292. doi:10.1136/thorax-2020-btsabstracts.105
16
17
- 18 **5** Weir EK, Thenappan T, Bhargava M, Chen Y. Does vitamin D deficiency increase the severity
19 of COVID-19? *Clinical Medicine*. 2020;20(4):107-108. doi:10.7861/clinmed.2020-0301
20
21
- 22 **6** Biesalski HK. Vitamin D deficiency and co-morbidities in COVID-19 patients – A fatal
23 relationship? *NFS Journal*. 2020; 20:10-21. doi: 10.1016/j.nfs.2020.06.001.
24
25
- 26 **7** Asyary A, Veruswati M. Sunlight exposure increased Covid-19 recovery rates: A study in the
27 central pandemic area of Indonesia. *Science of The Total Environment*. 2020; 729:139016. doi:
28 10.1016/j.scitotenv.2020.139016
29
30
- 31 **8** Ratnesar-Shumate S, Williams G, Green B, Krause M, Holland B, Wood S, et al. Simulated
32 Sunlight Rapidly Inactivates SARS-CoV-2 on Surfaces. *The Journal of Infectious Diseases*.
33 2020;222(2):214-222. doi:10.1093/infdis/jiaa274
34
35
- 36 **9** Shah K, Saxena D, Mavalankar D. Vitamin D supplementation, COVID-19 and disease
37 severity: a meta-analysis. *QJM: An International Journal of Medicine*. 2021;114(3):175-181.
38 doi:10.1093/qjmed/hcab009
39
40
- 41 **10** Arksey H, O'Malley L. studies: towards a methodological framework. *International Journal*
42 *of Social Research Methodology*. 2005;8(1):19-32. doi:10.1080/1364557032000119616
43
44
- 45 **11** Bayramoğlu E, Akkoç G, Ağbaş A, Akgün Özlem, Yurdakul K, Selçuk Duru HN, et al. The
46 association between vitamin D levels and the clinical severity and inflammation markers in
47 pediatric COVID-19 patients: single-center experience from a pandemic hospital. *European*
48 *Journal of Pediatrics*. 2021. doi:10.1007/s00431-021-04030-1
49
50
- 51 **12** Feketea G, Vlacha V, Bocsan IC, Vassilopoulou E, Stanciu LA, Zdrenghea M. Vitamin D in
52 Corona Virus Disease 2019 (COVID-19) Related Multisystem Inflammatory Syndrome in
53 Children (MIS-C). *Frontiers in Immunology*. 2021;12. doi:10.3389/fimmu.2021.648546
54
55
56
57
58
59

- 1
2
3 **13** Katz J, Yue S, Xue W. Increased risk for COVID-19 in patients with vitamin D
4 deficiency. *Nutrition*. 2021; 84:111106. doi: 10.1016/j.nut.2020.111106
5
6
7 **14** Nadiger M, Hassor S, Totapally B. 138: Vitamin D Levels in Children With COVID-19
8 Admitted to the PICU. *Critical Care Medicine*. 2020;49(1):54-54. doi:
9 10.1097/01.ccm.0000726440.30551.47
10
11 **15** Karakaya Molla G, Ünal Uzun Özlem, Koç N, Özen Yeşil B, Bayhan G İclal. Evaluation of
12 nutritional status in pediatric patients diagnosed with Covid-19 infection. *Clinical Nutrition*
13 *ESPEN*. 2021. doi: 10.1016/j.clnesp.2021.04.022
14
15
16 **16** Panfili FM, Roversi M, D'Argenio P, Rossi P, Cappa M, Fintini D. Possible role of vitamin D
17 in Covid-19 infection in pediatric population. *Journal of Endocrinological Investigation*.
18 2020;44(1):27-35. doi:10.1007/s40618-020-01327-0
19
20
21 **17** SÖBÜ E, KARAASLAN A, ÇETİN C, AKIN Y. Vitamin D Levels of COVID-19 Positive
22 Symptomatic Pediatric Cases. *Güncel Pediatri*. 2021;19(1):9-14. doi:10.4274/jcp.2021.0002
23
24
25 **18** YILMAZ kamil, en V. Is Vitamin D Deficiency a Risk Factor for Covid 19 in Children?
26 *Pediatr Pulmonol*. 2020;55(12):3595-3601. doi:10.22541/au.159318959.95163592
27
28
29 **19** Zhang W., Zhao Y., Zhang F., Wang Q., Li T., Liu Z. The use of anti-inflammatory drugs in
30 the treatment of people with severe coronavirus disease 2019 (COVID-19): The Perspectives
31 of clinical immunologists from China. *Clin Immunol*. 2020 May; 214:108393. doi:
32 10.1016/j.clim.2020.108393
33
34
35 **20** Chen S, Sims GP, Chen XX, Gu YY, Chen S, Lipsky PE. Modulatory Effects of 1,25-
36 Dihydroxyvitamin D₃ on Human B Cell Differentiation. *The Journal of Immunology*.
37 2007;179(3):1634-1647. doi:10.4049/jimmunol.179.3.1634;
38
39
40 **21** Lemire JM. Immunomodulatory role of 1,25-dihydroxyvitamin D₃. *Journal of Cellular*
41 *Biochemistry*. 1992;49(1):26-31. doi:10.1002/jcb.240490106
42
43
44 **22** Zhang Y, Leung DYM, Richers BN, Liu Y, Remigio LK, Riches DW, et al. Vitamin D Inhibits
45 Monocyte/Macrophage Proinflammatory Cytokine Production by Targeting MAPK
46 Phosphatase-1. *The Journal of Immunology*. 2012;188(5):2127-2135.
47 doi:10.4049/jimmunol.1102412
48
49
50 **23** Liu P, Stenger S, Li H, Wenzel L, Tan B, Krutzik S, et al. Vitamin D₃-Triggered Antimicrobial
51 Response—Another Pleiotropic Effect beyond Mineral and Bone Metabolism. *Journal of the*
52 *American Society of Nephrology*. 2006;17(11):2949-2953. doi:10.1681/asn.2006091030
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 24 Cui C, Xu P, Li G, Qiao Y, Han W, Geng C, et al. Vitamin D receptor activation regulates microglia polarization and oxidative stress in spontaneously hypertensive rats and angiotensin II-exposed microglial cells: Role of renin-angiotensin system. *Redox Biology*. 2019; 26:101295. doi: 10.1016/j.redox.2019.101295
- 25 Im J.H., Je Y.S., Baek J., Chung M.H., Kwon H.Y., Lee J.S. Nutritional status of patients with COVID-19. *Int J Infect Dis*. 2020 Nov; 100:390–393. doi: 10.1016/j.ijid.2020.08.018.
- 26 Raharusun P, Priambada S, Budiarti C, Agung E, Budi C. Patterns of COVID-19 Mortality and Vitamin D: An Indonesian Study. *SSRN Electronic Journal*. 2020. doi:10.2139/ssrn.3585561
- 27 Baqui A, Sabir A, Begum N, Arifeen S, Mitra S, Black R. Causes of childhood deaths in Bangladesh: an update. *Acta Paediatrica*. 2007;90(6):682-690. doi:10.1111/j.1651-2227.2001.tb02434.x
- 28 Lu D, Zhang J, Ma C, Yue Y, Zou Z, Yu C, et al. Link between community-acquired pneumonia and vitamin D levels in older patients. *Zeitschrift für Gerontologie und Geriatrie*. 2017;51(4):435-439. doi:10.1007/s00391-017-1237-z
- 29 Zhou Y-F, Luo B-A, Qin L-L. The association between vitamin D deficiency and community-acquired pneumonia. *Medicine*. 2019;98(38):17252. doi:10.1097/md.00000000000017252
- 30 Manion M, Hullsiek KH, Wilson EMP, Rhame F, Kojic E, Gibson D, et al. Vitamin D deficiency is associated with IL-6 levels and monocyte activation in HIV-infected persons. *PLOS ONE*. 2017;12(5). doi: 10.1371/journal.pone.0175517
- 31 Dalvi SM, Ramraje NN, Patil VW, Hegde R, Yeram N. Study of IL-6 and vitamin D3 in patients of pulmonary tuberculosis. *Indian Journal of Tuberculosis*. 2019;66(3):337-345. doi: 10.1016/j.ijtb.2018.05.018
- 32 Bikle D. Vitamin D Metabolism, Mechanism of Action, and Clinical Applications. *Chemistry & Biology*. 2014;21(3):319-329. doi: 10.1016/j.chembiol.2013.12.016
- 33 Morris HA, Anderson PH. Autocrine and paracrine actions of vitamin d. *Clin Biochem Rev*. 2010;31(4):129-38.
- 34 Tellioglu A, Basaran S, Guzel R, Seydaoglu G. Efficacy and safety of high dose intramuscular or oral cholecalciferol in vitamin D deficient/insufficient elderly. *Maturitas*. 2012;72(4):332-338. doi: 10.1016/j.maturitas.2012.04.011
- 35 Martineau A, Jolliffe D, Hooper R, Greenberg L, Aloia J, Bergman P, et al. S102 Vitamin d supplementation to prevent acute respiratory infections: systematic review and meta-analysis

1
2
3 of individual participant data. *Thorax*. 2016; 71:356-6583. doi:10.1136/thoraxjnl-2016-
4 209333.108
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Characteristic details of the studies included in the review

Sr. No	Author	Title	Country	Study type	Population	Study period	Total cases
1	Bayramoğlu et al 2021 ¹¹	The association between vitamin D levels and the clinical severity and inflammation markers in pediatric COVID-19 patients: single-center experience from a pandemic hospital	Germany	Retrospective cohort study	Pediatric (1 to 18 years)	March and May 2020	356
2	Feketea et al 2021 ¹²	Vitamin D in Corona Virus Disease 2019 (COVID-19) Related Multisystem Inflammatory Syndrome in Children (MIS-C)	Romania	Mini review	Children and adults	up to 30th December 2020	Not provided
3	Katz et al 2020 ¹³	Increased risk for COVID-19 in patients with vitamin D deficiency	USA	Cross sectional study	Children and adults	Up to June, 30, 2020	987849
4	Nadiger et al 2021 ¹⁴	Vitamin D Levels in Children With COVID-19 Admitted to the PICU	USA	Retrospective case series	Children	Jan 2021	14
5	Molla et al 2021 ¹⁵	Evaluation of nutritional status in pediatric patients diagnosed with Covid-19 infection	Turkey	Evaluation study	Pediatric (1 to 18 years)	May and 15 June 2020	49
6	Panfili et al 2020 ¹⁶	Possible role of vitamin D in Covid-19 infection in pediatric population	Italy	Short review	Children and adults	Not provided	Not provided
7	Söbü et al 2021 ¹⁷	Vitamin D Levels of COVID-19 Positive Symptomatic Pediatric Cases	Turkey	Retrospective study	Pediatric	March to April 2020	112
8	Yılmaz & Sen 2020 ¹⁸	Is vitamin deficiency a risk factor for COVID-19 in children?	Turkey	Retrospective study	Pediatric (1 to 18 years)	March 2020 and May 2020	85

Table 2: Key findings of the studies included in review

Sr. No.	Author	Key findings
1	Bayramoğlu et al 2021 ¹¹	<ul style="list-style-type: none"> • There is an association between vitamin D deficiency and clinical severity in pediatric COVID-19 cases. • Vitamin D level was correlated positively with the lymphocyte count, and negatively with age, CRP and fibrinogen levels. • Vitamin D deficiency is an independent predictor of severe clinical outcome.
2	Feketea et al 2021 ¹²	<ul style="list-style-type: none"> • Role of vitamin D was evaluated as biomarker and nutritional supplement for MIS-C • Lower concentration of serum vitamin D may be significantly associated with neonatal pneumonia and poorer outcome • Supplementation of vitamin D might improve the outcome, though more studies are needed to substantiate this
3	Katz et al 2020 ¹³	<ul style="list-style-type: none"> • Patients with vitamin D deficiency were 4.6 times more likely to be positive for COVID-19 than patients with no deficiency ($P < 0.001$). • The association of low vitamin D level remains significant even after adjusting effect of race, periodontal disease, diabetes and obesity.
4	Nadiger et al 2021 ¹⁴	Majority of the pediatric patients who tested positive for COVID19 and in need of ICU treatment were obese, vitamin D deficient adolescents.
5	Molla et al 2021 ¹⁵	<ul style="list-style-type: none"> • Vitamin D deficiency was the most common vitamin deficiency among COVID-19 pediatric patients. • It is possible that vitamin D deficiency increases susceptibility to the infection.
6	Panfili et al 2020 ¹⁶	<ul style="list-style-type: none"> • Vitamin D might have a role in prevention and control of COVID-19 in both adults and pediatric population • Role of vitamin D in preventing or even treating the disease urgently needs to be further addressed especially in patients with deficiencies
7	Söbü et al 2021 ¹⁷	<ul style="list-style-type: none"> • Vitamin D levels were assessed between healthy and children with COVID-19 infection. • Median vitamin D was significantly (8.9 vs 18.5 ng/mL; $p < 0.05$) lower in hospitalized COVID patients as compared to healthy control. • Age and gender did not differ among both the groups. • 93.3% of COVID positive patients were given Vitamin D supplementation and showed good outcome.
8	Yilmaz & Sen 2020 ¹⁸	Children with COVID-19 had significantly lower vitamin D levels as compared to their control counterparts [13.14 $\mu\text{g/L}$ (4.19–69.28) vs 34.81 (3.8–77.42) $\mu\text{g/L}$; ($p < 0.001$)].


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1: PRISMA chart of study selection

Figure 2: Forest plot of vitamin D deficiency (%) prevalence in paediatric COVID patients

Figure 3: Forest plot of association between low level of vitamin D and disease severity in paediatric COVID patients


Downloaded from <https://academic.oup.com/qjmed/advance-article/doi/10.1093/qjmed/hcab202/6325519> by guest on 23 July 2021



PRISMA chart

42x43mm (300 x 300 DPI)


Study	Sample size	Proportion (%)	95% CI	Weight (%)	
				Fixed	Random
Bayramoglu et al 2021	103	41.748	32.105 to 51.877	37.55	17.83
Katz et al 2020	35	5.714	0.700 to 19.157	13.00	16.74
Nadiger et al 2020	14	71.429	41.896 to 91.611	5.42	14.81
Molla et al 2021	49	26.531	14.947 to 41.082	18.05	17.19
Sobu et al 2021	30	70.000	50.604 to 85.265	11.19	16.49
Yilmaz & Sen 2020	40	72.500	56.112 to 85.399	14.80	16.93
Total (fixed effects)	271	42.580	36.683 to 48.636	100.00	100.00
Total (random effects)	271	45.912	25.148 to 67.450	100.00	100.00



Meta-analysis of vitamin D deficiency (proportion) in pediatric COVID patients

130x37mm (300 x 300 DPI)

Study	Intervention	Controls	Odds ratio	95% CI	Z	P	Weight (%)	
							Fixed	Random
Bayramoglu et al 2021	24/43	3/19	6.737	1.708 to 26.570			83.74	83.74
Yimaz and Sen et al 2020	2/29	0/11	2.091	0.0929 to 47.043			16.26	16.26
Total (fixed effects)	26/72	3/30	5.517	1.560 to 19.515	2.649	0.008	100.00	100.00
Total (random effects)	26/72	3/30	5.569	1.587 to 19.549	2.681	0.007	100.00	100.00



Forest plot of association between low level of vitamin D and disease severity in paediatric COVID patients
140x38mm (300 x 300 DPI)