

This Provisional PDF corresponds to the article as it appeared upon acceptance. Fully formatted PDF and full text (HTML) versions will be made available soon.

Recent incidence of type 1 diabetes mellitus in children 0--14 years in Newfoundland and Labrador, Canada climbs to over 45/100,000: a retrospective time trend study

BMC Research Notes 2012, **5**:628 doi:10.1186/1756-0500-5-628

Leigh A Newhook (lnewhook@mun.ca)
Sharon Penney (sharonj.penney@easternhealth.ca)
Jackie Fiander (fianderjj@hotmail.com)
Jeff Dowden (jeff.dowden@nlchi.nl.ca)

ISSN 1756-0500

Article type Short Report

Submission date 22 June 2012

Acceptance date 5 November 2012

Publication date 12 November 2012

Article URL <http://www.biomedcentral.com/1756-0500/5/628>

This peer-reviewed article can be downloaded, printed and distributed freely for any purposes (see copyright notice below).

Articles in *BMC Research Notes* are listed in PubMed and archived at PubMed Central.

For information about publishing your research in *BMC Research Notes* or any BioMed Central journal, go to

<http://www.biomedcentral.com/info/instructions/>

© 2012 Newhook *et al.*

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Recent incidence of type 1 diabetes mellitus in children 0–14 years in Newfoundland and Labrador, Canada climbs to over 45/100,000: a retrospective time trend study

Leigh A Newhook^{1,2,*}
Email: lnewhook@mun.ca

Sharon Penney²
Email: sharonj.penney@easternhealth.ca

Jackie Fiander¹
Email: fianderjj@hotmail.com

Jeff Dowden³
Email: jeff.dowden@nlchi.nl.ca

¹ Janeway Pediatric Research Unit, Memorial University, 300 Prince Philip Drive, St. John's, NL A1B 3V6, Canada

² Janeway Children's Health and Rehabilitation Centre, Eastern Health, St. John's, NL, Canada

³ Newfoundland and Labrador Centre for Health Information, St. John's, NL, Canada

* Corresponding author. Janeway Pediatric Research Unit, Memorial University, 300 Prince Philip Drive, St. John's, NL A1B 3V6, Canada

Abstract

Background

To study and update the provincial incidence of type 1 diabetes mellitus (T1DM) in Newfoundland and Labrador (NL), a province of Canada with a very high incidence previously reported in 2006, and one of the highest incidences reported worldwide. This is a retrospective time trend study of the incidence of T1DM, in children aged 0–14 years from 1987–2010 inclusive.

Findings

Over the study period 931 children aged 0–14 years were diagnosed with T1DM. The incidence of T1DM in this population over the period 1987 – 2010 inclusive was 37.7 per 100,000 per year (95% CI 35.3, 40.2)

The incidence from 2007–2010 was 49.9 per 100,000 per year (95% CI 42.2, 57.6). The incidence over this 24 year period increased by a factor of 1.03 per 100,000 per year.

Conclusion

NL has one of the highest incidences of T1DM reported worldwide. Potential reasons for the very high incidence could be related to the unique genetic background of the population, northern latitude and vitamin D insufficiency, low breastfeeding rates, and high rates of cesarean section.

Keywords

Type 1 diabetes mellitus, Incidence, Newfoundland and Labrador, Canada, Epidemiology

Findings

Background

In 2006, we reported a “very high” incidence of 35.1 per 100,000 per year of T1DM in the 0–14 year age group over the period 1987–2005 in Newfoundland and Labrador (NL) [1]. Other findings showed a significant difference between the incidence for boys versus girls in the 0–4-year age group (31.61 and 19.05 respectively per 100,000 per year, $p=0.0013$) and the estimated rate of incidence increase was 0.783 per 100,000 children ages 0–14 years per year [1]. The rates of T1DM in NL appear to be continually rising and this update was done to investigate this hypothesis.

Methods

This retrospective time trend study was performed across the Canadian province of NL. All children with T1DM who live in NL and were diagnosed with T1DM from 1987 to 2010 are included in the study. Patient sex, month of onset of diabetes, month of birth, and age of onset were collected and recorded by diabetes nurse educators in each facility that cares for children with T1DM and reports of new cases are faxed to the research nurse monthly at the Janeway Pediatric Research Unit at the tertiary care Janeway Children’s Health and Rehabilitation Centre, Eastern Health, in St. John’s. Classification, case definition and ascertainment are described elsewhere [1,2].

Ethical considerations

The Memorial University of Newfoundland and Labrador Human Investigations Committee and each of the hospital board’s ethics committees approved this study.

Statistical analysis

The data described is incidence per 100,000 of the age-specific population obtained from Canadian census data [3]. Age-standardized rates were computed across specified time periods using 1991 Canadian Census population as standard. 95% confidence intervals were

derived using Poisson distribution. Poisson log-linear regression was used to examine trends in annual incidence rates.

Results

A total of 931 cases were identified (Table 1); there were 483 males and 446 females with 1.1:1 ratio (M:F). Two cases did not specify gender. Incidence was significantly higher for males than females in the 0–4 year age group (m=32.7 vs. f=21.7 per 100,000 per year; p=0.006). Yearly incidence rates for ages 0–4 increased by a factor of 1.03 per 100,000 (95% CI: 1.01-1.04); whereas rates for ages 5–9 and 10–14 increased by factors of 1.04 per 100,000 (1.02-1.05), and 1.02 per 100,000 (1.01-1.04), respectively.

Table 1 Incidence, gender, and time trends of type 1 diabetes mellitus (0–14 yr) per 100,000 individuals in Newfoundland and Labrador, Canada, 1987-2010

Years	Sex	0-14 yr (n)	95% CI	0-4 yr (n)	95% CI	5-9 yr (n)	95% CI	10-14 yr (n)	95% CI
1987-2010	M+F	37.7 (931)	35.3-40.2	27.3 (196)	23.6-31.4	42.8 (350)	38.5-47.6	41.1 (385)	37.1-45.4
1987-2010	M	38.3 (485)	35.0-41.9	32.7* (120)	27.1-39.1	40.0 (167)	34.2-46.5	41.2 (198)	35.7-47.3
1987-2010	F	37.0 (446)	33.6-40.6	21.7* (76)	17.1-27.1	45.8 (183)	39.4-53.0	41.0 (187)	35.3-47.3
1987-1991	M+F	29.9 [†] (207)	24.4-35.5	21.3 (42)	15.3-28.7	33.2 (75)	26.1-41.7	35.3 (90)	28.4-43.4
1992-1996	M+F	31.9 [†] (190)	26.1-37.8	27.3 (47)	20.1-36.3	37.3 (73)	29.2-46.9	31.2 (70)	24.3-39.4
1997-2001	M+F	40.9 [†] (203)	34.3-43.4	28.0 (38)	19.8-38.4	46.7 (75)	36.7-58.5	48.1 (90)	38.7-59.1
2002-2006	M+F	42.3 [†] (178)	35.5-49.2	30.2 (36)	21.2-41.8	50.5 (68)	39.2-64.0	46.4 (74)	36.4-58.2
2007-2010	M+F	49.9 [†] (153)	42.2-57.6	35.8 (33)	24.6-50.2	59.1 (59)	45.0-76.3	54.7 (61)	41.9-70.4

CI, confidence interval; n, specified size of age group; F, Female; M, Male; [†] rates age-standardized to 1991 Canadian Census; * p=0.006

Trends were not significantly different across age groups. Age-standardized rates for census divisions ranged from 18.8 per 100,000 per year on the South Coast to 40.6 per 100,000 per year on the Northern Peninsula (Table 2).

Table 2 Incidence of type 1 diabetes mellitus (0–14 yr) by geographical census division[±], Newfoundland and Labrador, Canada, 1987-2010

Census Division	Incidence rate	95% CI
(1) Avalon Peninsula	40.2	34.9-45.4
(2) Burin Peninsula	33.3	24.6-42.0
(3) South Coast	18.8	10.7-26.9
(4) St. George's	33.7	24.7-42.6
(5) Humber District	33.6	25.7-41.5
(6) Central	45.8	37.1-54.5
(7) Bonavista/Trinity	33.3	25.3-41.4
(8) Notre Dame Bay	37.8	29.9-45.7
(9) Northern Peninsula	40.6	30.9-50.3
(10) Labrador	39.9	31.1-48.8

[±] Rates age-standardized to 1991 Canadian Census

Discussion

This study represents an updated analysis on the population of children 0–14 years with T1DM from the province of NL, Canada, confirming a very high and increasing incidence over the 24-year study period. The incidence from 1987 to 2010 inclusive is 37.7 per 100,000 per year, one of the highest reported worldwide. The incidence from 2007–2010 was 49.9 per 100,000 per year. The incidence increased significantly in all age groups and there are significantly more males diagnosed with T1DM than females in the 0–4 year age group. A male excess is more often found in populations with a high incidence of T1DM [4,5].

The incidence of T1DM is increasing worldwide, especially for the youngest ages and those with moderate genetic susceptibility. The rapidly rising incidence cannot be explained by genetic susceptibility alone and is hypothesized to be due to environmental, lifestyle, and or epigenetic factors that are triggering or accelerating the onset and progression of the disease [6]. The increasing incidence that we are reporting for NL is similar to reports from other large epidemiologic studies including DiaMOND [7], EURODIAB [8], and SEARCH [9]. Other regions of Canada have also reported high rates of T1DM but data on trends over time are not available [10,11]. Although we do not have an explanation for the unusually high incidence of T1DM in NL, there are several factors that have been identified in other regions that may be important for this population. Genetic factors are likely to be important and research is underway to explore this further. Newfoundland is an island and Labrador, although part of mainland Canada, is relatively isolated. The population of NL has a well-documented immigration history with 96% of residents having ancestors either from southeast Ireland or southwest England [12], as well as low intraprovincial migration rates [12,13] and founder effects [14,15].

The rising incidence over a relatively short period of time however suggests environmental factors are contributing to the increase in a genetically at risk population. Potential environmental risk factors that have been identified in other populations which may be associated with T1DM in NL include early infant nutrition, perinatal factors, northern latitude, and vitamin D deficiency. NL has the lowest rates of initiation and duration of breastfeeding in Canada [16]; exclusive breastfeeding has been shown have a weak protective association with T1DM risk [17-19]. Vitamin D deficiency has been associated with increased risk of T1DM [20,21], vitamin D supplementation may be protective against T1DM [22] and UVB irradiation and northern latitude are also associated factors [23]. NL has a northern latitude (49°N) and erythemal UVB radiation may be geospatially associated with the incidence of T1DM NL [24]. Vitamin D insufficiency has been documented in the population [25]. NL has a high rate of birth by C-section as compared to other regions in Canada. The provincial rate of births by C-section was 30.9% in 2005–2006 versus the Canadian rate of 26.3% [26]. The rates of C-section have increased in NL to 33% in 2010 [27]. Recent results from a population-based case–control study showed that birth by cesarean section is associated with T1DM in NL (HR 1.41, $p=0.015$) [28]. A recent meta-analysis of 20 studies found that the combined effect of C-section delivery on T1DM risk was 1.23 (95% CI 1.15-1.32) [29].

Conclusions

The results from this research continue to show a very high and increasing incidence. We continue to research the genetics, epidemiology and environmental risk factors of the disease in this population with hopes of contributing to a better understanding of the etiology and

pathogenesis of T1DM. Research into the relationship and underlying reasons for the association between birth by c-section and T1DM risk are continuing, as well as research on vitamin D insufficiency, early infant nutrition and growth. NL is a unique and important place to study T1DM with its geography, stable population, genetic founder effects and unusually high incidence.

Competing interest

There are no conflicts of interest to disclose.

Authors' contributions

LN is the principle investigator and was responsible for the intellectual conception and design of the study, funding application, manuscript preparation and is the guarantor of the research. SP was the research nurse who was responsible for study approvals, coordination of the study as well as gathering, confirming and entering of data. JF was responsible for overall coordination of the study and contributed intellectually to the methods, database development, logistics, approvals, and funding application submission. JD was responsible for the statistical analysis. All authors contributed to and approved the final manuscript submission.

Acknowledgements

Thank you to the Janeway Hospital Research Foundation for funding. Thank you to all the diabetes nurse educators throughout the province of Newfoundland and Labrador for your participation and continued efforts.

References

1. Newhook LA, Grant M, Sloka S, Hoque M, Paterson AD, Hagerty D, Curtis J: **Very high and increasing incidence of type 1 diabetes mellitus in Newfoundland and Labrador, Canada.** *Pediatr Diabetes* 2008, **9**:62–68.
2. Newhook LA, Curtis J, Hagerty D, Grant M, Paterson A, Crummel C, Bridger T, Parfrey P: **High incidence of childhood type 1 diabetes in the Avalon Peninsula, Newfoundland, Canada.** *Diabetes Care* 2004, **27**:885–888.
3. Statistics Canada: *Statistics Canada National Census*; 2011.
<http://www12.statcan.gc.ca/census-recensement/2011/as-sa/98-310-x2011001-eng.cfm>.
4. Karvonen M, Pitkaniemi M, Pitkaniemi J, Kohtamaki K, Tajima N, Tuomilehto J, for the WHO DIAMOND Project Group: **Sex difference in the incidence of insulin-dependent diabetes mellitus: an analysis of the recent epidemiological data.** *Diabetes Metab Rev* 1997, **13**:275–291.
5. Gale E, Gillespie K: **Diabetes and gender.** *Diabetologia* 2001, **44**:3–15.

6. Vehik K, Dabelea D: **The changing epidemiology of type 1 diabetes: why is it going through the roof?** *Diabetes Metab Res Rev* 2011, **27**:3–13.
7. DiaMOND Project Group: **Incidence and trends of childhood type 1 diabetes worldwide 1990–1999.** *Diabet Med* 2006, **23**:857–866.
8. Patterson C, Dahlquist G, Gyurus E, Green A, Soltesz G: **Incidence trend for childhood type 1 diabetes in Europe during 1989–2003 and predicted new cases 2005–20: a multicentre prospective registration study.** *Lancet* 2009, **373**:2027–2033.
9. Bell RA, Mayer-Davis EJ, Beyer JW, D'Agostino RB Jr, Lawrence JM, Linder B, Liu LL, Marcovina SM, Rodriguez BL, Williams D, Dabelea D: **Diabetes in non-Hispanic white youth: prevalence, incidence, and clinical characteristics: the SEARCH for diabetes in youth study.** *Diab care* 2009, **32**:s102–s111.
10. Blanchard JF, Dean H, Anderson K, Wadja A, Ludwig S, Depew N: **Incidence and prevalence of diabetes in children aged 0–14 years in Manitoba, Canada, 1985–1993.** *Diabetes Care* 1997, **20**:512–517.
11. Toth EL, Lee KC, Couch RM, Martin LF: **High incidence of IDDM over 6 years in Edmonton, Alberta, Canada.** *Diabetes Care* 1993, **16**:796–800.
12. Bear JC, Kennedy JC, Marshall W, Power AA, Kolonel VM, Burke GB: **Persistent genetic isolation in outport Newfoundland.** *Am J Med Genet* 1987, **27**:807–830.
13. Martin LJ, Crawford MH, Koertvelyessy T, Keeping D, Collins M, Huntsman R: **The population structure of ten Newfoundland outports.** *Hum Biol* 2000, **72**:997–1016.
14. Woods MD, Young TL, Parfrey PS, Hefferton D, Green JS, Davidson WS: **Genetic heterogeneity of Bardet Biedl syndrome in a district Canadian population: evidence for a fifth locus.** *Genomics* 1999, **55**:2–9.
15. Andermann EJ, Jacob JC, Andermann F, Carpenters S, Wolfe L, Berkovic S: **The Newfoundland aggregate of neuronal ceroid-lipofuscinosis.** *Am J Med Genet Suppl* 1988, **5**:111–116.
16. Newfoundland and Labrador Provincial Perinatal Program: *Eastern perinatal health surveillance report 2005.* <http://www.nlppp.ca/Eastern%20Avalon%20PHSR%202005.pdf>.
17. Cardwell CR, Stene LC, Ludvigsson J, Rosenbauer J, Cinek O, Svensson J, Perez-bravo F, Memon A, Gimeno SG, Wadsworth EJ, Strotmeyer ES, Goldacre MJ, Radon K, Chuang LM, Parslow R, Chetwynd A, Karavanaki K, Brigis G, Pozzilli P, Urbonaite B, Schober E, Devoti G, Sipetic S, Joner G, Ionescu-Tirgoviste C, de Beaufort CE, Harrild K, Benson V, Savilahti E, Ponsonby AL, Salem M, Rabiei S, Patterson C: **Breast-feeding and childhood-onset type 1 diabetes: a pooled analysis of individual participant data from 43 observational studies.** *Diab care* 2012, **35**(11):2215–2225.

18. Patelarou E, Girvalaki C, Brokalaki H, Patelarou A, Androulaki Z, Vardavas C: **Current evidence on the associations of breastfeeding, infant formula, and cow's milk introduction with type 1 diabetes mellitus: a systematic review.** *Nutr Rev* 2012, **70**:509–519.
19. Sadauskaite-Kuehne V, Ludvigsson J, Padaiga Z, Jasinskiene E, Samuelsson U: **Longer breastfeeding is an independent protective factor against development of type 1 diabetes mellitus in childhood.** *Diabetes Metab Res Rev* 2004, **20**:150–157.
20. Bener A, Alsaied A, Al-Ali M, Al-Kubaisi A, Basha B, Abraham A, Guiter G, Mian M: **High prevalence of vitamin D deficiency in type 1 diabetes mellitus and healthy children.** *Acta Diabetol* 2008, **46**(3):183–189.
21. Greer RM, Rogers MA, Bowling FG, Buntain HM, Harris M, Leong GM, Cotterill AM: **Australian children and adolescents with type 1 diabetes have low vitamin D levels.** *Med J Aust* 2007, **187**:59–60.
22. Zipitis CS, Akobeng AK: **Vitamin D supplementation in early childhood and risk of type 1 diabetes: a systematic review and meta-analysis.** *Arch Dis Child* 2008, **93**(6):512–517.
23. Mohr SB, Garland CF, Gorham ED, Garland FC: **The association between ultraviolet B irradiance, vitamin D status and incidence rates of type 1 diabetes in 51 regions worldwide.** *Diabetologia* 2008, **51**:1391–1398.
24. Sloka S, Grant M, Newhook LA: **Time series analysis of ultraviolet B radiation and type 1 diabetes in Newfoundland.** *Pediatr Diabetes* 2008, **9**:81–86.
25. Newhook LA, Sloka S, Grant M, Randell E, Kovacs C, Twells LK: **Vitamin D insufficiency common in newborns, children, and pregnant women living in Newfoundland and Labrador, Canada.** *Matern Child Nutr* 2009, **5**(2):186–191.
26. Canadian Institute for Health Information: *Giving birth in Canada: regional trends from 2001–2002 to 2005–2006.* 2007.
https://secure.cihi.ca/free_products/childbirth_aib_070725_e.pdf.
27. Research and Evaluation Department: *Newfoundland Centre for Health Information: live birth trends 2002-2006.* http://www.nlchi.nl.ca/images/PDFs/Live_Birth_Trends_2002-2006_Jan08FINAL.pdf.
28. Phillips J, Gill N, Sikdar K, Penney S, Newhook LA: **History of cesarean section associated with childhood onset of T1DM in Newfoundland and Labrador, Canada.** *J Environ Public Health* 2012, **2012**:635097.
29. Cardwell CR, Stene LC, Joner G, Cinek O, Svensson J, Goldacre MJ, Parslow RC, Pozzilli P, Brigis G, Stoyanov D, Urbonaite B, Sipetic S, Schober E, Ionescu-Tirgoviste C, Devoti G, de Beaufort CE, Buschard K, Patterson CC: **Caesarean section is associated with an increased risk of childhood-onset type 1 diabetes mellitus: a meta-analysis of observational studies.** *Diabetologia* 2008, **51**:726–735.