

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/364091064>

Anti-oxidants and Long Covid

Preprint · October 2022

CITATIONS

0

READS

6

1 author:



[Patrick W Chambers](#)

Torrance Memorial Medical Center

17 PUBLICATIONS 415 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



COVID-19 PATHOGENESIS [View project](#)



Anti-oxidants and Long Covid

Patrick Chambers, MD

Department of Pathology, Torrance Memorial Medical Center, Torrance, California

Email: pwc@gte.net

How to cite this paper: Author 1, Author 2 and Author 3 (2022) Paper Title. *Open Access Library Journal*, 9: ****.

https://dx.doi.org/10.4236/****.2022.****

Received: ****, **

Accepted: ****, **

Published: ****, **

Copyright © 2022 by author(s) and Open Access Library Inc.

This work is licensed under the Creative

Commons Attribution International License

(CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Long Covid has many symptoms that overlap with ME(myalgic encephalomyelitis)/CFS(chronic fatigue syndrome), FM(fibromyalgia), EBV(Epstein-Barr virus), CMV(cytomegalovirus), CIRS (chronic inflammatory response syndrome), MCAS(mast cell activation syndrome), POTS(postural orthostatic tachycardia syndrome), and post viral fatigue syndrome. They all portend a “long haul” with an antioxidant shortfall and elevated Ca:Mg. Oxidative stress is the root cause. Linkage between TGF(transforming growth factor)- β , IFN(interferon)- γ , the RAS(renin angiotensin system), and the KKS(kallikrein kinin system) is discussed. Technical explanations for the renin aldosterone paradox in POTS, the betrayal of TGF- β , and the commonality of markers for the Warburg effect are offered. The etiology of the common Long Covid symptoms of post exertional malaise, fatigue, and brain fog as well as anosmia, hair loss, and GI symptoms is technically discussed. Ca:Mg is critical to the glutamate/GABA(gamma amino butyric acid) balance. The role of GABA and butyrates from the “good” intestinal bacteria in the gut brain axis and its correlation with chronic fatigue diseases are explored. The crosstalk between the ENS(enteric nervous system) and the ANS(autonomic nervous system) and the role of the vagus in both are emphasized. HRV(heart rate variability), the fifth vital sign, points to an expanded gut-brain-heart/lung axis. A suggested approach to all of these - Long Covid, chronic fatigue diseases, post viral fatigue syndrome, and general health - is presented.

Subject Areas

Pathology

Keywords

Warburg effect, oxidative stress, magnesuria, inflammasome, butyrate

1. Introduction

Nobel laureate and anti Nazi Otto Warburg first presented his Warburg hypothesis in 1924. He believed that cancer should be interpreted as mitochondrial dysfunction. Mitochondria are the energy factories of the cell and their currency is ATP. Optimal mitochondrial function is at the heart of health. This same cancer related mitochondrial dysfunction also arises in a setting of chronic inflammation and oxidative stress. Antioxidants are protective, but under such conditions their consumption is accelerated. Oxygen is a very toxic element, due to the susceptibility of O_2 to form the superoxide radical $\cdot O_2^-$, which along with its partners, hydrogen peroxide H_2O_2 and the hydroxyl radical $\cdot OH$ constitute reactive oxygen species (ROS). These can also create reactive nitrogen species (RNS), e.g., peroxynitrite $ONOO^-$.

Pathogens and chronic exposure to biotoxins (CIRS) that elicit chronic inflammation can also produce cellular hypoxia and a Warburg environment (mitochondrial dysfunction)(1). Historically the two greatest scourges of mankind, tuberculosis(2,3) and malaria(4,5), can generate their own Warburg effect with elevated TGF- β and lactate (6).

2. Discussion

2.1 Oxidative Stress

The function of antioxidants is to reduce these oxidizing agents (oxidants), which can fatally overwhelm cellular defenses. Oxidative stress develops when oxidants outnumber antioxidants. Cellular hypoxia can develop due to inflammation and ROS production at the gas blood interchange (lungs), during delivery (erythrocytes and endothelial cells), or within the mitochondrion itself. During the latter, intracellular oxidant levels can quickly increase and overcome the onboard antioxidants, which may have been at marginal levels. This creates the hypoxic microenvironment. ROS are primarily produced in mitochondria where the energy of oxygen is transformed into ATP. These oxidizing agents, if not reduced, are very toxic to cells and threaten their death. In order to avoid this, cells shut down their mitochondria, the source of the ROS, to survive. The glycolytic pathway from glucose to pyruvate normally proceeds to the Krebs cycle for oxidative phosphorylation and ATP production within mitochondria. Instead pyruvate proceeds to lactate only. ATP production goes from 38 to 2 ATPs per glucose. Mitochondria are especially dense in muscle cells (skeletal, cardiac, smooth). Fatigue becomes unavoidable. Brain oxygen consumption represents 20% of the total. So, eventually some degree of cognitive compromise is inevitable.

The resulting hypoxia and increased lactic acid trigger release of TGF- β , the primary cytokine of the Warburg effect. This cytokine is elevated in CFS and many other chronic fatigue diseases. TGF- β and IFN- γ counterbalance and suppress each other. IFN- γ possesses C1 (complement component 1 of the CCP) inhibiting properties(7,8).

With the increase in TGF- β and the suppression of IFN- γ there is increased classic complement pathway (CCP) activity with cross talk to the KKS. There is no KKS crosstalk with either the alternative complement pathway or the lectin complement pathway(9). BKN (bradykinin) is the principal hormone of the KKS as angiotensin II is the principal hormone of the RAS. Estrogen down-regulates ACE, which degrades BKN. The subsequent angioedema appears to create the brain fog, post exertional malaise, and fatigue of Long Covid and its cousins.

The interface between insufficient magnesium and elevation of TGF- β in Long Covid begins with cellular hypoxia. Long Covid afflicts a younger age group, predominantly female, the opposite of the Covid-19 group. Perhaps more of the elderly males with Covid-19 died, affecting the gender and age breakdown for Long Covid. Perhaps it is due to decreased magnesium intake, more prevalent in females under 50, especially teens, and males over 50 (see figure1).




Figure 1. Magnesium deficiency is accentuated in females less than 50 and males over 50(10).

Or perhaps the hypoxia/lactate induced upregulation of TGF- β and consequent suppression of IFN- γ is the explanation? These last two are interrelated. If magnesium is insufficient, then the low grade inflammatory state has given TGF- β the upper hand over IFN- γ . Magnesium, as will be shown, is critical to the production of antioxidants.

2.2 Antioxidants

The primary problem for these chronic fatigue diseases appears to be a shortage of antioxidants aka mitochondrial optimizers or enhancers. Their inability to quench the increased ROS generated during a respiratory viral infection leads to oxidative stress and pressure on mitochondria.

Most endogenous antioxidants require methylation and SAdMe (S-adenosylmethionine) is the universal methyl donor. Magnesium is not only a required cofactor but also an ATP chelate for all SAdMe methylations, which occur in the mitochondria(11,12).

Glutathione, the master antioxidant, requires SAdMe, but does not cross the blood brain barrier. Each molecule of glutathione can be regenerated from several sources, e.g., alpha lipoic acid, cysteine, NAC, ..., but several ATPs are required for each pathway to glutathione synthesis. The currency of regeneration is in short supply under oxidative stress conditions.

Many antioxidants have been recommended for ME/CFS (13,14,15,16), FM(13,14,17), and EBV(18). Most of these are otherwise endogenously produced, but require methylation to attain active status. Those requiring methylation include melatonin, betaine, choline, cysteine, taurine, CoQ10, carnitine, creatine, creatinine, and lysine. Biochemical pathways for many of these are available(19). Others require just multiple ATPs, e.g., NADH, tryptophan. Unfortunately direct SAdMe supplementation has recently been shown to be counterproductive(20).

Vitamins B3,9,12 require methylations (Mg^{2+} as cofactor and ATP chelate) to attain active status. Vitamins B1,2,6 require phosphorylations (Mg^{2+} chelated to ATP) to attain this. In short, magnesium is critical to the synthesis of all endogenous antioxidants. However, some exogenous antioxidants have been suggested - D-ribose(21), zinc, quercetin, curcumin, resveratrol, selenium, zinc, vitamin C, all fat soluble vitamins (A,D,E,K), cannabinoids(22). D-Ribose - can create one ATP thru pentose phosphate shunt. One study suggested sodium as a nutritional supplement for CFS(23). Although not an antioxidant, this speaks to the likelihood of some degree of chronic dehydration in many with CFS(24). Sweat generates much more sodium loss than that of magnesium, but renal resorption of water can cause magnesuria.

This critical role for magnesium is inextricably entwined with serum Ca:Mg. The Western diet has seen this ratio escalate from 2.3-2.9 in 1977 to 2.9-3.5 in 2007(25) with a rise in magnesium deficiency .

2.3 POTS Paradox, Hypocortisolism, and Histamine

An elevated Ca:Mg and background oxidative stress may be responsible for the dysautonomic symptoms of POTS and the renin aldosterone paradox (low renin and aldosterone in the face of hypovolemia). Those with Long Covid and those with CFS have lower levels of cortisol(26,27), and those with POTS(28) in addition have inappropriately low aldosterone and renin in the face of hypovolemia. Aldosterone, corticosterone, and cortisol synthesis occur in mitochondria and all require 11-beta hydroxylase (see figure 2), i.e., ATP and magnesium are essential. Magnesium deficiency not only retards intramitochondrial aldosterone synthase but also increases Ca:Mg and the baroreflex blood pressure threshold => orthostatic hypotension(29).

In addition Mg^{2+} is cofactor for adenylyl/guanylyl cyclase and chelate for the substrate ATP/GTP in the synthesis of “second messengers” cAMP and cGMP. Endothelial cGMP and NO(nitric oxide) trigger renin secretion(30,31,32).

TGF- β is the master cytokine in the etiology of most chronic fatigue diseases, including Long Covid(39), CFS(40,41,42), EBV(43), CMV(44), CIRS (biotoxins from chronic mold exposure and chronic Lyme disease)(45).

TGF- β is elevated in Alzheimer's disease(46) and Alzheimer's disease progression is accelerated post Covid-19(47,48) and in EBV(49), CIRS(50), FM(51), and CMV(52).

2.5 TGF- β Channels and Inflammasomes

2.5.1.0 TRPM2 v TRPM5,7

The TGF- β connection between Long Covid and its chronic disease cousins has been demonstrated. An additional Alzheimer's disease connection has also been demonstrated. ROS and TRPMs are integral to neurodegenerative diseases(53,54) and the mechanism appears to involve a Ca:Mg imbalance in the Warburg microenvironment (mitochondrial dysfunction). TRPM channels mediate intracellular calcium and magnesium balance. TRPM2 is the calcium channel(55) and TRPM5,7 are the magnesium channels(56).




Figure 3. ROS facilitate dominance of TRPM2 and increase intracellular Ca^{2+} and mitochondrial dysfunction(53).

ROS induce TRPM2 activation in endothelial cells(55). Increased extracellular Ca:Mg and ROS facilitate TRPM2 activation, which promotes increased intracellular Ca^{2+} and a positive feedback loop with additional input from TRPM2 (see figure 3)(55,57).

Increased extracellular Mg^{2+} reverses the TRPM2 dominance over TRPM5,7 and reduces Ca^{2+} signaling in endothelial cells(58).

TRPM2 channels also contribute to the pathogenesis of inflammatory bowel disease(59).

A TRPM2 facilitated increase in intracellular Ca^{2+} leads to an assault on mitochondria via the permeability transition pore (see figure 4,5)(60,61).




Figure 4. Ca^{2+} enters mitochondria through the calcium uniporter and the mitochondrial permeability transition pore (mPT)⁽⁶⁰⁾.

F-ATP synthase dimers




Figure 5. Ca^{2+} competes with PTP-inhibitory Mg^{2+} and is an essential permissive factor for PTP (permeability transition pore) opening(61).

2.5.2.0 NLRP3 Inflammasome

TGF- β and the NLRP3 inflammasome are connected(62) and both of these are associated with the Warburg effect(63). The inflammasome is vital to the pathogenesis of ME/CFS(64), FM, Alzheimer's(65), IBD(66,67), autoimmune disease(68,69), EBV(70) and Long Covid(71). The Ca:Mg imbalance is at the root of all these diseases and magnesium therapy reverses the imbalance via CaSRs and TRPM5,7. The CaSR regulates the NLRP3 inflammasome(72) and can be downregulated by increasing Mg^{2+} (73,74,75). Calcium-sensing receptor (CASR) activates the NLRP3 inflammasome, mediated by increased intracellular Ca^{2+} and decreased cellular cyclic AMP (cAMP).

2.6 GABA

Glutamate is synthesized from TCA cycle substrates. GABA is synthesized directly from glutamate and requires cofactors P5P(pyridoxal-5-phosphate) and Mg^{2+} (see figure 7). Beta and gamma isomers of hydroxybutyrate (butyrate from “good“ intestinal bacteria) can replace glutamate(76,77) in the synthesis of GABA. The ketone body butyrate is hydroxylated in the liver to produce the isomers beta and gamma hydroxybutyrate. GABA cannot pass the blood brain barrier.



NMDA RECEPTOR ACTIVATION




Figure 6. Glutamate favors NMDA receptors. The Mg²⁺ plug closes the Ca²⁺ channel. Its removal opens the channel([78](#)).




Figure 7. GABA enabling roles for B6, Mg^{2+} , GABA supplements (butyrates), and probiotics are shown(79).

Mg^{2+} can bind to both NMDA receptors (see figure 6)(78) and GABAergic receptor sites(80). In addition P5P is a required cofactor for GAD (glutamic acid decarboxylase) and GABA synthesis (see figure 7). GABA plays a lead role in the gut-brain axis and probably determines the anosmia/ageusia, headaches, and depression of Long Covid. Post Covid anosmia/ageusia are thrice as likely in Caucasians versus Asians(81). Asians tend to exhibit lower serum Ca:Mg. Excitatory glutamatergic NMDA receptor activity reflects this ratio (see figure 6). Might this chemosensory dysfunction be due to excess glutamatergic tone(82,83) and not some olfactory bulb related etiology? Anosmia/ageusia may be symptoms of a minor seizure(84) that would otherwise have been inhibited by functioning GABAergic neurons. Gabapentin, an anticonvulsant and GABA analog, has been used to treat anosmia and ageusia. The other symptom frequently encountered in Long Covid is hair loss. This too is associated with magnesium deficiency(85). Calcium and magnesium share the same receptor, CaSR (calcium sensing receptor), and deliver the same message to the parathyroid glands to decrease PTH secretion. However, at low concentrations Mg^{2+} delivers the opposite message, i.e., to increase PTH secretion/vitamin D synthesis(86), exacerbating any pre-existing elevated Ca:Mg. Any attempt to rectify the ratio by increasing magnesium intake without addressing the calcium overage may quickly elicit the laxative effect.

2.7 HRV and the Gut-Brain-Heart/Lung Axis

HRV is the fifth vital sign and, like serum CRP(C reactive protein), is an early warning indicator of some health problem. Many studies have demonstrated a reinforcing relationship between CRP and HRV(87). Both can be excellent early indicators of deteriorating health, response to therapy, and prognosis on any behavioral, biologic, or epidemiological path(88). They offer high sensitivity but low specificity. The list of such detectable problems is quite comprehensive, e.g., cardiovascular disease(87), Covid-19(89), sudden cardiac death(90), seizures(91), Crohn's disease(92), ulcerative colitis(93), ME/CFS(94), depression(95).

The vagus nerve or the wandering nerve is the longest in the body. It links the ENS with the ANS and, when dysfunctional, is responsible for such diverse health issues as lone atrial fibrillation, orthostatic hypotension, diarrhea, and dysphagia. Many of these have been reported in Long Covid and a strong connection between vagus nerve dysfunction implicated(96).

Most, if not all, of these vagal correlations are due to a Ca:Mg imbalance or a glutamate/GABA imbalance, e.g., lone atrial fibrillation (97,98,99). Both imbalances are tightly linked and diet dependent. Intestinal “friendly” bacteria (Bifidobacterium pseudocatenulatum and F. prausnitzii) that produce butyrates were deficient in those with Long Covid(100).

2.8 Diet

Increased daily intake of omega-3 polyunsaturated fatty acids (DHA and EPA) significantly increased the density of bacteria that are known to produce butyrate(101,102). Aged cheese (up to 25 Ca:Mg) can aggravate Long Covid. Exogenous antioxidants that don't require additional energy to activate might be helpful in an energy challenged host. Careful attention to hydration is highly advisable, especially in the active and in the elderly, as the thirst reflex diminishes with age. The 30:1 gradient between intracellular K⁺ and that extracellular requires ATP and magnesium. If Mg²⁺ is low, then K⁺ is probably also low (lots of ectopic beats)(98).

The ketogenic diet, popular for weight loss, encourages dairy, but this increases Ca:Mg. The Mediterranean and Paleolithic diets, which encourage nuts and seeds (Mg²⁺ rich) and discourage dairy, might be better. But biologic individuality dictates an experimental approach.

2.9 Summary

In summary, Covid-19 severity is directly related to RAS activity. TGF- β , activated by angiotensin II type 1 receptors, is elevated in the elderly and those with comorbidities. Many of these never fully recover from the initial illness. Long Covid, characterized by symptoms such as brain fog, post exertional malaise, fatigue, anosmia, ageusia, headaches, hair loss, and many others, selects those with less RAS and more KKS. Magnesium, critical to the synthesis of endogenous antioxidants, is suboptimal. The virus that was at first asymptomatic or manifested few symptoms is not cleared(see figure 8).




Figure 8. PASC (posts-acute

sequelae of COVID-19) is associated with persistence of SARS Cov2(103).

Inflammation smolders. A Warburg microenvironment (mitochondrial dysfunction) develops. TGF- β is elevated in both groups. The primary risk factor for any illness including cancer is immune function, and antioxidant sufficiency is critical in this. The ability to properly methylate protein is paramount to preventing chronic inflammation. The ability to properly methylate DNA is paramount to preventing cancer. Following a diet that maintains the Ca:Mg between 1.7 and 2.6, one that preserves the excitatory glutamate/inhibitory GABA balance, one that slows the sympathetic takeover of the ANS is highly advisable. Cultivate some intestinal “friendlies” that produce butyrates. The efficacy of vitamin D is maximized with such a diet. The hydroxylations required to produce the active form of vitamin D, 1,25 (OH)₂ cholecalciferol, from either sunlight or D3 occur in mitochondria and the full efficacy of vitamin D is not realized outside this Ca:Mg range.

3.0 Conclusion

Long Covid and ME/CFS, FM, EBV, CMV, POTS, MCAS, CIRS, and post viral fatigue syndrome are linked by antioxidant deficiency, elevated TGF- β , and the Warburg effect. “Friendly” butyrate producing intestinal bacteria are in short supply. Butyrates rectify the glutamate-GABA imbalance. These two neurotransmitters determine autonomic tone via the vagus nerve, which reflects general health through the fifth vital sign, HRV. Perhaps the gut-brain axis should be expanded to include two other vital organs under vagal control - the gut-brain-heart/lung axis.

The slowly increasing Ca:Mg in the Western diet incriminates magnesium deficiency as a central player in the pathogenesis of most chronic fatigue diseases. With age there is a steady progression from parasympathetic to sympathetic tone, from GABAergic to glutamatergic predominance, and from a balanced Ca:Mg to a calcium predominant one, at least on a Western diet.

Improving Ca:Mg starts with knowing what it is. A comprehensive chem panel including serum calcium and magnesium in an otherwise healthy individual without renal disease or medication should provide this information (104). Addressing excess dietary calcium (Ca:Mg>2.6) first and then increasing magnesium intake might minimize any laxative effect. But for some, changing religion is easier than adjusting diet.

References

- 1 **The Warburg effect then and now: From cancer to inflammatory diseases**
<https://doi.org/10.1002/bies.201300084>
- 2 **Lactate Metabolism and Signaling in Tuberculosis and Cancer: A Comparative Review (2021)**
<https://doi.org/10.3389/fcimb.2021.624607>
- 3 **Relevance of the Warburg Effect in Tuberculosis for Host-Directed Therapy (2020)**
<https://doi.org/10.3389/fcimb.2020.576596>
- 4 **Transforming growth factor-beta profiles correlate with clinical symptoms and parameters of haemostasis and inflammation in a controlled human malaria infection (2020)**
<https://doi.org/10.1016/j.cyto.2019.154838>
- 5 **Etiology of lactic acidosis in malaria (2021)**
<https://doi.org/10.1371/journal.ppat.1009122>
- 6 **The Warburg Effect 97 Years after Its Discovery (2020)**
<https://doi.org/10.3390%2Fcancers12102819>
- 7 Lotz, M., and Zuraw, B.L. (1987) **IFN- γ Is a Major Regulator of C1-inhibitor Synthesis by Human Blood Monocytes.** The Journal of Immunology, 139, 3382-3387.
<https://pubmed.ncbi.nlm.nih.gov/3119706/>
- 8 Zuraw BL, Lotz M. **Regulation of the hepatic synthesis of C1 inhibitor by the hepatocyte stimulating factors interleukin 6 and interferon gamma.** J Biol Chem. 1990 Jul 25;265(21):12664-70.
[https://doi.org/10.1016/S0021-9258\(19\)38395-4](https://doi.org/10.1016/S0021-9258(19)38395-4)
- 9 **Long Covid, Short Magnesium**
<https://doi.org/10.4236/oalib.1108736>
- 10 **Grassrootshealth Nutrient Research Institute**
<https://www.grassrootshealth.net/blog/lack-magnesium-worsening-ability-handle-stress/>
- 11 **Protein methylation in mitochondria (2022)** <https://doi.org/10.1016%2Fj.jbc.2022.101791>
- 12 **Human METTL20 Methylates Lysine Residues Adjacent to the Recognition Loop of the Electron Transfer Flavoprotein in Mitochondria** <https://doi.org/10.1074/jbc.M114.580464>
- 13 **Mitochondria and the Future of Medicine: The Key to Understanding Disease, Chronic Illness, Aging, and Life Itself** by Lee Know
- 14 **Alternative medical interventions used in the treatment and management of myalgic encephalomyelitis/chronic fatigue syndrome and fibromyalgia (2010)**
<https://doi.org/10.1089/acm.2008.0376>
- 15 **A Comparative Study of Antifatigue Effects of Taurine and Vitamin C on Chronic Fatigue Syndrome**
<https://doi.org/10.4236/pp.2022.138023>
- 16 **Competition for glutathione precursors between the immune system and the skeletal muscle: pathogenesis of chronic fatigue syndrome**
<https://doi.org/10.1054/mehy.1998.0780>
- 17 **Creatine supplementation in fibromyalgia: a randomized, double-blind, placebo-controlled trial**
<https://doi.org/10.1002/acr.22020>

- 18 Glutathione peroxidase (GPx) and superoxide dismutase (SOD) activity in patients with diabetes mellitus type 2 infected with Epstein- Barr virus**
<https://doi.org/10.1371/journal.pone.0230374>
- 19 Homocysteine Metabolism: Nutritional Modulation and Impact on Health and Disease**, chapter 54, Alan L. Miller, ND, Gregory S. Kelly, ND, Jessica Tran, ND
<https://musculoskeletalkey.com/homocysteine-metabolism-nutritional-modulation-and-impact-on-health-and-disease/>.
- 20 S-adenosylmethionine inhibits methylation via catabolism to adenine (2022)** <https://doi.org/10.1038/s42003-022-03280-5>
- 21 Understanding D-Ribose and Mitochondrial Function** <http://dx.doi.org/10.7575/aiac.abcm.v6n.1p.1>
- 22 Synergistic and antagonistic antioxidant effects in the binary cannabinoids mixtures**
<https://doi.org/10.1016/j.fitote.2021.104992>
- 23 Chronic fatigue syndrome (CFS): Suggestions for a nutritional treatment in the therapeutic approach (2019)**
<https://doi.org/10.1016/j.biopha.2018.10.076>
- 24 Primary renal magnesium wasting: an unusual clinical picture of exercise-induced symptoms**
<https://doi.org/10.14814/phy2.12773>
- 25 Rising Ca:Mg intake ratio from food in USA Adults: a concern?**
<http://mgwater.com/Ca-Mg.pdf>
- 26 Distinguishing features of Long COVID identified through immune profiling (preprint August 10, 2022)**
<https://doi.org/10.1101/2022.08.09.22278592>
- 27 The Hypocortisolism in Chronic Fatigue Syndrome (ME/CFS) – Artifact or Central Factor?**
<https://phoenixrising.me/myalgic-encephalomyelitis-chronic-fatigue-syndrome-mecfs-research/pharmacogenomics/the-hypocortisolism-in-chronic-fatigue-syndrome-mecfs-artifact-or-central-factor/>
- 28 Postural orthostatic tachycardia syndrome is an under-recognized condition in chronic fatigue syndrome**
<https://doi.org/10.1093/qjmed/hcn123>
- 29 Calcium and Magnesium Sensitivity of the Carotid Baroreceptor Reflex in Cats**
<https://doi.org/10.1161/01.RES.45.6.815>
- 30 Nitric oxide synthase and cGMP-mediated stimulation of renin secretion**
<https://doi.org/10.1152/ajpregu.2001.281.4.R1146>
- 31 Effects of magnesium on nitric oxide synthase activity in endothelial cells**
<https://doi.org/10.1152/ajpcell.1995.269.3.C612>
- 32 Endothelium-Derived Nitric Oxide Supports Renin Cell Recruitment Through the Nitric Oxide–Sensitive Guanylate Cyclase Pathway**
<https://doi.org/10.1161/HypertensionAHA.111.00221>
- 33 Hydroxysteroid Dehydrogenases – Biological Role and Clinical Importance – Review**
<https://doi.org/10.5772/54149>
- 34 Pinto, M.D., Lambert, N., Downs, C.A., Abraham, H., Hughes, T.D. and Rahmani, A.M. (2022) Antihistamines for Post Acute Sequelae of SARS-CoV-2 Infection.** The Journal for Nurse Practitioners, 18, 335-338.
<https://doi.org/10.1016/j.nurpra.2021.12.016>
- 35 The Many Talents of TGF-β in the Kidney**
<https://doi.org/10.1097%2FNMNH.0000000000000490>
- 36 TGF-β receptors: In and beyond TGF-β signaling (2018)**
<https://doi.org/10.1016/j.cellsig.2018.09.002>

- 37 **TGFBR3 is an independent unfavourable prognostic marker in oesophageal squamous cell cancer and is positively correlated with Ki-67 (2020)**
<https://doi.org/10.1111/iep.12380>
- 38 **Promoter Hypomethylation of TGFBR3 as a Risk Factor of Alzheimer's Disease: An Integrated Epigenomic-Transcriptomic Analysis (2021)**
<https://doi.org/10.3389/fcell.2021.825729>
- 39 **A Review of Persistent Post-COVID Syndrome (PPCS)**
<https://doi.org/10.1007/s12016-021-08848-3>
- 40 **Cytokine signature associated with disease severity in chronic fatigue syndrome patients (July 2022)**
<https://doi.org/10.1073/pnas.1710519114>
- 41 **Activin B is a novel biomarker for chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME) diagnosis: a cross sectional study**
<https://doi.org/10.1186/s12967-017-1161-4>
- 42 **Up-regulation of TGF- β 1 mRNA expression in peripheral blood mononuclear cells of patients with chronic fatigue syndrome**
<https://doi.org/10.1016/j.jfma.2011.09.006>
- 43 **Transforming growth factor beta-induced reactivation of Epstein-Barr virus involves multiple Smad-binding elements cooperatively activating expression of the latent-lytic switch BZLF1 gene**
<https://doi.org/10.1128/jvi.01197-10>
- 44 **Cytomegalovirus production by infected astrocytes correlates with transforming growth factor-beta release**
<https://doi.org/10.1086/373995>
- 45 **Chronic Inflammatory Response Syndrome (CIRS) Evaluation and Treatment (2021)**
<https://hoffmancentre.com/chronic-inflammatory-response-syndrome-cirs-evaluation-and-treatment/>
- 46 **TGF- β 1 factor in the cerebrovascular diseases of Alzheimer's disease**
X. Zhang, W.-J. Huang, W.-W. Chen, Eur Rev Med Pharmacol Sci 2016 20(24):5178-5185
<https://pubmed.ncbi.nlm.nih.gov/28051272/>
- 47 **Association of COVID-19 with New-Onset Alzheimer's Disease (2022)**
<https://doi.org/10.3233/JAD-220717>
- 48 **COVID-19 and Alzheimer's disease: how one crisis worsens the other (2021)**
<https://doi.org/10.1186/s40035-021-00237-2>
- 49 **Herpes virus in Alzheimer's disease: relation to progression of the disease**
<https://doi.org/10.1016/j.neurobiolaging.2013.06.024>
- 50 **Inhalational Alzheimer's disease: an unrecognized—and treatable—epidemic**
<https://doi.org/10.18632/faging.100896>
- 51 **Fibromyalgia and Risk of Dementia-A Nationwide, Population-Based, Cohort Study**
<https://doi.org/10.1016/j.amjms.2017.09.002>
- 52 **Cytomegalovirus Infection and Risk of Alzheimer Disease in Older Black and White Individuals**
<https://doi.org/10.1093/finfdis/fjiu437>
- 53 **Transient Receptor Potential Channels as an Emerging Target for the Treatment of Parkinson's Disease: An Insight Into Role of Pharmacological Interventions. (figure 2)**
<https://doi.org/10.3389/fcell.2020.584513>

- 54 **TRPM2: a candidate therapeutic target for treating neurological diseases**
<https://doi.org/10.1038/aps.2018.31>
- 55 **Reactive Oxygen Species-Induced TRPM2-Mediated Ca²⁺ Signalling in Endothelial Cells.**
<https://doi.org/10.3390/antiox10050718>
- 56 **Magnesium Regulates Endothelial Barrier Functions through TRPM7, MagT1, and S1P1**
<https://doi.org/10.1002/adv.201901166>
- 57 **Regulation of TRPM2 by Extra- and Intracellular Calcium**
<https://doi.org/10.1085/jgp.200709836>
- 58 **Influences of Calcium and Magnesium Ions on Cellular Antioxidant Activity (CAA) Determination**
<http://dx.doi.org/10.1016/j.foodchem.2020.126625>
- 59 **TRP channels in inflammatory bowel disease: Potential therapeutic targets**
<https://doi.org/10.1016/j.bcp.2022.115195>
- 60 Figure 2 at **Mitochondrial pharmacology**
<http://dx.doi.org/10.1016/j.tips.2012.03.010>
- 61 Figure 1 at **Calcium and regulation of the mitochondrial permeability transition**
<https://doi.org/10.1016/j.ceca.2017.05.004>
- 62 **TGF- β activates NLRP3 inflammasome by an autocrine production of TGF- β in LX-2 human hepatic stellate cells**
<https://doi.org/10.1007/s11010-022-04369-5>
- 63 **The Warburg effect promoted the activation of the NLRP3 inflammasome induced by Ni-refining fumes in BEAS-2B cells (2020)**
<https://doi.org/10.1177/0748233720937197>
- 64 **Activation of the NLRP3 inflammasome in lipopolysaccharide-induced mouse fatigue and its relevance to chronic fatigue syndrome**
<https://jneuroinflammation.biomedcentral.com/articles/10.1186/s12974-016-0539-1>
- 65 **NLRP3 Inflammasome in Neurological Diseases, from Functions to Therapies**
<https://doi.org/10.3389/fncel.2017.00063>
- 66 **NLRP3 Inflammasome and Inflammatory Bowel Disease**
<https://doi.org/10.3389/fimmu.2019.00276>
- 67 **NLRP3 Inflammasome-Mediated Inflammatory Process in Patients with Irritable Bowel Syndrome**
<https://www.dissertationtopic.net/doc/663467>
- 68 **Microglial NLRP3 inflammasome activation in multiple sclerosis**
<https://doi.org/10.1016/bs.apcsb.2019.08.007>
- 69 **NLRP3: A promising therapeutic target for autoimmune diseases**
<https://doi.org/10.1016/j.autrev.2018.01.020>
- 70 **The danger molecule HMGB1 cooperates with the NLRP3 inflammasome to sustain expression of the EBV lytic switch protein in Burkitt lymphoma cells (2022)**
<https://doi.org/10.1016/j.virol.2021.12.002>
- 71 **NLRP3 inflammasome: A joint, potential therapeutic target in management of COVID-19 and fertility problems (2021)**
<https://doi.org/10.1016/j.jri.2021.103427>
- 72 **The calcium-sensing receptor regulates the NLRP3 inflammasome through Ca²⁺ and cAMP**

<https://doi.org/10.1038%2Fnature11588>

73 Effects of MgSO₄ on inhibiting Nod-like receptor protein 3 inflammasome involve decreasing intracellular calcium

<https://doi.org/10.1016/j.jss.2017.09.005>

74 Magnesium isoglycyrrhizinate blocks fructose-induced hepatic NF- κ B/NLRP3 inflammasome activation and lipid metabolism disorder

<https://doi.org/10.1016/j.ejphar.2017.05.032>

75 Magnesium lithospermate B acts against dextran sodium sulfate-induced ulcerative colitis by inhibiting activation of the NLRP3/ASC/Caspase-1 pathway

<https://doi.org/10.1016/j.etap.2015.10.009>

76 β -Hydroxybutyrate is the preferred substrate for GABA and glutamate synthesis while glucose is indispensable during depolarization in cultured GABAergic neurons <https://doi.org/10.1016/j.neuint.2011.06.002>

77 Gamma-Hydroxybutyrate modulates synthesis and extracellular concentration of gamma-aminobutyric acid in discrete rat brain regions in vivo

<https://pubmed.ncbi.nlm.nih.gov/10381791/>

78 Ketamine and Esketamine in depression - A Synopsis

<https://psychscenehub.com/psychinsights/ketamine-and-depression/>

79 Gamma-aminobutyric acid (GABA) Monograph

<https://www.fxmedicine.com.au/blog-post/gamma-aminobutyric-acid-gaba-monograph>

80 Magnesium potentiation of the function of native and recombinant GABA(A) receptors

<http://dx.doi.org/10.1097/00001756-200107200-00026>

81 von Bartheld, C. S., Hagen, M. M., & Butowt, R. (2020). Prevalence of chemosensory dysfunction in COVID-19 patients: A systematic review and meta-analysis reveals significant ethnic differences. ACS Chemical Neuroscience, 11(19), 2944–2961.

<https://doi.org/10.1021/acscemneuro.0c00460>

82 Levy, Lucien M., and Robert I. Henkin. "Brain Gamma-Aminobutyric Acid Levels Are Decreased in Patients With Phantageusia and Phantosmia Demonstrated by Magnetic Resonance Spectroscopy." Journal of Computer Assisted Tomography 28, no. 6 (November/December 2004): 721–27.

<https://doi.org/10.1097/00004728-200411000-00001>

83 Henkin, R.I., "Treatment of Distortions of Taste and Smell," 2006. <http://www.tasteandsmell.com/sep06.htm>.

84 Glutamatergic Mechanisms Associated with Seizures and Epilepsy

<https://doi.org/10.1101/cshperspect.a022863>

85 A Tataru and E Nicoara. Idiopathic diffuse alopecias in young women correlated with hypomagnesemia. J Eur Acad Dermatol Venereol. 2004 May;18(3):393-4.

<https://doi.org/10.1111/j.1468-3083.2004.00660.x>

86 Calcium, magnesium and the control of PTH secretion [https://doi.org/10.1016/0169-6009\(89\)90003-2](https://doi.org/10.1016/0169-6009(89)90003-2)

87 The relationship between heart rate variability and inflammatory markers in cardiovascular diseases

<https://doi.org/10.1016/j.psyneuen.2008.08.007>

88 The Vagus Nerve Can Predict and Possibly Modulate Non-Communicable Chronic Diseases: Introducing a Neuroimmunological Paradigm to Public Health

<https://doi.org/10.3390%2Fjcm7100371>

89 Heart-rate-variability (HRV), predicts outcomes in COVID-19

<https://doi.org/10.1371/journal.pone.0258841>

90 Heart rate variability as predictive factor for sudden cardiac death

<https://doi.org/10.18632/aging.101386>

91 Prediction of epileptic seizures based on heart rate variability

<https://doi.org/10.3233/thc-161225>

92 Autonomic Dysfunction Correlates with Clinical and Inflammatory Activity in Patients with Crohn's Disease

<https://doi.org/10.1097/mib.0000000000000508>

93 Longitudinal Autonomic Nervous System Measures Correlate With Stress and Ulcerative Colitis Disease Activity and Predict Flare (2021)

<https://doi.org/10.1093/ibd/izaa323>

94 Reduced heart rate variability predicts fatigue severity in individuals with chronic fatigue syndrome/myalgic encephalomyelitis (2020)

<https://doi.org/10.1186/s12967-019-02184-z>

95 Heart Rate Variability for the Prediction of Treatment Response in Major Depressive Disorder

<https://doi.org/10.3389/fpsyt.2020.00607>

96 Pilot study suggests long COVID could be linked to the effects of SARS-CoV-2 on the vagus nerve (2022)

<https://www.eurekalert.org/news-releases/943102>

97 'Lone' atrial fibrillation precipitated by monosodium glutamate and aspartame

<https://doi.org/10.1016/j.ijcard.2009.01.028>

98 Magnesium and Potassium in Lone Atrial Fibrillation

<http://www.mgwater.com/laf.shtml>

99 Lone atrial fibrillation: Pathologic or not?

<https://doi.org/10.1016/j.mehy.2006.07.030>

100 Gut microbiota composition reflects disease severity and dysfunctional immune responses in patients with COVID-19 (2021)

<https://gut.bmj.com/content/70/4/698>

101 Associations among Dietary Omega-3 Polyunsaturated Fatty Acids, the Gut Microbiota, and Intestinal Immunity (2021)

<https://doi.org/10.1155%2F2021%2F8879227>

102 Low omega-3 index and polyunsaturated fatty acid status in patients with chronic fatigue syndrome/myalgic encephalomyelitis

<https://doi.org/10.1016/j.plefa.2018.11.00>

103 Persistent circulating SARS-CoV-2 spike is associated with post-acute COVID-19 sequelae, Zoe Swank, Yasmee Senussi, Galit Alter, David R. Walt *Zoe Swank, Clinical Infectious Diseases, 2022, ciac722, 2022.06.14.22276401*

<https://doi.org/10.1093/cid/ciac722>

104 Ca:Mg + D, the Shield that Interdicts the Crown Viruses and Vaccines

<https://doi.org/10.4236/oalib.1109249>