



## Review

## A glimpse of vitamin D status in Mainland China

Weiguo Zhang M.D., Ph.D.<sup>\*</sup>, Elisabeth Stoecklin M.D., Manfred Eggersdorfer Ph.D.

DSM Nutritional Products, Human Nutrition and Health, Beijing, China and Kaiseraugst, Switzerland

## ARTICLE INFO

## Article history:

Received 12 November 2012

Accepted 11 January 2013

## Keywords:

Blood

25-Hydroxyvitamin D

Bone

Nutrient deficiency

Nutritional status

Population

Public health

## ABSTRACT

As an essential dietary micronutrient, vitamin D plays a pivotal role in promoting calcium absorption in the intestine and maintaining a healthy skeletal system throughout life. Beyond bone health, an emerging volume of scientific studies shows that vitamin D also may provide cardiovascular, metabolic, and immunologic benefits and reduce mortality. To our knowledge, in mainland China no national surveys have been conducted to date to depict the overall vitamin D status in the population. Therefore, the purpose of this contribution was to provide the best possible evaluation of vitamin D deficiency/insufficiency in China by reviewing publications that measured plasma/serum 25-hydroxyvitamin-D (25[OH]D) levels in various age groups and in different areas of China from January 2000 to June 2012. From these investigations conducted throughout the country and from newborns to adults to the elderly, it has been found that vitamin D deficiency/insufficiency is prevalent in the Chinese population in almost all age groups and areas if individuals are not taking vitamin D–fortified products/supplements or are lacking sufficient sunshine exposure. Some studies showed severe deficiency (25[OH]D <25 nmol/L) in Nanjing (north latitude 31) during the winter months and in Beijing (north latitude 40) in the fall. This unoptimistic situation represents a significant but modifiable public health risk that deserves greater attention and more efficient and timely management.

© 2013 Elsevier Inc. All rights reserved.

## Introduction

Vitamin D is an essential dietary micronutrient because it plays a pivotal role in aiding intestinal absorption of calcium and phosphorus, therefore, exerting a favorable effect on bone mineralization and musculoskeletal health. Other than the classical effects of vitamin D on calcium and phosphorus metabolism and on bone health and muscle strength, an emerging volume of scientific studies also shows that vitamin D deficiency/insufficiency may increase the risks for developing non-skeletal disorders such as cardiovascular disease, type 2 diabetes, autoimmune disease, selected cancers, high blood pressure, depression, and overall mortality [1,2].

In Mainland China, there has, to our knowledge, been no systematic national survey to depict the landscape of vitamin D deficiency, insufficiency, and adequacy. However, research interests in the topic developed and continued to mount over the past decade(s), and as a result, circulating vitamin D status has been examined in various populations and regions.

The purpose of this contribution is to provide best possible overview from these studies that measured circulating

25-hydroxyvitamin-D (25[OH]D) levels in Mainland China. Serum or plasma 25(OH)D concentration is widely accepted as the best biomarker to define vitamin D status, because it reflects both the endogenous vitamin D photosynthesis by skin in response to sun exposure and vitamin D ingestion via digestive track from dietary sources [3]. To this end, peer-reviewed original English publications from January 2000 to November 2012, which examined 25(OH)D values, were retrieved from databases and analyzed. Studies were not included when the number of participants was less than 30 and when they had known overt vitamin D intake. A deficient vitamin D level is defined as being 25(OH)D <25 nmol/L; insufficient <50 nmol/L; desirable >75 nmol/L (as recommended by International Osteoporosis Foundation [IOF], the Endocrine Society, and many scientists in the field) [4]. Therefore, levels between 50 nmol/L and 75 nmol/L were defined as inadequate (although they are defined as adequate by the Institute of Medicine). These cut-off points are also in line with the recently published global vitamin D status map by the IOF [5] (<http://www.iofbonehealth.org/facts-and-statistics/vitamin-d-studies-map>).

## Vitamin D status in the youth

As shown in Table 1 [6–12], vitamin D deficiency was phenomenal in the youth. There was no group with desirable

<sup>\*</sup> Corresponding author. Tel.: +86 10 5976 1028; fax: +86 5976 1088.

E-mail address: [weiguo.zhang@dsm.com](mailto:weiguo.zhang@dsm.com) (W. Zhang).

**Table 1**  
Vitamin D status in the youth

Publication	Age (yrs)	Number	Place	Latitude (north)	Season	D Use	Assay method	25-OH-D (nmol/l)	Deficiency (%)				
									<12.5	<25	<37.5	<50	<75
Wang J. 2010	Newborn	77 (MF)	Chengdu	30	Sept	Unknown	Enzyme immunoassay	<b>41.0 ± 18.9</b>		44.2			96.1
Song S. 2012	Newborn	58 (MF)	Beijing	40	Apr-May	No	ELISA	<b>27.9 ± 1.6*</b>	46.6		93.2	100	
Zhu Z. 2012	6-11	1440 (MF)	Hangzhou	30	All	Unknown	ELISA	<b>56.1 ± 19.9</b>	2.0		40.3	88.3	
Du X. 2001	12-16	183 (MF)	Hangzhou	30	All	Unknown	ELISA	<b>52.1 ± 17.0</b>	3.3		46.4	89.6	
	12.7	108 (F)	Beijing	40	Jan	Unknown	Competitive protein assay	<b>13.9 ± 9.6</b>	42.5				
					Sept-Oct	Unknown	Competitive protein assay	<b>30.2 ± 11.9</b>	5.1				
	13.0	57 (F)	Beijing	40	Jan	Unknown	Competitive protein assay	<b>12.7 ± 5.9</b>	49.6				
					Sept-Oct	Unknown	Competitive protein assay	<b>24.7 ± 10.6</b>	6.6				
	13.2	64 (F)	Beijing	40	Jan	Unknown	Competitive protein assay	<b>12.8 ± 6.7</b>	45.1				
					Sept-Oct	Unknown	Competitive protein assay	<b>23.8 ± 8.7</b>	9.2				
Foo L. 2009	15.0	301 (F)	Beijing	40	Mar-Apr	No	RIA	<b>34.0</b>	32.8	68.4	89.2		
Arguelles L. 2009	16.4	226 (FM)	Anqing	31	All	Unknown	HPLC	<b>45.0 ± 23.5</b>					90.3

M, male; F, female; All, the blood samples were collected in all seasons

The 25-OH-D values are expressed in mean ± SD (or mean ± SE when with \*) or median, and are converted if not in nmol/l. Different deficiency/insufficiency cutoffs and rates were provided by investigators

25(OH)D levels and as many as 40% to 90% had blood 25(OH)D levels lower than 50 nmol/L. A study in Beijing showed even worse vitamin D status: More than 40% of school-aged girls had 25(OH)D levels below 12.5 nmol/L in January. This significant vitamin D deficiency remained in 5% to 9% of them even in September and October [9].

Poor vitamin D status was not seen in participants taking vitamin D-fortified products or supplementation (Table 2A) [8, 13], unequivocally suggesting that vitamin D supplementation improved 25(OH)D status. Additionally, exposure to sunshine, thus allowing individuals to receive ultraviolet B (UVB), in the summer months improved 25(OH)D status in the same children who were previously deficient (Table 2B) [14]. Consequently, clinical rickets declined from 41.6% to 17%.

### Vitamin D status in adults

A larger percentage of adults had blood 25(OH)D levels below 50 nmol/L (Table 3) [6,7,15–27]. The best 25(OH)D levels (67.2 nmol/L) in adult groups were reported in Jinan [22]; whereas the worst (<25 nmol/L) was found in pregnant women in Nanjing and pregnant women with gestational diabetes mellitus in Beijing [15,20].

Surprisingly, in women of childbearing age, a lower 25(OH)D level also was found in lower latitude regions (29 nmol/L in

Beijing versus 34 nmol/L in Hong Kong where the latitude is 22 north) [18].

### Vitamin D status in the elderly

The elderly are fourfold less efficient in cutaneous vitamin D photosynthesis than youth [28,29] and are particularly vulnerable to bone fracture or mobility disorders. The 25(OH)D status was alarming in the fast-growing aging Chinese population (Table 4) [17,30–36]. There was no any study showing average 25(OH)D above 50 nmol/L. In two large-scale investigations in Beijing and Shanghai, as high as 70% to 90% of the participants had blood 25(OH)D levels below 50 nmol/L [30,35]. In 1460 elderly urban Shanghai residents, only 3.9% had plasma 25(OH)D levels >75 nmol/L [31]. Oral cholecalciferol (vitamin D<sub>3</sub>) at 925 IU/d, raised serum 25(OH)D from 40 nmol/L to 50 nmol/L in 3 mo in 45 postmenopausal women [34].

### Discussion

The data presented in this contribution are likely epitomic for the nation because the investigations were conducted in the east part of Hu's line (Fig. 1), which has been with <40% of the land and >90% of the population since 1930s [37]. In the global map of vitamin D deficiency/insufficiency [3,38], these data from China

**Table 2**  
Vitamin D status can be improved in children

A. Effect of supplementation											
Publication	Age (yrs)	Number	Place	Latitude (north)	Season	D Use	Assay method	25-OH-D (nmol/l)	Deficiency (%)		
									<25	<50	<75
Zhu Z. 2012	0-1	2116 (MF)	Hangzhou	30	All	Yes	ELISA	<b>98.7 ± 47.1</b>	0.4	5.4	33.6
	2-5	2269 (MF)	Hangzhou	30	All	Yes	ELISA	<b>69.6 ± 30.4</b>	1.1	21.9	68.6
Liang GY. 2011	0-10	76 (MF)	Nanjing	32	Nov-Mar	Yes	EIA	<b>80.5 ± 29.3</b>	1.3	10.5	
	0-10	66 (MF)	Nanjing	32	Nov-Mar	Yes	EIA	<b>65.7 ± 32.3*</b>	16.7	30.8	
B. Effect of sunshine exposure											
Publication	Age (month)	Number	Place	Latitude (north)/Elevation (meter)	Season	D Use	Assay method	25-OH-D (nmol/l)	Deficiency (%)		
									<12.5	<30	<50
Strand MA. 2009	18.2	177 (MF)	Yuci	37/797	Apr	Unknown	Radioimmunoassay	<b>34.5 ± 74.3</b>	33.5	65.3	84.3
	23.3	172 (MF)	Yuci	37/797	Sept	4.2%	Radioimmunoassay	<b>127.8 ± 143.5</b>	0	2.9	8.1

For footnote: please see Table 1

\* With respiratory infection.

**Table 3**  
Vitamin D status in adults

Publication	Age (yrs)	Number	Place	Latitude (north)	Season	D Use	Assay method	25-OH-D (nmol/l)	Deficiency (%)			
									<25	<50	<75	<80
Jiang L. 2012	26.3	78 (F)	Nanjing	31	Winter	99 IU	ELISA	<b>22.6 ± 12.7</b>	65.8	96.1		
Woo J. 2008	27.7	78 (F)	Nanjing	31	Summer	117 IU	ELISA	<b>31.8 ± 9.2</b>	22.4	94.7		
	26.9	220 (F)	Beijing	40	Feb-May	No	Radio-immunoassay	<b>29</b>	40	94		
Shao H. 2012	27.9	221 (F)	Hong Kong	22	Feb-May	No	Radio-immunoassay	<b>34*</b>	18	92		
	26.8	130 (F)	Shanghai	31	Jul-Dec	No	Chemiluminescence	<b>30.2<sup>†</sup></b>				
Wang J. 2010	27.3	139 (F)	Shanghai	31	Jul-Dec	No	Chemiluminescence	<b>29.7</b>				
	End pregnancy	77 (F)	Chengdu	30	Sept	Unknown	Enzyme immunoassay	<b>36.0 ± 18.9<sup>†</sup></b>		97.4		
Tao M. 2012	28.1	1695 (F)	Shanghai	31	All	No	Chemiluminescence	<b>43.9 ± 28.6<sup>†</sup></b>	69.0	91.0		
Yang B. 2012	29.4	41 (M)	Xi'an	34	All	Unknown	ELISA	<b>52.5 ± 15.9</b>				
	30.3	314 (M)	Xi'an	34	All	Unknown	ELISA	<b>53.3 ± 14.5</b>				
	30.5	195 (M)	Xi'an	34	All	Unknown	ELISA	<b>54.1 ± 14.3</b>				
Song S. 2012	29.9	70 (F)	Beijing	40	Apr-May	No	ELISA	<b>28.6 ± 1.4<sup>†</sup></b>	54.3	90.2	100	
Yan L. 2000	30.9	48 (F)	Shenyang	42	Apr-May	No	Radio-immunoassay	<b>40.7 ± 14.1</b>	13.0			
	31.1	48 (M)	Shenyang	42	Mar-May	No	Radio-immunoassay	<b>31.4 ± 10.4</b>	29.0			
Wang O. 2012	31.0	200 (F)	Beijing	40	All	Unknown	ELISA	<b>25.9<sup>†</sup></b>				
	32.0	200 (F)	Beijing	40	All	Unknown	ELISA	<b>22.4<sup>‡</sup></b>	53.8	96.3		
Lu H. 2012	43.0	2588 (MF)	Shanghai	31	Feb-Mar	No	Chemiluminescence	52.2		30 (M)	84 (M)	
										46 (F)	89 (F)	
Yin X. 2012	49.4	601 (MF)	Jinan	37	Nov-Dec	No	RIA	<b>67.2 ± 25.3</b>	28.6	66		
Lin S. 2012	56.5	1101 (MF)	Linxian	36	Spring	No	Enzyme immunoassay	<b>31.7</b>				
Li L. 2012	Adults	1420 (MF)	Dali	25	Mar-May	No	RIA	<b>54.9</b>				
Ren C. 2012	Adults	197 (MF)	Guangzhou	23	All	No	ELISA	<b>48.9 ± 23.7<sup>§</sup></b>	57.9	91.9		
Huang Y. 2012	Adults	49 (F)	Shanghai	31	Jan-Apr	No	Chemiluminescence	<b>29.4</b>				

Footnote: please see Table 1

The elevation in Dali is 2007 meters

\* The data from Hong Kong were used for head-to-head comparison with Beijing.

† Pregnancy.

‡ Pregnancy with gestational diabetes.

§ Gastric cancer.

would stand for a sizable geographic area and the largest population.

Vitamin D deficiency/insufficiency in China might have been caused by multiple factors, some of which are unique and worthy of being recognized. First, with the rapid transformation from an agrarian to an industrialized society, the urban proportion of population increased from 33% in 2000 to 50% in 2013 (the fifth and sixth National Population Survey <http://www.stats.gov.cn>). Accordingly, the number of people engaged in an outdoor profession with sun exposure has dramatically decreased. Second, due to air pollution accompanied by industrialization and urbanization, people were discouraged from spending time outdoors and UVB was prevented from penetrating the atmosphere; Third, the preference for lighter skin color (fair skin) in

the society remains unchanged, leading to popular use of hats, umbrellas, and sunscreens when outdoors. UVB radiation-related skin carcinomas also are increasingly a concern. Fourth, despite massive and successful improvements in macronutrient intakes in the Chinese population, micronutrient intake did improve in parallel [39]. In fact, natural food rich in vitamin D in the Chinese diet remain scarce, and food and beverages fortified with vitamin D are limited in the market.


Over the past several decades, the life span and life expectancy of the Chinese people has increased significantly, which brought the consequences of vitamin D deficiency, like osteoporosis and bone fracture, into greater prominence [40]. Accompanied by poor 25(OH)D status, a bone mineral density (BMD) survey with dual-energy X ray absorptiometry (DXA) in

**Table 4**  
Vitamin D status in the elderly

Publication	Age (yrs)	Number	Place	Latitude (north)	Season	D Use	Assay method	25-OH-D (nmol/l)	Deficiency (%)			
									<25	<50	<75	<80
Lu L. 2009	50-70	3262 (MF)	Shanghai	31	Apr-Jun	Unknown	Radio-immunoassay	<b>40.4</b>		69.2		
Dorjgochoo T. 2012	61	1460 (MF)	Shanghai	31	All	Unknown	chemiluminescence	<b>34.7</b>		96.1		
	61.5	32 (F)	Beijing	40	Feb-May	No	Chemiluminescence	<b>33.1 ± 15.5</b>				
Kruger M. 2012	62.7	31 (F)	Beijing	40	Feb-May	No	Chemiluminescence	<b>29.3 ± 12.0</b>				
	62.1	181 (MF)	Beijing	40	All	No	ELISA	<b>35.8 ± 12.4</b>	20.4	86.1	100	
Zhou X. 2012	63.6	193 (MF)	Beijing	40	All	No	ELISA	<b>32.1 ± 10.8*</b>	21.2	94.3	100	
	63.8	100 (F)	Shanghai	31	Dec-Mar	<600IU	Chemiluminescence	<b>42.0 ± 13.5</b>				
Zhang H. 2012	64.1	1724 (F)	Beijing	40	Unknown	Unknown	Chemiluminescence	<b>33.0 ± 13.5</b>		89.7 99.4		
Yan L. 2003	65.2	110 (F)	Shenyang	42	Feb-Apr	14.5%	Radio-immunoassay	<b>30.9 ± 13.5</b>	39.1			
Yan L. 2000	67.9	108 (M)	Shenyang	42	Feb-Apr	9.3%	Radio-immunoassay	<b>27.1 ± 11.5</b>	52.8			
	66.9	48 (F)	Shenyang	42	Apr-May	No	Radio-immunoassay	<b>42.9 ± 21.2</b>	15.0			
	68.9	50 (M)	Shenyang	42	Mar-May	No	Radio-immunoassay	<b>28.4 ± 12.5</b>	48.0			

Footnote: please see Table 1

\* With COPD.



**Fig. 1.** Distribution of studies. In some places like Beijing and Shanghai, multiple studies were conducted. The original blank map is from [http://en.wikipedia.org/wiki/File:China\\_edcp\\_relief\\_location\\_map.jpg#file](http://en.wikipedia.org/wiki/File:China_edcp_relief_location_map.jpg#file). The map is licensed under Creative Commons ShareAlike 3.0 <http://creativecommons.org/licenses/by-sa/3.0/deed.en>.

10 Chinese cities estimated that the proportion of osteoporosis in men and women over age 50 y was 10.4% and 31.2%, respectively [41]. Furthermore, the incidence of hip fracture in Beijing in the period 2002 to 2006 escalated much more rapidly compared with 1990 to 1992 [42]. Vitamin D deficiency increases the risk for both osteoporosis and falls [43,44], which are the two major risks for bone fracture.

In summary, from published investigations in Mainland China since the year 2000, vitamin D deficiency/insufficiency was found to be widely prevalent, which constitutes a significant but modifiable public health risk that deserves greater awareness and more efficient and timely management. To attain a desirable vitamin D status at the population level, it requires multiple approaches (e.g., promotion of a healthy lifestyle, implementation of recommended daily intake, development of voluntary and mandatory fortification programs, and regulation establishment and reinforcement).

## References

- [1] Zittermann A, Gummert JF. Nonclassical vitamin D action. *Nutrients* 2010;2:408–25.
- [2] Schottker B, Ball D, Gellert C, Brenner H. Serum 25-hydroxyvitamin D levels and overall mortality. A systematic review and meta-analysis of prospective cohort studies. *Ageing Res Rev*; 2012 Feb 17. <http://dx.doi.org/10.1016/j.arr.2012.02.004> [Epub ahead of print].
- [3] Troesch B, Hoefft B, McBurney M, Eggersdorfer M, Weber P. Dietary surveys indicate vitamin intakes below recommendations are common in representative Western countries. *Br J Nutr* 2012;108:692–8.
- [4] Vieth R. Why the minimum desirable serum 25-hydroxyvitamin D level should be 75 nmol/L (30 ng/ml). *Best Pract Res Clin Endocrinol Metab* 2011;25:681–91.
- [5] Wahl DA, Cooper C, Ebeling PR, Eggersdorfer M, Hilger J, Hoffmann K, et al. A global representation of vitamin D status in healthy populations. *Arch Osteoporos* 2012;7:155–72.
- [6] Wang J, Yang F, Mao M, Liu DH, Yang HM, Yang SF. High prevalence of vitamin D and calcium deficiency among pregnant women and their newborns in Chengdu, China. *World J Pediatr* 2010;6:265–7.
- [7] Song SJ, Si S, Liu J, Chen X, Zhou L, Jia G, et al. Vitamin D status in Chinese pregnant women and their newborns in Beijing and their relationships to birth size. *Public Health Nutr*; 2012:1–6.
- [8] Zhu Z, Zhan J, Shao J, Chen W, Chen L, Li W, et al. High prevalence of vitamin D deficiency among children aged 1 month to 16 years in Hangzhou, China. *BMC Public Health* 2012;12:126.
- [9] Du X, Greenfield H, Fraser DR, Ge K, Trube A, Wang Y. Vitamin D deficiency and associated factors in adolescent girls in Beijing. *Am J Clin Nutr* 2001;74:494–500.
- [10] Foo LH, Zhang Q, Zhu K, Ma G, Hu X, Greenfield H, et al. Low vitamin D status has an adverse influence on bone mass, bone turnover, and muscle strength in Chinese adolescent girls. *J Nutr* 2009;139:1002–7.
- [11] Foo LH, Zhang Q, Zhu K, Ma G, Trube A, Greenfield H, et al. Relationship between vitamin D status, body composition and physical exercise of adolescent girls in Beijing. *Osteoporos Int* 2009;20:417–25.
- [12] Arguelles LM, Langman CB, Ariza AJ, Ali FN, Dilley K, Price H, et al. Heritability and environmental factors affecting vitamin D status in rural Chinese adolescent twins. *J Clin Endocrinol Metab* 2009;94:3273–81.
- [13] Liang GY, Qin R, Li J, Liang GX, Guan YJ, Gao ZH. Optimal level of 25-(OH)D in children in Nanjing (32 degrees N Lat) during winter. *Pediatr Int* 2011;53:541–5.
- [14] Strand MA, Perry J, Zhao J, Fischer PR, Yang J, Li S. Severe vitamin D-deficiency and the health of North China children. *Matern Child Health J* 2009;13:144–50.
- [15] Jiang L, Xu J, Pan S, Xie E, Hu Z, Shen H. High prevalence of hypovitaminosis D among pregnant women in southeast China. *Acta Paediatr* 2012;101:e192–4.
- [16] Tao M, Shao H, Gu J, Zhen Z. Vitamin D status of pregnant women in Shanghai, China. *J Matern Fetal Neonatal Med* 2012;25:237–9.
- [17] Yan L, Prentice A, Zhang H, Wang X, Stirling DM, Golden MM. Vitamin D status and parathyroid hormone concentrations in Chinese women and men from north-east of the People's Republic of China. *Eur J Clin Nutr* 2000;54:68–72.
- [18] Woo J, Lam CW, Leung J, Lau WY, Lau E, Ling X, et al. Very high rates of vitamin D insufficiency in women of child-bearing age living in Beijing and Hong Kong. *Br J Nutr* 2008;99:1330–4.
- [19] Yang B, Sun H, Wan Y, Wang H, Qin W, Yang L, et al. Associations between testosterone, bone mineral density, vitamin D and semen quality in fertile and infertile Chinese men. *Int J Androl* 2012;35:783–92.
- [20] Shao H, Tao M, Fan Y, Jing J, Lu J. Vitamin D levels and other factors related to bone mineral density during pregnancy. *Aust N Z J Obstet Gynaecol* 2012;52:571–5.
- [21] Wang O, Nie M, Hu YY, Zhang K, Li W, Ping F, et al. Association between vitamin D insufficiency and the risk for gestational diabetes mellitus in pregnant Chinese women. *Biomed Environ Sci* 2012;25:399–406.
- [22] Yin X, Sun Q, Zhang X, Lu Y, Sun C, Cui Y, et al. Serum 25(OH)D is inversely associated with metabolic syndrome risk profile among urban middle-aged Chinese population. *Nutr J* 2012;11:68.
- [23] Lin SW, Chen W, Fan JH, Dawsey SM, Taylor PR, Qiao YL, et al. Prospective study of serum 25-hydroxyvitamin d concentration and mortality in a Chinese population. *Am J Epidemiol* 2012;176:1043–50.
- [24] Li L, Yin X, Yao C, Zhu X, Wu X. Vitamin D, parathyroid hormone and their associations with hypertension in a Chinese population. *PLoS One* 2012;7:e43344.

- [25] Ren C, Qiu MZ, Wang DS, Luo HY, Zhang DS, Wang ZQ, et al. Prognostic effects of 25-hydroxyvitamin D levels in gastric cancer. *J Transl Med* 2012;10:16.
- [26] Huang Y, Yu H, Lu J, Guo K, Zhang L, Bao Y, et al. Oral supplementation with cholecalciferol 800 IU ameliorates albuminuria in Chinese type 2 diabetic patients with nephropathy. *PLoS One* 2012;7:e50510.
- [27] Lu HK, Zhang Z, Ke YH, He JW, Fu WZ, Zhang CQ, et al. High prevalence of vitamin D insufficiency in China: relationship with the levels of parathyroid hormone and markers of bone turnover. *PLoS One* 2012;7:e47264.
- [28] MacLaughlin J, Holick MF. Aging decreases the capacity of human skin to produce vitamin D3. *J Clin Invest* 1985;76:1536–8.
- [29] Aksnes L, Rodland O, Aarskog D. Serum levels of vitamin D3 and 25-hydroxyvitamin D3 in elderly and young adults. *Bone Miner* 1988;3:351–7.
- [30] Lu L, Yu Z, Pan A, Hu FB, Franco OH, Li H, et al. Plasma 25-hydroxyvitamin D concentration and metabolic syndrome among middle-aged and elderly Chinese individuals. *Diabetes Care* 2009;32:1278–83.
- [31] Dorjgochoo T, Ou Shu X, Xiang YB, Yang G, Cai Q, Li H, et al. Circulating 25-hydroxyvitamin D levels in relation to blood pressure parameters and hypertension in the Shanghai Women's and Men's Health Studies. *Br J Nutr* 2012;108:449–58.
- [32] Kruger MC, Ha PC, Todd JM, Kuhn-Sherlock B, Schollum LM, Ma J, et al. High-calcium, vitamin D fortified milk is effective in improving bone turnover markers and vitamin D status in healthy postmenopausal Chinese women. *Eur J Clin Nutr* 2012;66:856–61.
- [33] Zhou X, Han J, Song Y, Zhang J, Wang Z. Serum levels of 25-hydroxyvitamin D, oral health and chronic obstructive pulmonary disease. *J Clin Periodontol* 2012;39:350–6.
- [34] Zhang H, Huang QR, Gu JM, Hu WW, Liu YJ, Hu YQ, et al. Comparison of the effects of cholecalciferol and calcitriol on calcium metabolism and bone turnover in Chinese postmenopausal women with vitamin D insufficiency. *Acta Pharmacol Sin* 2012;33:490–5.
- [35] Zhao J, Xia W, Nie M, Zheng X, Wang Q, Wang X, et al. The levels of bone turnover markers in Chinese postmenopausal women: Peking vertebral fracture study. *Menopause* 2011;18:1237–43.
- [36] Yan L, Zhou B, Wang X, D'Ath S, Laidlaw A, Laskey MA, et al. Older people in China and the United Kingdom differ in the relationships among parathyroid hormone, vitamin D, and bone mineral status. *Bone* 2003;33:620–7.
- [37] Tu P, Peng X. China's population: status and challenges. *China Popul Today* 1994;11:13–5.
- [38] van Schoor NM, Lips P. Worldwide vitamin D status. *Best Pract Res Clin Endocrinol Metab* 2011;25:671–80.
- [39] Li LM, Rao KQ, Kong LZ, Yao CH, Xiang HD, Zhai FY, et al. [A description on the Chinese national nutrition and health survey in 2002]. *Zhonghua Liu Xing Bing Xue Za Zhi* 2005;26:478–84.
- [40] Dhingra V, Lau E. The Asian Audit: Epidemiology, costs and burden of osteoporosis in Asia 2009. Switzerland: International Osteoporosis Foundation; 2009.
- [41] Zhu H, Fang J, Luo X, Yu W, Zhao Y, Li X, et al. A survey of bone mineral density of healthy Han adults in China. *Osteoporos Int* 2010;21:765–72.
- [42] Xia WB, He SL, Xu L, Liu AM, Jiang Y, Li M, et al. Rapidly increasing rates of hip fracture in Beijing, China. *J Bone Miner Res* 2012;27:125–9.
- [43] EFSA Panel on Dietetic Products NaAN. Scientific Opinion on the substantiation of a health claim related to vitamin D and risk of falling pursuant to Article 14 of Regulation (EC) No 1924/2006. *EFSA Journal* 2011;9:2382.
- [44] Bischoff-Ferrari HA, Willett WC, Orav EJ, Lips P, Meunier PJ, Lyons RA, et al. A pooled analysis of vitamin D dose requirements for fracture prevention. *N Engl J Med* 2012;367:40–9.