

Higher Serum Levels of Vitamin D Are Associated With a Reduced Risk of Diverticulitis

LILLIAS H. MAGUIRE,* MINGYANG SONG,† LISA E. STRATE,§ EDWARD L. GIOVANNUCCI,‡ and ANDREW T. CHAN||

*Department of Surgery, †Division of Gastroenterology, Massachusetts General Hospital, Boston, Massachusetts; ‡Department of Nutrition, Harvard School of Public Health, Boston, Massachusetts; and §Department of Gastroenterology, University of Washington, Seattle, Washington

BACKGROUND & AIMS: Recent studies have shown geographic and seasonal variations in hospital admissions for diverticulitis. Because this variation parallels differences in ultraviolet light exposure, the most important contributor to vitamin D status, we examined the association of prediagnostic serum levels of vitamin D with diverticulitis.

METHODS: Among patients within the Partners Healthcare System who had blood drawn and serum levels of 25-hydroxyvitamin D (25-[OH]D) measured, from 1993 through 2012, we identified 9116 patients with uncomplicated diverticulosis and 922 patients who developed diverticulitis that required hospitalization. We used multivariate logistic regression to estimate relative risks and 95% confidence intervals to compare serum 25(OH)D levels between these groups.

RESULTS: Patients with uncomplicated diverticulosis had significantly higher mean prediagnostic serum levels of 25(OH)D (29.1 ng/mL) than patients with diverticulitis who required hospitalization (25.3 ng/mL; $P < .0001$). Compared with patients in the lowest quintile of 25(OH)D, the multivariate-adjusted relative risk for diverticulitis hospitalization was 0.49 (95% confidence interval, 0.38–0.62; P for trend $< .0001$) among patients in the highest quintile of 25(OH)D level. Compared with patients with uncomplicated diverticulosis, the mean level of 25(OH)D was significantly lower for patients with acute diverticulitis without other sequelae (25.9 ng/mL; $P < .0001$; $n = 594$), for patients with diverticulitis with abscess (25.8 ng/mL; $P = .0095$; $n = 124$), for patients with diverticulitis requiring emergent laparotomy (22.7 ng/mL; $P = .002$; $n = 65$), and for patients with recurrent diverticulitis (23.5 ng/mL; $P < .0001$; $n = 139$).

CONCLUSIONS: Among patients with diverticulosis, higher prediagnostic levels of 25(OH)D are associated significantly with a lower risk of diverticulitis. These data indicate that vitamin D deficiency could be involved in the pathogenesis of diverticulitis.

Keywords: Vitamin D; Diverticulitis; Diverticulosis; Risk Factor; Epidemiology.

See related articles on pages 1532, 1609, 1614, and 1622 in this issue of *Clinical Gastroenterology and Hepatology*.

Diverticulosis is a nearly ubiquitous colorectal condition of older individuals in the United States and has been reported in more than 70% of colonoscopies in patients older than age 80.¹ Diverticulitis, one of its potential consequences, results in more than 200,000 hospitalizations at a cost of more than 2 billion dollars per year.² Potential complications of diverticulitis, including fistula, abscess, intestinal perforation, stricture, and sepsis, generate substantial morbidity and potential mortality. In recent years, diverticulitis has been increasing in frequency, particularly among younger individuals.³

Long-standing theories regarding the etiology of diverticulitis, such as seed and nut impaction in diverticula,⁴ have been undermined through etiologic studies that also have highlighted novel associations between diverticulitis and lifestyle factors such as obesity⁵ and use of nonsteroidal

anti-inflammatory drugs.⁶ Recently, an analysis of a national database of hospital admissions observed a seasonal variation in hospitalizations for diverticulitis.⁷ Seasonal variation also reflects variation in ultraviolet (UV) light exposure, the greatest contributor to vitamin D status.

Vitamin D is a fat-soluble vitamin dependent largely on exposure of the skin to UV light for its production, with dietary sources playing an important secondary role. The active metabolite, 1,25-dihydroxyvitamin D [1,25(OH)₂D], is synthesized from 25-hydroxyvitamin D [25(OH)D] both in the kidneys and in extrarenal tissues including the colonic epithelium and many cell types in the immune system.⁸ Although its role in

Abbreviations used in this paper: CI, confidence interval; CT, computed tomography; ICD-9, International Classification of Diseases, 9th revision; OR, odds ratio; RPDR, research patient data registry; 25(OH)D, 25-hydroxyvitamin D; UV, ultraviolet.

© 2013 by the AGA Institute

1542-3565/\$36.00

<http://dx.doi.org/10.1016/j.cgh.2013.07.035>

maintaining bone health is well characterized, compelling data support important extraskeletal effects, including an influence on the colon. Previous studies have shown associations between vitamin D and other colonic disease, including colorectal neoplasia^{9,10} and inflammatory bowel disease.¹¹ Based on these data, we hypothesized that a similar link might exist between prediagnostic levels of 25(OH)D and diverticulitis that required hospitalization.

Methods

Assembly of Study Cohort

We identified eligible patients through the Partners Healthcare Research Patient Data Registry (RPDR). Partners Healthcare is an integrated academic health care system that provides care for more than 1.8 million patients in the greater Boston area. The RPDR is a centralized data warehouse for patient information from multiple hospital systems within Partners Healthcare. Access is provided to demographic data, billing codes, problem lists, and narrative notes generated within affiliated hospitals and outpatient facilities. Details regarding the RPDR have been published previously.¹² We queried the RPDR for individuals who had at least one measurement of serum 25(OH)D level from 1993 to 2012 and defined a group of patients with uncomplicated diverticulosis by identifying 9116 patients who had a code for diverticulosis (International Classification of Diseases, 9th revision [ICD-9] code 562.10) and did not have an associated hospitalization for diverticulitis over at least 5 years of follow-up evaluation after diagnosis. We also defined a group of patients hospitalized with diverticulitis as individuals with an inpatient admission associated with the ICD-9 code 562.11 or 562.13 after measurement of 25(OH)D. Within this group, we further examined subtypes of diverticulitis admissions according to the following categories: *uncomplicated diverticulitis*, defined as acute diverticulitis without abscess or perforation (ICD-9 codes 562.11 and 562.13); *complicated diverticulitis*, defined as diverticulitis with abscess or perforation (ICD-9 codes 567, 569.5, and 569.83); *surgical diverticulitis*, defined as diverticulitis requiring laparotomy for peritonitis or pneumoperitoneum (ICD-9 codes 17.3, 45.7, and 54); and *recurrent diverticulitis*, defined as more than

one inpatient admission for diverticulitis with admissions separated by at least 30 days. To assess the validity of our identification of patients with diverticulosis using billing codes and patient problem lists, we examined the medical records of a randomly selected sampling of 100 patients in our control group. We were able to verify the diagnosis of diverticulosis in 95 of 100 patients. Among the 95 patients, the diagnosis was confirmed in 77 (81%) by colonoscopy or sigmoidoscopy, with the remainder confirmed by computed tomography (CT) scan. Of the 5 patients who could not be confirmed, 2 had physician notes documenting diverticulosis without mention of a confirmatory study and 3 did not have documentation in a limited set of clinician notes. Among the 100 cases, there were no cases of diverticulitis. We also examined the medical records of a randomly selected sampling of 100 patients with diverticulitis. We were able to support the diagnosis among 92 patients. Among the 92 patients, 68 (74%) had CT evidence of diverticulitis, with the remaining patients diagnosed by history and physical examination features consistent with diverticulitis and/or increased white blood cell count and fevers.

This study was approved by the Partners Human Research Committee.

Assessment of Exposures and Covariates

For our primary analyses, we examined the first measured 25(OH)D level in relation to risk of diverticulitis. In sensitivity analyses, we also examined the association of mean 25(OH)D levels with diverticulitis among patients with more than one measurement of 25(OH)D and among patients who underwent a CT scan within 7 days of diagnosis. We collected information on demographics (age, sex, race) and selected comorbidities as reported by patient providers in the medical record, including coronary artery disease, obesity, chronic renal failure, and diabetes.

Statistical Analysis

In univariate analyses, we compared the group with uncomplicated diverticulosis with the group hospitalized with diverticulitis using *t* tests for continuous variables and chi-squared tests for categorical variables. We also defined

Table 1. Patient Characteristics

	Group				
	Uncomplicated diverticulosis (N = 9116)	Acute diverticulitis (N = 594)	Complicated diverticulitis (N = 124)	Surgical diverticulitis (N = 65)	Recurrent diverticulitis (N = 139)
Age, mean (SD), y	63.6 (11.6)	67.3 (13.7)	65.8 (11.7)	65.4 (13.0)	66.0 (12.8)
Sex, n (%)					
Male	3515 (38.6)	190 (32.0)	48 (38.7)	19 (29.2)	44 (31.6)
Female	5601 (61.4)	404 (68.0)	76 (61.3)	46 (70.8)	95 (68.4)
Race, n (%)					
White	7775 (85.1)	495 (83.3)	104 (83.9)	57 (87.7)	115 (82.7)
Black	463 (5.1)	40 (6.7)	8 (6.4)	5 (7.7)	8 (5.7)
Other ^a	898 (9.8)	59 (9.9)	12 (9.7)	3 (4.6)	16 (11.5)
Comorbidity, n (%)					
Coronary artery disease	2125 (23.3)	83 (13.9)	11 (8.9)	10 (15.4)	13 (9.3)
Chronic renal failure	1062 (11.6)	61 (10.3)	14 (11.3)	19 (29.2)	9 (6.5)
Diabetes	1497 (16.4)	101 (17.0)	17 (13.7)	14 (21.5)	17 (12.2)
Obesity	85 (9.4)	32 (5.4)	7 (5.6)	3 (4.6)	1 (0.7)

^aOther race includes Hispanic, Asian, Native American, unknown, other, and not documented.

Table 2. First and Average Vitamin D Levels for Each Group

	Group				
	Uncomplicated diverticulosis (N = 9116)	Acute diverticulitis (N = 594)	Complicated diverticulitis (N = 124)	Surgical diverticulitis (N = 65)	Recurrent diverticulitis (N = 139)
First vitamin D level, mean (SD), ng/mL	29.1 (14.0)	25.9 (13.4)	25.8 (12.8)	22.7 (14.9)	23.5 (14.0)
P value ^a		<.0001	.0095	.002	<.0001
Average vitamin D level, mean (SD), ng/mL	33.0 (12.2)	28.1 (12.7)	28.8 (12.7)	23.9 (13.9)	25.5 (13.7)
P value ^a		<.0001	.002	<.0001	<.0001

^aP value compares each group with the uncomplicated diverticulosis group.

quintiles for serum 25(OH)D levels using cut-off points determined according to the distribution among patients with uncomplicated diverticulosis. We used logistic regression to calculate odds ratios (ORs) and 95% confidence intervals (CIs) for the association between quintiles of serum 25(OH)D and risk of hospitalization for diverticulitis. In the multivariate logistic regression, we adjusted for other potential risk factors, including age, sex, race, diabetes, chronic renal failure, obesity, coronary artery disease, and history of a prior colonoscopy. Because diverticulitis was a relatively rare outcome, we interpreted the OR as an estimate of the relative risk. *P* values less than .05 were considered significant. We used SAS version 9.2 (SAS Institute, Inc, Cary, NC) for all analyses.

Results

Table 1 shows the baseline characteristics of our study population. Compared with the 9116 patients with uncomplicated diverticulosis, the 922 patients hospitalized with diverticulitis were older, more likely to be female, and less likely to be obese or have coronary artery disease.

Table 2 compares the prediagnostic serum 25(OH)D level between patients with uncomplicated diverticulosis and those hospitalized with diverticulitis. For individuals with more than one measurement of 25(OH)D, we used the first measured vitamin D level. Among patients with uncomplicated diverticulosis, the mean 25(OH)D was 29.1 ng/mL compared with 25.3 ng/mL among the 922 patients with diverticulitis that required hospitalization (*P* < .0001). Compared with patients with uncomplicated diverticulosis, the mean serum 25(OH)D level was significantly lower for each subtype of diverticulitis: 25.9 ng/mL (*P* < .0001) for acute diverticulitis without other sequelae (*n* = 594), 25.8 ng/mL (*P* = .009) for diverticulitis with abscess (*n* = 124), 22.7 ng/mL (*P* = .002) for those who required emergent laparotomy (*n* = 65), and 23.5 ng/mL (*P* < .0001) for

recurrent diverticulitis (*n* = 139). In sensitivity analyses, for individuals with more than one prediagnostic measurement of 25(OH)D, we used the mean of all 25(OH)D levels for our comparisons and obtained similar results (Table 2). The mean vitamin D level for patients with acute uncomplicated diverticulitis with a documented CT scan within 7 days of diagnosis (*n* = 344) was 24.8 ng/mL compared with 27.4 ng/mL for patients with diverticulitis who did not undergo a CT scan (*n* = 250) as part of their diagnostic evaluation (*P* = .02).

We considered the possibility that our observed associations may be the result of a higher prevalence of healthy behaviors among individuals with asymptomatic diverticulosis compared with those with diverticulitis. Because such health-seeking behavior may be reflected by a higher likelihood of undergoing screening colonoscopy, we performed a secondary analysis limited to individuals who also had a colonoscopy (before admission for case patients with diverticulitis). Among such patients, the association between 25(OH)D and risk of diverticulitis was not significantly different (Table 3).

Table 4 shows the association between quintiles of 25(OH)D level in relation to risk of hospitalization for diverticulitis, adjusting for other risk factors. Compared with patients in the lowest quintile, the age-adjusted ORs for diverticulitis hospitalization were 0.71 (95% CI, 0.59–0.87) among patients in the second quintile of 25(OH)D, 0.56 (95% CI, 0.46–0.69) among patients in the third quintile, 0.56 (95% CI, 0.46–0.69) among patients in the fourth quintile, and 0.45 (95% CI, 0.36–0.56) among patients in the highest quintile (*P* trend < .0001). Additionally, adjusting for available information on comorbidities and health-modifying behaviors such as the presence of coronary artery disease, obesity, chronic renal failure, diabetes, and having undergone a colonoscopy did not materially alter these risk estimates (Table 4).

Compared with patients with acute diverticulitis without other sequelae, patients in the subgroups who developed

Table 3. Vitamin D Levels for Diverticulosis and Diverticulitis Patients With and Without Colonoscopy

	Colonoscopy		No colonoscopy	
	Diverticulosis (N = 7563)	Diverticulitis (N = 486)	Diverticulosis (N = 1553)	Diverticulitis (N = 436)
First vitamin D level, mean (SD), ng/mL	29.7 (13.9)	26.4 (12.9)	26.3 (13.9)	24.1 (14.1)
P value ^a		<.0001		.003
Average vitamin D level, mean (SD), ng/mL	33.7 (10.8)	29.3 (12.2)	29.4 (12.3)	25.8 (13.3)
P value ^a		<.0001		<.0001

^aP value compares the diverticulitis group with the diverticulosis group.

Table 4. Risk of Diverticulitis According to Quintiles of Serum Vitamin D

	Quintile					P trend
	1	2	3	4	5	
First 25-(OH)D						
Quintile median, ng/mL ^a	13.0	21.0	29.0	36.0	47.0	
Age-adjusted RR (95% CI)	1	0.71 (0.59–0.87)	0.56 (0.46–0.69)	0.56 (0.46–0.69)	0.45 (0.36–0.56)	<.0001
Multivariate RR (95% CI) ^b	1	0.71 (0.58–0.87)	0.60 (0.49–0.74)	0.61 (0.49–0.76)	0.49 (0.38–0.62)	<.0001
Average 25-(OH)D						
Quintile median, ng/mL	18.2	27.3	32.8	38.2	47.8	
Age-adjusted RR (95% CI)	1	0.41 (0.34–0.50)	0.39 (0.32–0.48)	0.31 (0.25–0.38)	0.30 (0.25–0.38)	<.0001
Multivariate RR (95% CI) ^a	1	0.44 (0.36–0.54)	0.45 (0.36–0.55)	0.36 (0.28–0.45)	0.34 (0.27–0.43)	<.0001

RR, relative risk.

^aQuintiles are based on the distribution of 25(OH)D level among individuals with uncomplicated diverticulosis.

^bAdjusted for age, sex, race, diabetes, chronic renal failure, obesity, coronary artery disease, and history of a prior colonoscopy.

abscess, required surgery, or had recurrent attacks were observed to have lower prediagnostic levels of vitamin D. These differences did not reach significance, with the exception of surgical patients. Surgical patients had significantly lower average levels of vitamin D as compared with patients with acute diverticulitis without other sequelae (23.9 vs 28.1 ng/mL; $P = .01$).

Conclusions

Diverticulitis is a common disease in developed countries, including the United States. However, factors that lead to diverticulitis among patients with diverticulosis are poorly understood. In this study, we show that prediagnostic serum 25(OH)D level is associated inversely with risk of hospitalization for diverticulitis. Compared with patients in the lowest quintile of serum 25(OH)D level, patients in the highest quintile had a substantially lower risk of developing diverticulitis requiring hospitalization. The lowest levels of 25(OH)D were observed in subgroups with more severe disease. Risk of diverticulitis decreased sharply as 25(OH)D levels increased to 25 to 30 ng/mL, with additional small reductions in risk seen with levels greater than 30 ng/mL.

We have shown an association between measured circulating 25(OH)D level and risk of diverticulitis. Our results extend prior indirect evidence of such an association derived from ecologic studies that have shown that hospitalization for diverticulitis varies according to season of the year as well as geographic location. Both of these factors are associated with variation in exposure to UV-B light, the principal contributor to vitamin D status.

Our results are biologically plausible based on prior data showing the importance of vitamin D in colonic physiology. Previous studies, including those by our group, have shown that low levels of circulating 25(OH)D are associated with an increased incidence of colorectal cancer,^{10,13} as well as lower colorectal cancer-specific survival.¹⁴ At the cellular level, vitamin D has been shown to have a pro-apoptotic and anti-proliferative effect, and expression of synthetic and catabolic enzymes is altered in malignant colonocytes.¹⁵ Vitamin D also appears to play a role in inflammatory bowel disease. We, and others, have shown that patients living at lower latitudes with greater UV-B light exposure have a lower risk for Crohn's disease.¹⁶ Recently, we also showed that a lower predicted plasma level of 25(OH)D is associated with an increase in

the risk of CD.¹¹ In parallel, dietary supplementation with vitamin D in murine models reduces the severity of colitis, modulates the gut immune response, and improves intestinal epithelial barrier function.¹⁷ Taken together, these data support a critical role for vitamin D in maintaining colonic homeostasis by modulating inflammation, maintaining epithelial integrity, and regulating intestinal proliferation.

Our study had several strengths. First, we examined associations between 25(OH)D levels collected before hospitalizations for diverticulitis, minimizing the likelihood of reverse causation (ie, diverticular disease resulting in low vitamin D levels). Second, we used a large database of patients with measured 25(OH)D levels and follow-up evaluation for diverticular outcomes. Third, our findings were robust to a sensitivity analyses, which showed a strong association of both mean plasma 25(OH)D levels taken over time as well as a single assessment of 25(OH)D level in relation to risk.

This study had several limitations. First, we were unable to examine the association of 25(OH)D levels in relation to risk of milder cases of diverticulitis that did not require hospitalization. Thus, we cannot exclude the possibility that our findings are the result of a higher likelihood of patients with low vitamin D levels requiring hospital admission for diverticulitis. Nonetheless, our end point of hospitalizations for diverticulitis is perhaps the most clinically important. Second, our reliance on billing coding likely underestimates the true incidence of uncomplicated diverticulosis. Third, we were limited in our ability to adjust for other potential diverticulitis risk factors. Although we used available information on the presence of comorbidities, such as diabetes mellitus, coronary artery disease, and obesity, to reflect overall patient health, such data are likely incomplete because they required entry and coding by treating clinicians. We also were unable to consider other important lifestyle variables within our data set. One marker of health-promoting behaviors, receipt of colonoscopy, did not appear to alter our results materially. However, we could not consider other potentially important factors, such as diet and physical activity, that were not recorded reliably in the medical record. Physical activity has been associated with both higher 25(OH)D levels and diverticulitis. However, physical activity is unlikely to completely account for the magnitude of our observed inverse associations between 25(OH)D level and diverticulitis. Moreover, the mechanism by which physical activity may be associated with a lower risk of diverticulitis is unclear. It is plausible that such an association could, at least in part, be

mediated by higher 25(OH)D levels related to greater exposure to UV light among more physically active individuals.

In summary, we show that higher prediagnostic serum 25(OH)D levels are associated with a lower risk of requiring hospitalization for diverticulitis. Taken together with prior studies showing an inverse association of 25(OH)D and risk of colorectal cancer and inflammatory bowel disease, these results highlight the potential importance of vitamin D in the maintenance of colonic health. Additional studies in cohorts with more detailed information on potential confounders of this association are warranted.

References

1. Everhart JE, Ruhl CE. Burden of digestive diseases in the United States part II: lower gastrointestinal diseases. *Gastroenterology* 2009;136:741-754.
2. Peery AF, Dellon ES, Lund J, et al. Burden of gastrointestinal disease in the United States: 2012 update. *Gastroenterology* 2012;143:1179-1187.
3. Nguyen GC, Sam J, Anand N. Epidemiological trends and geographic variation in hospital admissions for diverticulitis in the United States. *World J Gastroenterol* 2011;17:1600-1605.
4. Strate LL, Liu YL, Syngal S, et al. Nut, corn, and popcorn consumption and the incidence of diverticular disease. *JAMA* 2008;300:907-914.
5. Strate LL, Liu YL, Aldoori WH, et al. Obesity increases the risks of diverticulitis and diverticular bleeding. *Gastroenterology* 2009;136:115-122.
6. Strate LL, Liu YL, Huang ES, et al. Use of aspirin or nonsteroidal anti-inflammatory drugs increases risk for diverticulitis and diverticular bleeding. *Gastroenterology* 2011;140:1427-1433.
7. Ricciardi R, Roberts PL, Read TE, et al. Cyclical increase in diverticulitis during the summer months. *Arch Surg* 2011;146:319-323.
8. Bises G, Kállay E, Weiland T, et al. 25-hydroxyvitamin D3-1alpha-hydroxylase expression in normal and malignant human colon. *J Histochem Cytochem* 2004;52:985-989.
9. Hong SN, Kim JH, Choe WH, et al. Circulating vitamin D and colorectal adenoma in asymptomatic average-risk individuals who underwent first screening colonoscopy: a case-control study. *Dig Dis Sci* 2012;57:753-756.
10. Jenab M, Bueno-de-Mesquita HB, Ferrari P, et al. Association between pre-diagnostic circulating vitamin D concentration and risk of colorectal cancer in European populations: a nested case-control study. *BMJ* 2010;340:b5500.
11. Ananthakrishnan AN, Khalili H, Higuchi LM, et al. Higher predicted vitamin D status is associated with reduced risk of Crohn's disease. *Gastroenterology* 2012;142:482-489.
12. Nalichowski R, Keogh D, Chueh HC, et al. Calculating the benefits of a Research Patient Data Repository. *AMIA Annu Symp Proc* 2006:1044.
13. Lee JE, Li H, Chan AT, et al. Circulating levels of vitamin D and colon and rectal cancer: the Physicians' Health Study and a meta-analysis of prospective studies. *Cancer Prev Res (Phila)* 2011;4:735-743.
14. Fedirko V, Riboli E, Tjønneland A, et al. Prediagnostic 25-hydroxyvitamin D, VDR and CASR polymorphisms, and survival in patients with colorectal cancer in western European populations. *Cancer Epidemiol Biomarkers Prev* 2012;21:582-593.
15. Cross HS, Nittke T, Kallay E. Colonic vitamin D metabolism: implications for the pathogenesis of inflammatory bowel disease and colorectal cancer. *Mol Cell Endocrinol* 2011;347:70-79.
16. Khalili H, Huang ES, Ananthakrishnan AN, et al. Geographical variation and incidence of inflammatory bowel disease among US women. *Gut* 2012;61:1686-1692.
17. Zhao H, Zhang H, Wu H, et al. Protective role of 1,25(OH)₂ vitamin D₃ in the mucosal injury and epithelial barrier disruption in DSS-induced acute colitis in mice. *BMC Gastroenterol* 2012;12:57.

Reprint requests

Address requests for reprints to: Andrew T. Chan, MD, MPH, Division of Gastroenterology, Massachusetts General Hospital, 55 Fruit Street, GRJ 825C, Boston, Massachusetts 02114. e-mail: achan@partners.org; fax: (617) 724-6832.

Conflicts of interest

These authors disclose the following: Lisa Strate has served as a consultant to Shire Pharmaceuticals, and Andrew Chan has served as a consultant to Bayer Healthcare; Pfizer, Inc; Millennium Pharmaceuticals; and Pozen, Inc. The remaining authors disclose no conflicts.

Funding

Supported by grants from the American College of Gastroenterology and NIH/NIDDK K24 DK098311.