

Received Date : 17-Dec-2013

Revised Date : 24-Jan-2014

Accepted Date : 21-Feb-2014

Article type : Original

Avoidance of sun exposure is a risk factor for all-cause mortality: results from the MISS cohort

Pelle G. Lindqvist¹, Elisabeth Epstein², Mona Landin-Olsson³, Christian Ingvar⁴, Kari Nielsen⁵, Magnus Stenbeck⁶ & Håkan Olsson⁷

From the ¹Department of Obstetrics and Gynecology, Clintec, Karolinska University Hospital, Huddinge, Stockholm, Sweden; ²Department of Obstetrics and Gynecology, Mothers and Childrens Health, Karolinska University Hospital, Solna, Stockholm, Sweden; ³Department of Endocrinology, Clinical Science, Lund University Hospital, Lund, Sweden; ⁴Department of Surgery, Clinical Scienc, Lund, University Hospital, Lund, Sweden; ⁵Department of Dermatology, Helsingborg Hospital, Clinical Sciences, Lund University, Lund, Sweden; ⁶Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden; and ⁷Departments of Oncology and Cancer Epidemiology, Lund University Hospital, Lund, Sweden

Running head: Sun exposure habits and all-cause mortality

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/joim.12251

This article is protected by copyright. All rights reserved.

Correspondence: Pelle G. Lindqvist

Department of Obstetrics and Gynecology, Clintec

Karolinska University Hospital, Huddinge

Kvinnokliniken K 57

SE-14186 Stockholm

Sweden

Tel: +46 708 992545 or +46 40 158910

E-mail: Pelle.lindqvist@ki.se

Abstract

Background. Sunlight exposure and fair skin are major determinants of human vitamin D production, but they are also risk factors for cutaneous malignant melanoma (MM). There is epidemiological evidence that all-cause mortality is related to low vitamin D levels.

Methods. We assessed the avoidance of sun exposure as a risk factor for all-cause mortality for 29,518 Swedish women in a prospective 20-year follow-up of the Melanoma in Southern Sweden (MISS) cohort. Women were recruited from 1990 to 1992, and were aged 25 to 64 years at the start of the study. We obtained detailed information at baseline on their sun exposure habits and potential confounders. Multivariable flexible parametric survival analysis was applied to the data.

Results. There were 2545 deaths among the 29,518 women who responded to the initial questionnaire. We found that all-cause mortality was inversely related to sun exposure habits. The mortality rate among avoiders of sun exposure was approximately two-fold higher compared to the highest sun exposure group, resulting in excess mortality with a population attributable risk of 3%.

Conclusion. The results of this study provide observational evidence that avoiding sun exposure is a risk factor for all-cause mortality. Following sun exposure advice that is very restrictive in countries with low solar intensity might in fact be harmful to women's health.

Keywords: evolution, longevity, melanoma, population attributable risk, UV radiation, vitamin D.

Introduction

Ultraviolet (UV) radiation from the sun is known to heighten the risk of developing malignant melanoma (MM) of the skin. This condition is primarily responsible for increased mortality due to UV radiation exposure. The risk of MM varies widely among people of different skin colour, depending on the type of melanin in their skin. MM is most common among Northern Europeans with pale skin and is rare among Africans with very dark skin [1]. Individuals with red hair or a tendency to develop freckles are at increased risk of developing MM. The highest risk has been found among those of European ancestry living in Northern Australia [1]. This has been the basis for considering UV radiation as the major cause of MM. Programmes for avoiding UV radiation are widely implemented in societies in which a large proportion of the population is descended from Europeans [2].

Nevertheless, exposure to sunlight remains the main source of vitamin D. Sunlight UVB radiation with a wavelength between 290 and 315 nm penetrates the skin and converts 7-dehydrocholesterol to 25-hydroxycholecalciferol vitamin D₃ via previtamin D [3]. Low vitamin D levels have been associated with an increase in both all-cause and cardiovascular disease mortality, but results are inconsistent [4-8]. Low vitamin D concentrations have also

Accepted Article
been linked to thicker, more aggressive melanomas with shorter survival times [9]. Thus, opposing mechanisms may act with regard to MM and sunlight exposure.

Our longitudinal cohort study was carried out in order to assess how all-cause mortality risk may be influenced by avoidance of sun exposure in women.

Methods

Study design

The study was approved by the Ethics Committee of Lund University (LU 632-03). In 1990 the Melanoma in Southern Sweden (MISS) study was initiated. Our subjects consisted of approximately 1000 Sweden-born women of each age from 25 to 64 years ($n = 39,973$) with no history of malignancy. The subjects were selected from the general population registry of the South Swedish Health Care Region by random computerized selection, and represented 20% of the female population of South Sweden in the selected age groups.

The women were invited to complete a standardized written questionnaire concerning risk factors for MM. The initial inquiry was carried out from 1990 to 1992 and 29,518 women subsequently entered the study (response rate 74%). A written follow-up was then conducted between 2000 and 2002. A total of 184 women emigrated during the study period and were censored after emigration. The questionnaire was a detailed inquiry into several factors of potential interest for longevity, such as sun exposure habits, marital status, educational level, smoking habits, alcohol consumption, number of pregnancies, body mass index (BMI) and physical exercise. As sun exposure may depend on economic resources, we collected information on mean personalized family income between 1990 and 1993, i.e. approximately

the time of study initiation, from official income and taxation records at Statistics Sweden (for further details see <http://www.scb.se/en/>). Four questions were asked regarding sun exposure: (i) How often do you sunbathe during the summertime? (never, 1–14 times, 15–30 times, >30 times); (ii) Do you sunbathe during the winter, such as on holiday to the mountains? (no; 1–3 days, 4–10 days; >10 days); (iii) Do you use tanning beds? (never; 1–3 times/year; 4–10 times/year; >10 times/year); and (iv) Do you go abroad on holiday to swim and sunbathe? (never; once/1–2 years; once/year; two or more times/year). The four questions were dichotomized into yes/no in the analysis (i.e. 'no/never' or 'sometimes'). As a measure of total sun exposure, we created a four-score variable depending on the number of 'yes' responses to the above questions from 0 (avoiding sun exposure: reference) to 4 (highest sun exposure). For the purpose of presentation in the figures and in order to estimate population attributable risk (PAR), sun exposure habits were categorized into three groups: zero 'yes' responses (avoidance of sun exposure); 'yes' response to one or two questions (moderate exposure); and 'yes' response to three or four questions (highest exposure). Hereditary disposition towards MM was defined as having a first-degree relative with MM. Red hair was considered a risk factor for developing MM. Known prognostic factors for MM include gender, *in situ*/invasive MM, Breslow thickness, location of MM, ulceration and age at diagnosis. In the present study we included only women, and Breslow thickness was dichotomized into >0.8 mm or <0.8 mm. In the analysis of case fatality for MM, characteristics of MM were dichotomized into invasive MM and Breslow thickness >0.8 mm versus other, due to an empty group. As the hazard ratio (HR) for the extremities and head/neck area were similar, location was dichotomized into trunk or other. We did not have access to information on whether or not ulceration was present.

Cases of MM were identified up to 1 January 2011 by cross-linking with the National Cancer Register and Cause of Death Register. Vital statistics were determined from the National Population Register and the Cause of Death Register also up to 1 January 2011.

Smoking habits at baseline were recorded and categorized into lifetime number of cigarettes smoked: none (reference), <100 000 and >100 000. This was based on self-reported estimations of mean cigarette consumption over 5-year intervals.

As a measure of comorbid illness at the start of the study, we created a dummy variable 'comorbidity' identifying women who have been treated with antidiabetic [Anatomical Therapeutic Chemical Classification System (ATC) A:10] or anticoagulant (ATC B:01) drugs or medications for cardiovascular disease (ATC C:01–C:10) for more than 1 month.

Weight and height were recorded at the second questionnaire in the year 2000, and BMI was calculated (in kg/m²). BMI was classified into three groups: <25 (reference), 25 to <30 (overweight) and >30 kg/m² (obese). The level of regular exercise was categorized at the second interview into three groups: none; walking at least once per week; or strenuous exercise [10].

Statistical analysis

Descriptive statistical analysis was performed using cross-tabulation with the 95% confidence interval (CI) (Table 1). Because we were interested in evaluating both differences in mortality rate with attained age as timescale and relative risk, we chose flexible parametric models for survival analysis [11]. All-cause mortality was used as a dependent variable and time variable as time from inception, and at attained age in Figs 1 and 2, respectively. Time from inception

was calculated from inclusion (see above definition) to death, emigration or 1 January 2011, whichever occurred first. As shown in Table 2, model 2 included BMI and physical exercise from the second interview; data before year 2000 were censored. Missing values were represented by a dummy variable and included in the analysis. The model was fitted with a combination of backward and forward selection. Red hair, heredity for MM, parity and alcohol consumption were not found to be significant in multivariable analysis of all-cause mortality and were not included in the final model (Table 2). Red hair and hereditary disposition towards MM are known risk factors for MM and were included as adjustments in model 1, as shown in Table 3. In model 2 (see Table 3), comorbidity and earlier known variables such as; location, Breslow thickness, *in situ*/invasive MM and age at diagnosis, were included. As Breslow thickness and invasiveness are correlated, a new dummy variable ‘MM characteristics’ was created and categorized into two groups: *in situ* MM/invasive MM and Breslow <0.8 mm or invasive MM and Breslow >0.8mm. A logistic regression analysis was performed (with the same adjustments as in the models shown in Table 2) in order to estimate PAR. To compute PAR, the following formula was used: $PAR = P_e \times [adjusted\ odds\ ratio\ (OR) - 1] / adjusted\ OR$ (Table 2, model 2), where P_e is the proportion of cases assigned to a given variable [12].

IBM SPSS 21 (Statistical Package for the Social Sciences, SPSS Inc., Chicago IL, USA) software was used for descriptive analysis and logistic regression analysis, and STATA 12 (Statacorp, College Station TX, USA) was used for the multivariable flexible parametric regression model. *P* values < 0.05 were considered statistically significant.

Results

There were 2545 deaths among the 29,518 women – representing 540,577 person-years – who responded to the initial questionnaire. As can be seen in Table 1, all the potential

Accepted Article

confounders were significantly related to sun exposure habits and were initially included in the model. In model 1 (see Table 2), we show how sun exposure relates to all-cause mortality. Most comparisons showed significantly lower HR values among those with the highest sun exposure habits, as compared to those who avoided sunshine. In model 2 (see Table 2), BMI and physical exercise were included and the data were censored before the year 2000. HR values were similar and the CIs were wider. In both models the summary sun exposure variables showed a ‘dose-dependent’ inverse relation between sun exposure and all-cause death.

In Fig. 1 all-cause mortality was presented as an adjusted flexible parametric regression survival model with three sun exposure groups. Adjustments were made for comorbidity, age, smoking habits, education level, marital status and disposable income. Fig. 2 shows all-cause mortality rate by attained age, in the three sun exposure groups. As compared to the highest sun exposure group, the mortality rate was doubled (2.0, 95% CI 1.6 – 2.5) among avoiders of sun exposure and increased by 40% (1.4, 95% CI 1.1– 1.7) in those with moderate exposure. We found that the assumption of proportional hazards seemed reasonable. We estimated that PAR for mortality for those avoiding sun exposure was 3%.

Table 3 shows the risk of developing MM by sun exposure habits (as both number of women and HR values). In addition the overall prognosis and MM case fatality-related deaths among the 267 women who developed MM during the study is presented (Table 3, models 2 and 3). Adjustments were made for known risk factors including age at diagnosis, MM characteristics, location (trunk or other) and comorbidities.

Discussion

We found that all-cause mortality was inversely related to sun exposure habits in a ‘dose-dependent’ manner. The mortality rate was increased two-fold among avoiders of sun exposure as compared to those with the highest sun exposure habits. In this study focusing on avoidance of sun exposure, women with ‘normal’ sun exposure habits were not at significantly increased risk of MM or of MM-related death.

In the present study we did not focus on the cause of death. However, the effect was presumably attributed to cancer, heart disease and cerebrovascular disease. There is a known seasonal variation in both heart disease and cerebrovascular disease, with increased risk during winter/spring as compared to summer [13, 14]. There is some epidemiological evidence of an increased risk of cardiovascular disease and early death among individuals with low levels of vitamin D, however these results are inconsistent [5, 6, 15-18]. Only women without a history of diagnosed cancer were included in the present study. Thus, the comparison between those with and without cancer is biased in our study. Previously, it has been suggested that season-related differences in diseases are mainly due to cold or humidity, but we recently proposed that lack of sun exposure might be the environmental factor causing the increased risk of incident thromboembolism, type 2 diabetes and eclampsia in winter [10, 19, 20]. We speculated that insufficient vitamin D levels among those who avoid sun exposure might be the mechanism that increases the mortality rate. However, our findings are of association and not necessarily causal. Regarding diabetes, for example, the melatonin system might be involved in the lower risk of type 2 diabetes among those with the highest sun exposure habits [10, 21]. Another possible explanation is that UVB exposure induces cutaneous endorphins. It has been reported that the use of tanning beds induces mood enhancement, feelings of relaxation and socialization, due to UVB exposure [22, 23]. The

presence of an awarding system at UVB exposure may also be interpreted as a mechanism for avoiding vitamin D deficiency. Another possibility is that solar UVA radiation causes an increase in skin-derived NO bioactivity lowering blood pressure and cardiovascular morbidity [24].

Skin tumors, the major concern of excessive sun exposure, consist of three types (with increasing severity): basal cell carcinoma, squamous cell carcinoma and MM. Sun exposure increases the risk of MM mainly through episodic sunburn or frequent use of tanning beds in individuals with unprotected fair skin. The incidence of MM represents the greatest increase of all cancers affecting women at lower age than men. The disease is fatal in approximately 20% of patients with MM. Basal cell carcinoma and squamous cell carcinoma often affect older individuals and are mostly associated with cumulative exposure to UV light. The incidences of both these types of tumour are increasing. Basal cell carcinoma is generally not fatal but could be locally aggressive, while surgery or radiotherapy provides a cure for most patients with squamous cell carcinoma [25].

US Navy personnel have been reported to be at high risk of skin cancer, but at a reduced risk of all-cause mortality (−26%) and other forms of cancer [26, 27]. However, by using the contraction of basal cell carcinoma as a proxy for heavy sun exposure, without taking into consideration differences in *p53* polymorphisms, the finding of a lower risk of internal cancer has been questioned in a large well-executed case–control study [28]. Support for our results comes from Yang and co-workers, who reported a 30% lower all-cause mortality in individuals who took sunbathing holidays at least once a year over the course of three decades [29]. These authors reported no difference in all-cause mortality among those who did not use tanning beds, as compared to those who did so <12 times per year. We focused on

the low sun exposure group as a risk factor for mortality, whereas most other studies focused on the disadvantages of high sun exposure (so-called overuse).

In agreement with our results, Newton-Bishop and co-workers reported thinner, less aggressive melanoma with greater survival rates in the presence of higher vitamin D levels [9]. Our findings are also consistent with the results of a study by Måsbäck and co-workers; these authors concluded that sun-associated MM was related to superficial spreading MM, in contrast to nodular MM, which had a worse prognosis [30]. Because non-MM skin cancer risk increases with cumulative sun exposure, the finding of an increased incidence of this type of skin cancer with increasing vitamin D levels does not contradict our results [31-33].

Public health implications

Over the last 30 to 40 years, Sweden and other western countries have produced national guidelines provided restrictive advice on sun exposure habits in order to lower the risk of skin cancer. These recommendations are based on Australian/English guidelines known as SunSmart [2], which may be appropriate in a country with a high UV index such as Northern Australia, where dense pigmentation would be an advantage. However, following generally restrictive guidelines in Sweden, a country located at the northern latitude of between 55° and 67° with limited sunshine and a low UV index, might not be optimal. In fact, our findings indicate that these guidelines may indeed be harmful in terms of overall health of the population. The Southern Australia (Tasmania) Cancer Council recommendations including daily short sun exposure might be more suitable in regions with low sun intensity levels [34]. Moreover, in areas with less intense sunlight, genetic variations appear to play a role as a risk factor for MM and recent evidence shows that non-pigmentation-associated polymorphisms may be of major importance in risk of developing MM [35-37].

Our study population of women born in Sweden before 1966, i.e. before widespread immigration took place, consisted almost entirely of light-skinned Caucasian women. If avoidance of sun exposure is a major risk factor for all-cause mortality in the case of Caucasian women, the problem may even be more serious among women who traditionally cover their skin or women more densely pigmented. In the USA, black women were reported to have a 26% excess all-cause mortality, as compared to Caucasian women [38]. Such data should direct our attention to a risk group whose health we may be in a position to improve.

Strengths and weaknesses

A strength of the present study is its use of an unselected large population-based cohort drawn from the Swedish National Population Register and followed for 20 years. The information on sun exposure habits was gathered at the start of the study, as a result of which there is no recall bias. The assumption was made that sun exposure habits did not change over time, and consequently information from one assessment alone was used in the models. This is a common assumption in cohort studies and tends to underestimate risk. Although a causal relationship has not been established, it cannot be ruled out. The inverse dose–response association between sun exposure and all-cause mortality increases the likelihood of a causal relationship (Table 2 and Fig. 1). However, additional evidence and possibly other study designs are needed to draw causal conclusions.

A limitation of this study is the lack of information on physical exercise habits and BMI from baseline, although such information was included from the second interview. Further, we included a variable of comorbidity to adjust for differences in health at baseline. In addition, similar results were found after censoring the first 10 years (Table 2, model 2), further minimizing any differences that might have occurred due to illness at the start of the study.

We have no data on vitamin D supplementation or vitamin D levels, but it is well known that

Accepted Article

sun exposure is the most important determinant of vitamin D status. Despite having adjusted for several potential confounders, the major shortcoming of this study is the inability to distinguish between the consequences of an unhealthy lifestyle and of avoidance of sun exposure.

Evolutionary perspective

From an evolutionary perspective, there must be a selection advantage in being less pigmented when living far from the equator; otherwise, differences in skin pigmentation would not have emerged. Fair skin might confer an evolutionary advantage in regions with low solar intensity by preventing vitamin D deficiency and thereby increasing longevity and reducing infections [39, 40]. However, a fair-skinned individual, for example emigrating from the UK, would not derive any benefit from light skin in Australia, where pigmented skin is an advantage due to high solar intensity.

Conclusion

We conclude that women who avoid sun exposure are at an increased risk of all-cause death with a two-fold increased mortality rate as compared to those with the highest sun exposures. The implementation of restrictive sun exposure advice in countries with low solar intensity might not be beneficial to women's health.

Author contributions

HO was the initiator of the MISS cohort. PGL and HO were responsible for the study design. PGL performed data analysis, with input from all authors, and wrote the initial draft. All authors contributed to the literature search, data interpretation and writing the paper. Additionally, all authors have approved the final version of this paper.

This article is protected by copyright. All rights reserved.

Conflict of interest statement

None of the authors has any conflicts of interest to declare.

Acknowledgements

This study was supported by funds from Karolinska Institutet, the Faculty of Medicine, Lund University, Region Skåne (ALF), the Swedish Cancer Society and the Swedish Medical Research Council. Funding was also received from Lund University Hospital, Region Skåne, the Gustav V Jubilee Fund, the Gunnar Nilsson Foundation, the Kamprad Foundation and the European Research Council Advanced Grant ERC-2011-294576.

Financial disclosure

The funding bodies had no role in the study design, data collection, analysis and interpretation, decision to publish or preparation of the manuscript.

References

1. Buettner PG, Raasch BA. Incidence rates of skin cancer in Townsville, Australia. *Int J Cancer* 1998; **78**: 587-93.
2. CR-UK. SunSmart programme. 2006 [cited 2011; Available from: www.cancerresearchuk.org/sunsmart/]
3. Holick MF. Vitamin D deficiency. *N Engl J Med* 2007; **357**: 266-81.
4. Dobnig H, Pilz S, Scharnagl H, et al. Independent association of low serum 25-hydroxyvitamin d and 1,25-dihydroxyvitamin d levels with all-cause and cardiovascular mortality. *Arch Intern Med* 2008; **168**: 1340-9.
5. McGreevy C, Williams D. New insights about vitamin D and cardiovascular disease: a narrative review. *Ann Intern Med* 2011; **155**: 820-6.
6. Eaton CB, Young A, Allison MA, et al. Prospective association of vitamin D concentrations with mortality in postmenopausal women: results from the Women's Health Initiative (WHI). *Am J Clin Nutr* 2011; **94**: 1471-8.
7. Schottker B, Ball D, Gellert C, et al. Serum 25-hydroxyvitamin D levels and overall mortality. A systematic review and meta-analysis of prospective cohort studies. *Ageing research reviews* 2013; **12**: 708-18.
8. Zittermann A, Iodice S, Pilz S, et al. Vitamin D deficiency and mortality risk in the general population: a meta-analysis of prospective cohort studies. *Am J Clin Nutr* 2012; **95**: 91-100.

9. Newton-Bishop JA, Beswick S, Randerson-Moor J, et al. Serum 25-hydroxyvitamin D3 levels are associated with breslow thickness at presentation and survival from melanoma. *J Clin Oncol* 2009; **27**: 5439-44.
10. Lindqvist PG, Olsson H, Landin-Olsson M. Are active sun exposure habits related to lowering risk of type 2 diabetes mellitus in women, a prospective cohort study? *Diabetes Res Clin Pract* 2010; **90**: 109-14.
11. Lambert PC, Royston P. Further development of flexible parametric models for survival analysis. *The Stata Journal* 2009; **9**: 265-90.
12. Miettinen OS. Proportion of disease caused or prevented by a given exposure, trait or intervention. *Am J Epidemiol* 1974; **99**: 325-32.
13. Crawford VL, McCann M, Stout RW. Changes in seasonal deaths from myocardial infarction. *QJM* 2003; **96**: 45-52.
14. Turin TC, Kita Y, Rumana N, et al. Stroke case fatality shows seasonal variation regardless of risk factor status in a Japanese population: 15-year results from the Takashima Stroke Registry. *Neuroepidemiology* 2009; **32**: 53-60.
15. Wang TJ, Pencina MJ, Booth SL, et al. Vitamin D deficiency and risk of cardiovascular disease. *Circulation* 2008; **117**: 503-11.
16. Brondum-Jacobsen P, Benn M, Jensen GB, et al. 25-Hydroxyvitamin D Levels and Risk of Ischemic Heart Disease, Myocardial Infarction, and Early Death: Population-Based Study and Meta-Analyses of 18 and 17 Studies. *Arterioscler Thromb Vasc Biol* 2012; **32**: 2794-802.
17. Pilz S, Gaksch M, O'Hartaigh B, et al. The role of vitamin D deficiency in cardiovascular disease: where do we stand in 2013? *Archives of toxicology* 2013; **87**: 2083-103.
18. Wang L, Song Y, Manson JE, et al. Circulating 25-hydroxy-vitamin D and risk of cardiovascular disease: a meta-analysis of prospective studies. *Circulation Cardiovascular quality and outcomes* 2012; **5**: 819-29.
19. Lindqvist PG, Epstein E, Olsson H. Does an active sun exposure habit lower the risk of venous thrombotic events? A D-lightful hypothesis. *J Thromb Haemost* 2009; **7**: 605-10.
20. Rylander A, Lindqvist PG. Eclampsia is more prevalent during the winter season in Sweden. *Acta Obstet Gynecol Scand* 2011; **90**: 114-7.
21. Lyssenko V, Nagorny CL, Erdos MR, et al. Common variant in MTNR1B associated with increased risk of type 2 diabetes and impaired early insulin secretion. *Nat Genet* 2009; **41**: 82-8.
22. Poorsattar SP, Hornung RL. UV light abuse and high-risk tanning behavior among undergraduate college students. *J Am Acad Dermatol* 2007; **56**: 375-9.
23. Kaur M, Liguori A, Lang W, et al. Induction of withdrawal-like symptoms in a small randomized, controlled trial of opioid blockade in frequent tanners. *J Am Acad Dermatol* 2006; **54**: 709-11.
24. Liu D, Fernandez BO, Hamilton A, et al. UVA Irradiation of Human Skin Vasodilates Arterial Vasculature and Lowers Blood Pressure Independently of Nitric Oxide Synthase. *J Invest Dermatol* 2014.
25. Reszko A, Aasi SZ, Wilson LD, et al. Cancer of the Skin. In: DeVita VT, Lawrence TS, Rosenberg SA, DePinho R, A,, Weinberg R A, editors. *Cancer Principles & Practice of Oncology*. Philadelphia: LIPPINCOTT WILLIAMS & WILKINS, a WOLTERS KLUWER business; 2011.
26. Peller S, Stephenson CS. Skin irritation and Cancer in the United States Navy. *Am J M Sc* 1937; **194**: 326-33.
27. Groves FD, Page WF, Gridley G, et al. Cancer in Korean war navy technicians: mortality survey after 40 years. *Am J Epidemiol* 2002; **155**: 810-8.
28. Lindelöf B, Krynitz B, Ayoubi S, et al. Previous extensive sun exposure and subsequent vitamin D production in patients with basal cell carcinoma of the skin, has no protective effect on internal cancers. *Eur J Cancer* 2012; **48**: 1154-8.

29. Yang L, Lof M, Veierod MB, et al. Ultraviolet exposure and mortality among women in Sweden. *Cancer Epidemiol Biomarkers Prev* 2011; **20**: 683-90.
30. Masback A, Westerdahl J, Ingvar C, et al. Clinical and histopathological characteristics in relation to aetiological risk factors in cutaneous melanoma: a population-based study. *Melanoma Res* 1999; **9**: 189-97.
31. Srikanth V, Fryer J, Venn A, et al. The association between non-melanoma skin cancer and osteoporotic fractures--a population-based record linkage study. *Osteoporos Int* 2007; **18**: 687-92.
32. Asgari MM, Tang J, Warton ME, et al. Association of prediagnostic serum vitamin D levels with the development of basal cell carcinoma. *J Invest Dermatol*; **130**: 1438-43.
33. Eide MJ, Johnson DA, Jacobsen GR, et al. Vitamin D and Nonmelanoma Skin Cancer in a Health Maintenance Organization Cohort. *Arch Dermatol* 2011; **147**: 1379-84.
34. Tasmania CC. Vitamin D in Tasmania - getting the right UV balance. Tasmania: Cancer Council Tasmania; 2012.
35. Bertolotto C, Lesueur F, Giuliano S, et al. A SUMOylation-defective MITF germline mutation predisposes to melanoma and renal carcinoma. *Nature* 2011; **480**: 94-8.
36. Barrett JH, Iles MM, Harland M, et al. Genome-wide association study identifies three new melanoma susceptibility loci. *Nat Genet* 2011; **43**: 1108-13.
37. Bishop DT, Demenais F, Iles MM, et al. Genome-wide association study identifies three loci associated with melanoma risk. *Nat Genet* 2009; **41**: 920-5.
38. Grant WB, Peiris AN. Possible role of serum 25-hydroxyvitamin D in black-white health disparities in the United States. *J Am Med Dir Assoc*; **11**: 617-28.
39. Jablonski NG, Chaplin G. The evolution of human skin coloration. *J Hum Evol* 2000; **39**: 57-106.
40. Bergman P, Norlin AC, Hansen S, et al. Vitamin D3 supplementation in patients with frequent respiratory tract infections: a randomised and double-blind intervention study. *BMJ open* 2012; **2**.

Fig. 1 Adjusted all-cause survival plot of 29,518 women recruited in the Melanoma in Southern Sweden (MISS) cohort (1990 to 1992) and followed for sun exposure habits for 20 years. Sun exposure habits were classified by the number of 'yes' answers to the following questions: (i) Do you sunbathe during the summer? (ii) Do you sunbathe during the winter, such as during holidays to the mountains? (iii) Do you use tanning beds? (iv) Do you go abroad on holiday to swim and sunbathe? Significance of difference: $P < 0.001$ among all three sun exposure groups. Adjustments were made for comorbidity, age, smoking habits, education level, marital status and disposable income.

Fig. 2 Mortality rate by sun exposure habits with attained age as the time variable for all 29,518 women included in the study. As compared to the highest sun exposure group, the mortality rate was two-fold higher (2.0, 95% confidence interval (CI) 1.6–2.5) among avoiders of sun exposure and increased by 40% (1.4, 95% CI 1.1–1.7) in those with moderate exposure.

Table 1 Demographic characteristics of women with active and not active sun exposure

habits at inception of study					
	Not active		Active		Significance of difference (<i>P</i>)
	sun exposure habits		sun exposure habits**		
	(<i>n</i> = 1721)	%	(<i>n</i> = 27,797)	%	
<u>Womens' characteristics and habits*</u>					
<u>Education</u>					<0.001
≤9 years	741	43.1	4,842	17.4	
9 years	143	8.3	2,630	9.5	
10–12 years	219	12.7	7,440	26.8	
≥12 years	344	20.0	9,353	33.6	
Other	274	15.9	3,532	12.7	
<u>Marital status</u>					<0.001
Unmarried	149	8.7	2,414	8.7	
Married	1,239	72.0	21,771	78.3	
Divorced	140	8.1	2,527	9.1	
Widow	149	8.7	1004	3.6	
Unknown*	44	2.6	81	0.3	
<u>Parity</u>					<0.001
0	325	18.9	4,647	16.7	
1 to 2	820	47.6	15,637	56.3	
≥3	576	33.5	7,513	27.0	
<u>Smoking</u>					<0.001
Non-smokers	947	55.0	11,370	40.9	
<100 000 ciggarettes	355	20.6	9298	33.4	
≥100,000 cigarettes	334	19.4	6490	23.3	
Unknown*	85	4.9	639	2.3	

<u>Alcohol consumption</u>					<0.001
None or <5 g/day	1,280	74.4	17,171	61.8	
5 to <10 g/day	73	4.2	4822	17.3	
10 to <15 g/day	32	1.9	1788	6.4	
≥15 g/day	27	1.6	1261	4.5	
Unknown*	309	18.0	2,755	9.9	
<u>Red Hair</u>					0.001
Yes	92	5.3	1,028	3.8	
<u>Heredity risk of melanoma</u>					0.01
Yes	86	5.0	1,046	3.8	
<u>Disposable income</u>					< 0.001
Low	787	45.7	5,661	20.4	
Moderate	459	26.7	8,454	30.4	
High	475	27.6	13,682	49.2	
<u>Comorbidity***</u>					
Yes	1,351	20.4	2,571	9.2	<0.001
<u>Data from second interview</u>		<i>n</i> = 1210		<i>n</i> = 22,888	
<u>BMI</u>					<0.001
<25 kg/m ²	421	34.8	12,662	55.3	
25 – <30 kg/m ²	346	28.6	6434	28.1	
≥30 kg/m ²	237	19.6	2242	9.8	
Unknown	206	17.0	1550	6.8	
<u>Physical exercise</u>					<0.001
None	167	13.8	1892	8.3	
Light	484	40.0	10,322	45.1	
Strenuous	224	18.5	7476	32.7	
Unknown	335	27.7	3198	14.0	

*Some women did not answer all questions.

**Answering 'yes' to at least one of the sun exposure questions.

***Women who have been treated for >1 month with drugs with the ATC codes A10, B01 or C01–10.

Table 2 Sun exposure habits and risk of all-cause mortality: multivariate analysis

	Women alive	Women dead	Model 1		Model 2	
			HR	95% CI	HR	95% CI
<u>Use of sunbeds?*</u>						
No	11,117	1,825	1.0	ref	1.0	ref
Yes	12,856	720	0.77	0.7–0.8	0.87	0.8–0.98
<u>Sunbathing during winter holiday?*</u>						
No	21,777	2,288	1.0	ref	1.0	ref
Yes	5,196	257	0.81	0.7–0.9	0.85	0.7–1.02
<u>Sunbathing during summer?*</u>						
No	1,719	425	1.0	ref	1.0	ref
Yes	25,254	2,120	0.74	0.7–0.8	0.80	0.7–0.9
<u>Sunbathing during holiday abroad?*</u>						
No	11,331	1,323	1.0	ref	1.0	ref
Yes	15,642	1,222	0.83	0.8–0.9	0.88	0.8–0.97
<u>Summary sun exposure (number of 'yes' answers to above four questions)</u>						
0	1,352	364	1.0	ref	1.0	ref
1	6,229	771	0.80	0.7–0.9	0.83	0.7–1.0
2	8,384	782	0.71	0.6–0.8	0.74	0.6–0.9
3	8,081	508	0.61	0.5–0.7	0.66	0.5–0.8

4	2,927	115	0.53	0.4–0.7	0.62	0.5–0.8
0	1,352	369	1.0	ref	1.0	ref
1 to 2	14,613	1553	0.76	0.7–0.8	0.79	0.7–0.9
3 to 4	11,008	623	0.60	0.5–0.7	0.66	0.5–0.8

Cox regression analysis: *not all women answered all questions.

Model 1 adjusted for age, smoking, marital status, educational level, disposable income and comorbidity.

Model 2 adjusted for all the above and BMI and physical exercise among those answering the second questionnaire i.e. censoring outcomes before second interview (in year 2000).

Table 3 Relation between sun exposure habits and risk of malignant melanoma (MM), all-cause and MM fatality

	1. Women with MM			2. Overall case fatality among women with MM			3. MM case fatality among women with MM		
	n	HR	95% CI	n	HR	95% CI	n	HR	95% CI
	<u>Use of sunbeds?</u>								
No	143	1.0	ref	24	1.0	ref	15	1.0	ref
Yes	124	1.2	1.0–1.6	15	1.0	0.5–2.0	9	1.2	0.5–2.9
<u>Sunbathing during winter holiday?</u>									
No	206	1.0	ref	32	1.0	ref	18	1.0	ref
Yes	61	0.9	0.7–1.3	7	1.2	0.5–2.6	6	1.6	0.6–4.1
<u>Sunbathing during summer?*</u>									
No	20	1.0	ref	7	1.0	ref	3	1.0	ref
Yes	247	1.1	0.7–1.8	32	0.6	0.2–1.3	21	0.9	0.3–3.2

Sunbathing during holiday abroad?

No	89	1.0	ref	15	1.0	ref	8	1.0	ref
Yes	178	1.0	0.8–1.3	24	1.3	0.6–2.5	16	2.0	0.8–4.8

Summary sun exposure adjusted*

0	14	1.0	ref	5	1.0	ref	2	1.0	ref
1	51	1.3	0.7–2.5	8	0.5	0.2–1.7	4	0.7	0.1–3.7
2	76	1.1	0.6–2.1	12	0.8	0.3–2.4	4	1.2	0.2–6.0
3	97	1.2	0.6–2.2	10	0.7	0.2–2.0	3	1.2	0.2–5.8
4	29	1.5	0.7–3.0	4	0.8	0.2–2.9	3	1.9	0.3–3.5

Summary sun exposure adjusted*

0	14	1.0	ref	5	1.0	ref	2	1.0	ref
1 to 2	127	1.18	0.6–2.2	20	0.6	0.2–1.8	12	0.9	0.2–4.3
3 to 4	126	1.24	0.7–2.3	14	0.44	0.08–2.5	10	1.3	0.3–6.2

Active sun exposure*

No	14	1.0	ref	5	1.0	ref	2	1.0	ref
Yes 1 to 4	250	1.2	0.7–2.2	34	0.7	0.2–1.7	22	1.1	0.3–4.8

*Summary sun exposure = number of 'yes' answers to above four questions.

Model 1: incidence of MM adjusted for age at diagnosis, red hair, hereditary disposition towards MM and smoking habits for the entire study population.

Model 2: all-cause mortality among women with MM; model 3: MM case fatality adjusted for age at diagnosis, MM characteristics (*in situ*/invasive and Breslow thickness <0.8 mm /invasive and Breslow thickness \geq 0.8 mm), location (trunk or other) and comorbidity (being on antidiabetic, anticoagulant or cardiovascular disease medication);

model 3: same as model 2 but MM characteristics dichotomized into: invasive and Breslow thickness \geq 0.8 mm or not, due to an empty cell.



