

Prophylactic Effect of Low Dose Vitamin D in Osteopenia of Prematurity: A Clinical Trial Study

Paymaneh Alizadeh¹, Negar Sajjadian^{1*}, Bahram Beyrami¹, Mamak Shariat²

¹ Department of Pediatrics, Bahrami Children Hospital, Tehran University of Medical Sciences, Tehran, Iran

² Research Maternofetal Center, Vali Asr Hospital, Tehran University of Medical Sciences, Tehran, Iran

Received: 10 Jul 2013; Accepted: 2 Nov. 2013

Abstract- Osteopenia of prematurity (OOP) is a preventable disease. Improved survival of newborns with OOP is associated with an increased incidence of OOP. The purpose of this study was to compare the prophylactic effects of two low doses of vitamin D (200 and 400 IU/Day) on the clinical, biochemical and radiological indices of the rickets of prematurity. In a randomized clinical trial, 60 preterm newborns with birth weight < 2000 g & gestational age < 37 weeks were randomly divided in two groups. Thirty newborns received 200 IU/d of vitamin D in group one and 30 ones received 400 IU/day of vitamin D in group two. On the 6th and 8th weeks of life, serum calcium, phosphate, alkaline phosphates, and 25 - hydroxy vitamin D concentrations were measured and x- ray of left wrist and physical examination were performed. Both groups had no difference in biochemical, radiological or clinical presentation of rickets. Current study indicated that low dose vitamin D (200 IU/Day) is enough for prevention of OOP.

© 2014 Tehran University of Medical Sciences. All rights reserved.

Acta Medica Iranica, **Article in Press** 16 May 2014.

Keywords: Osteopenia; Prematurity; Dose; Vitamin D

Introduction

Premature newborns are at risk of developing rickets or osteopenia that inversely is related to the intrauterine gestational age. Clinical rickets usually appears between the 6 to 8 weeks of postnatal age. A wrist x-ray at 6th to 8th wks of age remains a practical assessment of the presence of overt rickets. The rates of rickets among premature newborns under 1500gr and 1000gr are 30% and 50%, respectively. This rate increases among newborns with weight of 800gr to 73% (2, 4).

The etiology of rickets is multifactorial and includes phosphorous (P), calcium (Ca) and vitamin D deficiency. When premature babies are fed with human milk, both the Ca and phosphorous P supplies are insufficient. Human milk provides 25% of Ca and P needed for normal bone mineralization. To prevent OOP, vitamin D, Ca and P supplementation is needed. Risk factors for bone disease in preterm infants include: Maternal vitamin D deficiency, placental insufficiency and genetic etiology.

Postnatal risk factors include: inadequate supply of Ca and P by prolonged total parenteral nutrition (TPN), vitamin D deficiency, feeding with UN supplemented

human milk, and taking medications (diuretics, corticosteroid) and immobility.

Recommendations for vitamin D supply are different in Europe and America. The European Society of Pediatric Gastroenterology and Nutrition (ESPGAN) recommend a dose of 800-1600 IU/d (6). In literature the recommended dose of vitamin D for prevention of OOP is 400 to 1000 IU/day (2,3,4,7,11-16). The least dose that is reported to prevent OOP is 100IU/Day (6). The American Academy of Pediatrics (AAP) recommends 200 IU/d for infants except neonates (8). Several studies indicated that a daily vitamin D dose 200 IU/d is sufficient to maintain vitamin D status and normal activity, and higher doses of vitamin D may cause hypervitaminosis D that involves risk of hypercalcemia with subsequent complications (2).

The only study about OOP was performed by Alizadeh P et al, in Iran (1). In that study 400 IU/d of vitamin D could prevent OOP as well as 1000 IU/Day.

The goal of present study was to compare the prophylactic effects of two low doses of vitamin D (200 -400 IU/Day) on the clinical, biochemical and radiological indices of OOP.

Corresponding Author: N. Sajjadian

Department of Pediatrics, Bahrami Children Hospital, Tehran University of Medical Sciences, Tehran, Iran
Tel: +98 21 9125957275, Fax: +98 21 88633039, E-mail address: nsajjadian@yahoo.com

Materials and Methods

In this randomized clinical trial authors compared the effects of two low doses of vitamin D on biochemical, radiological and clinical manifestations of OOP. Sixty preterm infants (inpatients or outpatients) referred to Shariati Hospital, Tehran, Iran from May 2010 to May 2012, were enrolled in the study. The inclusion criteria were gestational age of <37 wks and birth weight < 2000 g. The exclusion criteria were taking specific medications interacting with vitamin D metabolism (eg: anticonvulsants, diuretics, corticosteroids and so on) in mother, diabetes mellitus in mother, previous IUGR or SGA baby, chronic use of furosemide in infant and being NPO (non per oral) for more than 2 wks. The withdrawal criteria were failure of taking vitamin D supplements according to the protocol of the study and failure to follow orders such as performing radiography.

Gestational age was determined based on the history of mothers' last menstrual periods, prenatal sonographic findings and postnatal physico-neurological examination. The newborns were randomly divided into two groups by block randomization of two, to receive a 200 IU/d vitamin D (group1) and 400 IU/d vitamin D (group 2) from the time they tolerated full enteral nutrition.

All parents received written instructions about the purpose and protocol of the study after signing an informed consent form which was approved by the local institutional review board for human investigations. All infants in group1 and 29 (96.7%) infants in group 2 could tolerate breast milk and received Ca supplement as Ca gluconate (90-120 mg/kg/d) and phosphate supplement as phosphate sandose, effervescent tablet (55-75 mg/kg/d). For infants deprived of human milk, premature formula (Prenon), enriched in Ca and phosphate, was prescribed. In the 6 to 8 wks of age, plasma Ca, serum inorganic phosphate and alkaline phosphatase and 25 (OH) vitamin D3 were measured; x-ray of left wrist and a through physical examination were carried out.

Ca and P concentration was analyzed using photometry analyser. Serum alkaline phosphatase

activity was measured using paranitrophenol reaction method and 25(OH) vitamin D3 by ELZA. Physical examination and x-ray evaluation were made by blinded expert neonatologists. Diagnosis of OOP was based on: 1-abnormal radiographic findings (osteopenia, bone fractures, intracortical resorption), losing the sharp border (fraying), changing from a convey or flat surface to more concave surface (cupping) of metaphyseal edge, widening of the distal end of the metaphysis.

2- Biochemical results including normal or low serum Ca, low serum phosphate and high serum alkaline phosphatase and low Serum 25 (OH) Vitamin D3 concentrations.

3-Signs and symptoms of rickets (craniotabes, rickets rosary, wide fontanel, Harrison groove, kyphosis/scoliosis, Potts belly).

Statistical analyses were performed by SPSS version12, software package (SPSS Inc, Chicago, IL), using Chi-square and student t-test. P values < 0.05 were considered statistically significant.

Results

In this study, 60 preterm infants in two groups received two prophylactic different doses of vitamin D 200 IU/Day in the first group and 400 IU/Day in the second group. All participated infants had GA of 26 to 36 weeks, and birth weight of 700 gr to 2000 gr.

No difference in two groups in the serum Ca (p. value=0.23), P (p.value = 0.20), Alkaline phosphatase (p.value = 0.91) and 25 - Hydroxy vitamin D (p.value = 0.86) were seen (Table 1).

No difference in two groups in the decreased calcification of the bone cortex (p.value = 0.25), fraying (p.value = 1), increased distance metaphyseal of radius and ulnar from wrist bones and density decrease (p.value=0) were found (Table 2).

No difference in two groups in the clinical features including wide fontanele (p.value= 0.71), widening of the wrist (p. value =0.55), Harrison groove (p.value = 0.31) were seen (Table3).

Rickets Rosary was not observed in any participants of the two groups (Table 3).

Table 1. Comparison of Radiological Results in two Groups

	Reduced Bone Content	Fraying	Increased Distance of Metaphyseal	Reduced Bone Density
200 IU/d	2 (%6.7)	(%16.7) 5	(%0) 0	(%0) 0
400 IU/d	(%20) 6	(%16.7) 5	(%0) 0	(%0) 0
P.Value	0.25	1	0	0

Table 2. Comparison of Biochemical Results in two Groups

	mean±SD Ca	mean±SD Serum P	mean±SD Serum Alkaline Phosphate	mean±SD 25(OH)D Serum
200 IU/d	9.49 ± 0.48	6.12 ± 0.82	796.5 ± 315.45	80.26 ± 24.5
400 IU/d	6.65±0.52	5.78±1.15	788.4±261.7	79.16 ± 26.15
P.Value	0.23	0.20	0.91	0.86

Table 3. Comparison of Clinical Results in two Groups

	Wide fontanels	Widening of the wrist	Harrison groove	Craniotables
200 IU/d	43 (%13.3)	(%3.3) 1	0	0
400 IU/d	(%16.7) 5	(%6.7) 2	(%3.3) 1	0
P.Value	0.71	0.55	0.31	0

Discussion

Rickets is one of the most important complications of prematurity. Incidence of the OOP is estimated to be 23-32% in very low birth weight infants (< 1500g) and 50% in infants < 1000 gr (2). Human milk and formula both supply insufficient Ca and P for normal bone mineralization (6,7,9,14,17-20). Clinical rickets appears between the 6th to 8th postnatal weeks.

DEXA (Dual Energy X-ray Absorptiometry) can measure Bone Mineral Content (BMC) but in routine clinical practice it is performed in few centers and has its own 'baggage', such as cost, availability and logistic difficulties (4, 8). Standard x-ray films are not an accurate assessment for bone demineralization because BMC must be decreased by 20% to 30% or more to be diagnosed by this method (2), however they can detect bone fractures, osteopenia, intracortical resorption, cupping, fraying and increased distance of the distal of the metaphysis (4). A wrist x-ray film at 6 to 8 weeks of age remains a practical assessment of the presence of rickets (2).

Simple biochemical indicators of bone mineralization such as serum alkaline phosphatase, and to some extent, serum phosphate and serum Ca and Serum 25 (OH) vitamin D may be an easy way of identifying metabolic bone disease in premature infants (4).

Provision of adequate amounts of Ca, P, and vitamin D significantly decreases the risk of rickets of prematurity (2-4).

Backstrom et al. compared the effects of low (200 IU/Day) and high (960 IU/Day) doses of vitamin D on bone mineral accretion, using DEXA, in two small groups of premature infants until three months old. They

noticed that there were no differences in bone mineral content and in bone mineral density at three and six months corrected age between the infants receiving low or high dose of vitamin D (15).

AAP has recommended a vitamin D dose of for preterm newborns (14). Koo et al., showed that even as little as 160 IU/Day of vitamin D maintain normal and stable vitamin D status in preterm infants received adequate mineral intake (13). According to Alizadeh P et al, two different doses of vitamin D (400IU/Day and 1000IU/Day) had the same effect in prevention of OOP (1).

Current study indicated that low dose of vitamin D intake; 200 IU/Day has the same protection in bone mineral content as 400IU/Day of vitamin D. The authors used biochemical, radiological and clinical criteria to diagnose OOP and there were no significant differences between the control and interventional group for biochemical and radiological features, as well as clinical manifestations.

In present study serum 1,25 (OH) D₃ concentration in both groups were within the reference limits, indicating sufficient 1-hydroxylation of vitamin D even with a low dose of vitamin D.

The limitation of this study was that authors didn't use DEXA for evaluating BMC.

In conclusion, the authors recommend 200 IU/Day of vitamin D supplement for premature newborns until they gain a normal term birth weight (3000- 3500 g); this low dose of vitamin D as well as 400IU/Day of vitamin D can prevent OOP.

References

1. Alizadeh P, Naderi F, Sotoudeh K. A Randomized Clinical

- Trial Prophylactic Effects of Vitamin D on Different Indices of Osteopenia of Prematurity. *Iranian J Public Health* 2006;35(3):58-63.
2. Rigo J, Mohamed MM, Curtis M. Disorders of calcium, phosphorus and magnesium Metabolism. In: Martin RJ, Fanaroff AA, Walsh MC, editors. *Neonatal Perinatal Medicine Diseases of the Fetus and Infant*. 9th ed, Philadelphia: Mosby Elsevier; 2010: p. 1550-3.
 3. Winston WK, Reginald C. Calcium and Magnesium Homeostasis. In: MacDonald MG, Seshia MM, Mullett MD, editors. *Avery's Neonatology patho physiology & Mangement of the Newborn*. 6th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2005: p. 845-71.
 4. Greenbaum LA. Rickets and hypervitaminosis D. In: Kliegman RM, Jenson HB, Behrman RH, et al, editors. *Nelson Textbook of Pediatrics*. 18th ed. Philadelphia: Elsevier; 2011: p. 261-3.
 5. Harrison CM, Gibson AT. Osteopenia in Preterm Infants. *Arch Dis Child Fetal Neonatal Ed* 2013;98(3):F272-5.
 6. Haschke F, Schilling R, Pietschnig B, et al. Calcium, phosphorus and vitamin D administration in infancy. *Monatsschr Kinderheilkd* 1992;140(9 suppl1):S13-6.
 7. Vachharajani AJ, Mathur AM, Rao R. Metabolic Bone Disease of Prematurity. *Neo Rev* 2009;10(8):e402-11.
 8. Sivert WL. Joint Compression Therapy in the Prevention of Osteopenia of Prematurity: Current research and future considerations. *SOPHIA*. (Accessed in May 2, 2014, at http://sophia.stkate.edu/cgi/viewcontent.cgi?article=1006&context=ma_nursing).
 9. Eliakim A, Nemet D. Osteopenia of Prematurity the role of exercise in prevention and treatment. *Pediatr Endocrinol Rev* 2005;2(4):675-82.
 10. Law Hs, So KW, Ng PC. Osteopenia in Neonates: A Review. *HK J Pediatr* 2007;12(2):118-24.
 11. Taylor SN, Hollis BW, Wanger CL. Vitamin D Needs of Preterm Infants. *Neo Rev* 2009;10(12):e590-9.
 12. Woonso K, Cheung P. Treatment and prevention of neonatal osteopenia. *Curr Paediatr* 2005;15(2):106-13.
 13. Koo WW, Krug-Wispé S, Neylan M, et al. Effect of Three levels of Vitamin D Intake in Preterm Infants Receiving High Mineral-Containing milk. *J Pediatr Gastroenterol Nutr* 1995;21(2):182-9.
 14. Gartner LM, Chairperson Kramer MS, Chalmers B, et al. American Academy of Pediatrics. Committee on Nutrition. *JOGN* 2005;34:367-72.
 15. Backstrom MC, MAKI R, Kuusela AL, et al. Maci. Randomised Controlled Trial of Vit D Supplementation on Bone Density and Biochemical Indices in Preterm Infants. *Arch Dis Child Fetal Neonatal Ed* 1999;80(3):F161-6.
 16. Cross BK, Vasquez E. Osteopenia of Prematurity Prevention & Treatment. Internet Scientific publications (Accessed in May 17, 2014, at <http://ispub.com/IJANP/4/2/7117>).
 17. Jacobus CH, Holick MF, Shao Q, et al. Hypervitaminosis D Associated with Drinking Milk. *N Engl J Med* 1992;326(18):1173-7.
 18. Laing IA, Glass EJ, Hendry GM, et al. Rickets of Prematurity: Calcium and Phosphorus Supplementation. *J Pediatr* 1985;106(2):265-8.
 19. Haschke F, Schilling R, Pietschnig B, et al. Calcium, Phosphorus and Vitamin D Administration in Infancy. Unsolved questions. *Monatsschr Kinderheilkd* 1992;140(9 Suppl 1):S13-6.
 20. Markestad T, Aksnes L, Finne PH, et al. Vitamin D Nutritional Status of Premature Infants supplemented with 500 IU vitamin D2 per day. *Acta Paediatr* 1983;72(4):517-20.
 21. Oyatsi DP, Musoke RN, Wasunna AO. Incidence of rickets Prematurity at Kenyatta National Hospital, Nairobi. *East Afr Med J* 1999;76(2):63-6.
 22. Agostoni C, Buonocore G, Carnielli VP, et al. Enteral Nutrient Supply for Preterm Infants: commentary from the European Society of Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. *J Pediatr Gastroenterol Nutr* 2010;50(1):85-91.