

PUBLISHED BY

# INTECH

open science | open minds

World's largest Science,  
Technology & Medicine  
Open Access book publisher



**3,100+**  
OPEN ACCESS BOOKS



**103,000+**  
INTERNATIONAL  
AUTHORS AND EDITORS



**102+ MILLION**  
DOWNLOADS



**BOOKS**  
DELIVERED TO  
151 COUNTRIES

AUTHORS AMONG  
**TOP 1%**  
MOST CITED SCIENTIST



**12.2%**  
AUTHORS AND EDITORS  
FROM TOP 500 UNIVERSITIES



Selection of our books indexed in the  
Book Citation Index in Web of Science™  
Core Collection (BKCI)

**WEB OF SCIENCE™**

Chapter from the book *A Critical Evaluation of Vitamin D - Basic Overview*

Downloaded from: <http://www.intechopen.com/books/a-critical-evaluation-of-vitamin-d-basic-overview>

Interested in publishing with InTechOpen?  
Contact us at [book.department@intechopen.com](mailto:book.department@intechopen.com)

---

# Vitamin D and Physical Activity

---

Nikolaos E. Koundourakis and Andrew N. Margioris

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/65103>

---

## Abstract


Vitamin D is synthesized in the skin following exposure to ultraviolet radiation, producing cholecalciferol, while only a small percentage of the circulating vitamin D is of exogenous origin deriving from food. Following two sequential hydroxylations, in the liver and in the kidneys, vitamin D is fully activated. Although its role in bone physiology and calcium homeostasis is well documented, there is emerging evidence that vitamin D exerts a plethora of additional effects on most tissues regulating the musculoskeletal, cardiovascular, and immune systems as well as energy homeostasis. Its deficiency/insufficiency poses a major public health problem observed in all age groups and regardless of latitude and insolation. In muscles, vitamin D deficiency is associated with a decline in neuromuscular function including muscular strength, walking speed, balance, jumping and sprinting performance, and aerobic capacity, although the evidence is still weak regarding its effects in the young and the athletes. Supplementation counteracts the negative effects of vitamin D deficiency on performance although in individuals with adequate levels of vitamin D, additional supplementation does not appear to enhance further physical capabilities. The aim of this chapter is to review our current understanding of diverse effects of vitamin D in physical performance in athletic and nonathletic populations.

**Keywords:** vitamin D, Physical activity, athletes, musculoskeletal system, cardiovascular system, immune system

---

## 1. Introduction

Vitamin D is synthesized in the skin from 7-dehydrocholesterol following exposure to ultraviolet radiation (cholecalciferol or D<sub>3</sub>), and only a small percentage is exogenous originating from food (ergocalciferol, D<sub>2</sub>) [1–3]. Following to sequential hydroxylations in the liver and the kidneys, vitamin D is activated (**Figure 1**). Vitamin D plays a crucial role in calcium



**Figure 1** Biochemical pathways of vitamin D production after exposure to UVB.

homeostasis in addition to its multiple effects in a plethora of tissues [3]. Indeed, vitamin D receptors (VDRs) are present in most tissues affecting the transcription of many genes [4–7]. Vitamin D is transported in the blood via vitamin D-binding protein [8, 9]. Vitamin D plays a role in the mineralization of type I collagen matrix in the skeleton [10]. Vitamin D and its nuclear receptor affect the expression of more than one thousand genes acting as a transcription factor [11]. Non-genomic effects of vitamin D have also been described [12–19].

## 2. Vitamin D deficiency

A surprisingly high prevalence of vitamin D insufficiency has been recently reported worldwide (**Table 1**) regardless of their insolation [20–23]. In Canada, 30–50 % of children and adults are vitamin D deficient and need vitamin D supplementation [24]. Similar data have been reported in Africa, Australia, Brazil, Middle East, Mongolia, and New Zealand documenting a high risk for vitamin D deficiency in adults and children [24–26]. Cross-sectional studies of vitamin D status in adolescents have found deficiency in 17–47 % with an increased risk in black and Hispanic teenagers [24–26]. It should be noted here that there are fewer reports regarding the prevalence of vitamin D in athletes. A high prevalence of vitamin D deficiency has been documented in athletes in both outdoor and indoor sports. For instance, it has been reported that more than 42 % of distance runners in Baton Rouge, Louisiana, were vitamin D insufficient or deficient [21]. Indeed, athletes may be more vulnerable to vitamin D deficiency than age-matched non-athletes even in regions with insolation. A recent meta-analysis pooling 23 studies composed of 2313 athletes found that 56 % of athletes had inadequate vitamin D [23]. In addition, a sizeable number of athletes do not meet the US dietary reference intake for vitamin D in addition to their inadequate endogenous synthesis when they train indoors [27]. In a recent study, Morton et al. [26] examined the prevalence of seasonal variation in vitamin D levels in 20 FA Premier League soccer players residing at a latitude of 53°N. Serum 25-

hydroxyvitamin D (25(OH)D) levels decreased in winter to insufficient levels in 65 % of the soccer players (<50 nmol/L). Koundourakis et al. [28] showed that professional Greek soccer players' training at a latitude of 35.9°N had insufficient vitamin D levels, while nearly identical levels were reported in American national football league players [29], in elite gymnasts in Australia [30], and in young Hawaiian skateboarders [31] and a variety of other athletes worldwide [32–35]. These findings were observed irrespective of sun exposure. In a recent study conducted in Israel in a favorable latitude (31.8°N) for insolation, 73 % of the athletes were vitamin D deficient. In Qatar, 84 % of soccer players were vitamin D insufficient. Furthermore, vitamin D deficiency in regions of high sunlight (high insolation) may be due to clothing, air pollution, and failure to meet the dietary reference intake for vitamin D in addition to their inadequate endogenous synthesis when they train indoors. Indeed, high prevalence of low vitamin D levels has been reported in the general population in the United Arab Emirates, India, and Egypt [36, 37].

Study	Population	Indoor/ outdoor	Season	Country, latitude	Vitamin D levels
Lovell [30]	18 female gymnasts, age = 13.6 ± 1.2 y	Indoor	Spring	Australia, 35.27°S	Insufficient 83.33 %, deficient 33.33 %
Constantini et al. [35]	98 Israeli male and female athletes and dancers, age = 14.7 ± 3.0 y	Outdoor/ indoor	Winter	Israel, 31°N	Insufficient 73 %, insufficiency was higher in dancers (94 %), basketball players (94 %), Tae Kwon Do fighters (67 %), athletes from indoor sports
Kopeć et al. [32]	24 Caucasian male Polish professional soccer players 26.24±3.41 y	Outdoor	Summer Winter	Poland, 51.10°N Cyprus, 30.11°N	Insufficient 37.5 %, deficient 12.5 % Insufficient 45.8 %, deficient 37.5 %
Morton et al. [26]	20 FA Premier League soccer players	Outdoor/ indoor	Winter	United Kingdom, 53°N	Insufficient 65 %
Galan et al. [40]	28 male soccer players, 26.7 ± 3.6 y	Outdoor/ indoor	Winter	Spain, 37°N	Deficient 64 %
Koundourakis et al. [28]	67 Caucasian male professional soccer players, age = 25.6 ± 6.2 y	Outdoor	Spring	Greece, 35.9° N	Insufficient 55.22 %
Halliday et al. [34]	93 male athletes (football, handball, shooting, squash, cycling, body building, martial arts), age = 21.3 ± 6.5 y	Outdoor/ indoor	Spring, summer	Middle East 29°N	Insufficient 91.39 %, deficient 8.60 %
Willis et al. [33]	19 male and female endurance-trained runners, age = 28.3 ± 3.4 y	Outdoor	N/A	United States, 30°N	Insufficient 42.10 %, deficient 10.52 %
Close et al. [211]	61 male athletes of soccer, rugby, jockeys, age = 24.3 ± 4.8 y	Outdoor/ indoor	Winter	United Kingdom, 50°N	Deficient 62 %
Close et al. [214]	30 male soccer and rugby athletes, age = 21.3 ± 1.3 y	Outdoor/ indoor	Winter	United Kingdom, 53°N	Deficient 57 %

Insufficient: vitamin D levels <30 ng/ml [75 nmol/L]; deficient: vitamin D levels < 20 ng/ml [50 nmol/L]; y = years.

**Table 1.** Vitamin D levels in various athletic populations.

Generally speaking, vitamin D insufficiency is defined (**Table 2**) as levels <30 ng/ml (80 nmol/L), whereas levels below 20 (50 nmol/L) and 10 ng/mL (25 nmol/L) represent deficiency and severe deficiency, respectively [7]. Levels between 40 and 60 ng/mL are the preferred range, while vitamin D intoxication usually does not occur until 25(OH)D3 reaches levels higher than 150 ng/ml [24].

Status	25-hydroxyvitamin D (ng/ml)	25-hydroxyvitamin D (nmol/L)
Severe deficiency	<10	<25
Deficiency	<20	<50
Insufficiency	20–30	50–75
Sufficiency	>30	>75
Toxicity	>150	>375

**Table 2.** Classification of vitamin D (25-hydroxyvitamin D) levels.

### 3. Seasonal variation of vitamin D levels

Seasonal variation of vitamin D is a well-documented phenomenon [38]. In the winter months, endogenous vitamin D production is drastically reduced, as a result of reduced exposure to UVB radiation, while in summer exposure to UVB radiation is adequate for vitamin D synthesis by the skin [5]. Similar variations have been reported for athletes. In soccer players, the levels of 25(OH)D3 are normal in 50 % of the athletes, while this number drops to 16.7 % in winter [32]. A study performed in a high latitude (Laramie, WY 41.3°N) revealed vitamin D insufficiency of 63 % in winter compared to only 12 % in autumn and 20 % in spring in both indoor and outdoor athletes [38]. Similar findings were reported in a study conducted at higher latitudes (Ellensburg, WA 46.9°N), using exclusively outdoor athletes. The authors reported that a percentage of 25–30 % of the athletes were vitamin D insufficient from fall to winter [39]. Morton et al. [26] also reported a significant drop of serum levels of 25(OH)D3 in a group of professional soccer players of the English Premier League at the latitude of 53°N between summer and winter. In agreement with these data were findings in professional soccer players in Spain that have been training at a latitude of 37°N showing a statistically significant reduction in serum levels of 25(OH)D3 between October and February [40].

Seasonal variation of athletic performance has been found to parallel that of the seasonal fluctuation of vitamin D levels. Indeed, athletic performance peaks when vitamin D levels are the highest, i.e., during summer [41]. Koch and Raschka [42] reviewed the German literature on the seasonality of physical performance. They reported that early studies indicated that strength and maximal oxygen uptake peaks ( $VO_{2max}$ ) in late summer. Comparable findings have been reported by Erikssen and Rodahl [43]. More specifically, in a population of 1835 Norwegian men, physical performance exhibited seasonal variability peaking in summer. Similar data were reported for the Swedish national track and field teams where  $VO_{2max}$  was observed in summer [44].

Most authors explain the enhanced athletic performance in summer to the higher vitamin D levels. However, other authors disagree [41]. According to Moran et al. [26], the higher physical performance in summer is a consequence of outdoor physical activity in the warmer weather which also results in exposure of athletes to UV, i.e., these authors see the two phenomena as parallel but independent. Athletes participating in athletic events such as track and field competitions have to be in their best conditioning in the summer since most major competitions take place at this time period. As a consequence, enhanced training and competition in summer would result in both better athletic performance and higher vitamin D levels [1, 45]. In contrast, participants in other sports such as soccer, American football, or basketball attain their best performance in winter while in summer recuperate resulting in a reduction and/or cessation of their training and to a deterioration of exercise performance and capacity [28]. Nevertheless, even when controlling for seasonal fluctuations in time spent exercising, variation in wrist flexor strength and muscle trainability still correlated with seasonal variations of vitamin 25(OH)D levels [41].

#### **4. Effects of vitamin D in calcium absorption and bone physiology**

Vitamin D plays an essential role in calcium homeostasis and the mineralization of bones [46, 47]. In the small intestine, vitamin D augments calcium uptake via an energy-dependent transcellular pathway and by a passive paracellular pathway through tight junctions [24, 48].

Insufficient levels of circulating vitamin D cause an increase of serum parathyroid hormone (PTH) due to secondary hyperparathyroidism, i.e., in a desperate effort of the body to keep circulating levels of calcium normal borrowing them from bones via an increase of bone turnover and increased bone resorption [26]. In chronic vitamin D-deficient states, individuals are susceptible to osteopenia, osteoporosis, and increased incidence of bone fractures [49, 50]. Administration of vitamin D is recommended as a first-line strategy reducing the incidence of osteoporotic fractures [51]. Indeed, low 25(OH)D levels in women could be viewed as a general marker of impaired health as the Malmö study has documented correlating with increased overall mortality [52].

Furthermore, low serum vitamin D is associated with cortical bone porosity indicative of increased bone fragility [53]. Vitamin D also enhances the activity of insulin-like growth factor 1 (IGF-1) via an induction the expression of its receptor, an affect crucial in bone formation both in vitro and in vivo [54].

##### **4.1. Implications of these data on physical activity and exercise performance**

Low vitamin D levels are related with osteoporosis and the risk of fractures in elderly individuals. Data from cross-sectional and longitudinal studies report that vitamin D insufficiency impairs bone health [4, 7]. As a result of this deterioration in bone health, the increased incident of falls in the elderly can easily result to bone fractures. However, recent evidence indicates that this is also evident in young physically active individuals. A study on Finish military recruits reported that those with insufficient vitamin D levels (<30 ng/mL) were susceptible

with a significant risk of developing stress fracture [55]. Similar findings have been reported in a number of other studies [56] including athletes [57, 58]. Deterioration in bone strength could be a problem apart from non-athletes, also in athletic populations. Hypothetically, athletes with low bone mineral density are predisposed to skeletal injury especially in contact individual and/or team sports [4, 7]. The latter could have several implications for the athletic performance especially in professional athletes. Notably the association between the incident of fractures and low vitamin D levels in athletes has not yet been determined, and there are also a number of studies in nonathletic populations that failed to support the hypothesis that high adequate vitamin D levels protect against stress fractures [4].

## 5. Effects of vitamin D on the skeletal muscle

The potential association between vitamin D and muscle function was also based on early clinical descriptions that myopathy was strongly associated with severe vitamin D deficiency [59]. Currently, vitamin D has been shown to be a potent modulator of skeletal muscle physiology [5]. Although the expression of the vitamin D receptors in the skeletal muscle tissue was questioned [60], recent data strongly indicate that they are also expressed in the skeletal muscle [61]. As we mentioned earlier,  $1,25(\text{OH})_2\text{D}$  by binding to its nuclear receptor augments the transcription of key muscular proteins [5–7]. It should be noted here that both genomic and non-genomic effects of vitamin D are crucial for muscular performance. Indeed, vitamin D affects muscular calcium and phosphate transport across cell membranes, phospholipid metabolism, and muscle cell proliferation and differentiation [62]. In addition, vitamin D downregulates the expression of myostatin (a negative regulator of muscle mass) and upregulates the expression of follistatin and insulin-like growth factor 2 (IGF-2) [5–7, 63]. Exposure of skeletal muscles to  $1,25$ -dihydroxyvitamin D<sub>3</sub> induces the expression of several myogenic markers and transcription factors [63]. A 10-day incubation of muscle cells with  $1,25$ -dihydroxyvitamin D<sub>3</sub> induces myosin heavy chain (MHC) type II, a late myogenic marker, while the mean diameter and width of muscular fibers increase compared to parallel controls. Indeed, vitamin D increases the size of MHC type II-positive myotubes [61, 64] and the cross-sectional area (CSA) of skeletal muscle fibers [65]. Vitamin D signaling has been also reported to alter the expression of C2C12 myotube size [5, 61], indicating a direct positive effect on the contractile filaments and thus muscle strength. In addition, it affects the diameter and number of type II muscle fibers [5–7] and in particular type IIa fibers [61]. In severe vitamin D deficiency, a proximal myopathy is observed characterized by type II (fast twitch) muscle fiber atrophy [5, 61]. A recent vitamin D supplementation study showed that in young male populations, a significant increase in the percentage of type IIa fibers was evident in the vitamin D group compared to the placebo group, demonstrating a positive effect on the muscle fiber type morphology, calcium transport, and regulation of intracellular calcium [61].

### 5.1. Implications of these data on physical activity and exercise performance

The data presented above suggest that vitamin D exerts beneficial effects on skeletal muscles and thus may improve physical performance capacity in both physically nonathletic individ-



uals and athletes. The documented vitamin D-mediated induction of muscle protein synthesis and myogenesis could result in muscles of higher quality and quantity. This could be in turn translated into increased muscle strength since it is well documented that there is a strong linear associations between muscle mass and strength [66]. In addition the association between vitamin D levels and muscle anabolism could also be related with another positive effect, which is particularly important for athletic populations in that it could accelerate muscle recovery from the stress of intense exercise [26]. This hypothetically could be of major importance in periods of continues competitions since it could enable them to maintain their performance capacity at a better level. In the elderly on the other hand, low vitamin D levels could cause a reduced muscular anabolism, sarcopenia, and thus negatively altered muscle mass which in turn leads to a negative effects in the sense of balance, strength, and stability. In addition, in vitamin D-insufficient states, active individuals may not have adequate energy provisions to sustain a moderate- to high-intensity muscular effort. Lastly, hypertrophy of type II muscle fibers could result in enhanced neuromuscular performance [4, 66]. These types of fibers are a major determinant of the explosive type of human movements resulting in high-power output. It is well documented that type II muscle fibers display a markedly faster muscle contraction velocity and thus higher force than type I muscle fibers [67]. Therefore the anaerobic maximal-intensity short-burst activities, such as jumping, sprinting, acceleration, deceleration, and change of direction, which are of crucial importance for the majority of the athletic events, are highly related with the type II muscle fibers. Interestingly, apart from athletes the importance of type II muscle fibers is highlighted also on the elderly. Reversal of type II fiber atrophy as a result of vitamin D supplementation is thought to account for approximately 20 % lower risk of falling [68]. Vitamin D deficiency apart from the atrophy of type II fibers in these individuals enables fat infiltration and consequently fibrosis (fibrosis is observed in different tissues including the muscle due to excess fat accumulation) which are important factors underpinning muscle quality and also may be a predictor of muscle function in older adult [69]. In closing, based on the available literature, we propose that vitamin D is beneficial for individuals since it increases muscle protein synthesis, adenosine triphosphate (ATP) concentration, strength, jump height, jumping and sprinting velocity and power, and aerobic and anaerobic exercise capacity. Physical performance could be significantly enhanced and/or preserved with adequate vitamin D. Vitamin D also prevents muscular degeneration and reverses myalgia [68, 70].

## **6. Effects of vitamin D on the cardiovascular system**

Vitamin D receptors are expressed in human myocardium as well as in vascular smooth muscles and the endothelium [5, 6]. The activated form of vitamin D, i.e., 1,25(OH)<sub>2</sub>D, participates in the structural remodeling of cardiac muscles and vascular tissues [71] and results in improvements in flow-mediated dilation and blood pressure [72]. Improved cardiac muscle function has been reported in patients with severe vitamin D deficiency following treatment [71]. In addition, animal studies have demonstrated that 1,25(OH)<sub>2</sub>D directly alters



myocyte contractility with accelerated relaxation, which is crucial in the normal cardiac diastolic function [6]. Vitamin D has been also found to regulate the function of calcium channels in cardiac myocytes, providing a rapid influx of calcium into cells promoting myocyte contractility [73]. Data from both animal and human studies show that vitamin D can act as a negative regulator of the renin-angiotensin-aldosterone system (RAAS) [6]. It is well documented that the RAAS maintains vascular resistance since it is a strong vasoconstrictor agent, due to angiotensin II synthesis, blood pressure, electrolyte, and intravascular fluid volume homeostasis. Vitamin D has the ability to decrease RAA activity by suppressing renin gene expression resulting to the downregulation of the RAA system [6, 45]. This is a vital mechanism that is regulated by vitamin D since it has been observed that the elevation in renin synthesis is highly related with left ventricular hypertrophy and the development of hypertension [45]. Another mechanism via which vitamin D insufficiency may induce left ventricular hypertrophy is by affecting and in particular elevating PTH levels. Observational studies have suggested that apart from hypertension, elevated PHT levels also result to left ventricular hypertrophy [74]. Vitamin D insufficiency has been found to be related with increased arterial stiffness and endothelial dysfunction in the conductance and resistance blood vessels in humans [75, 76]. Vitamin D has also some antiatherogenic functions by promoting HDL transport and inhibiting cholesterol uptake by the macrophages and the formation of foam cells [77], while its deficiency stimulates systemic and vascular inflammation, enabling atherogenesis [78]. In contrast, vitamin D has been shown to suppress inflammation via several pathways, such as inhibition of prostaglandin and cyclooxygenase pathways, upregulation of anti-inflammatory cytokines, decrease of cytokine-induced expression of adhesion molecules, reduction of matrix metalloproteinase, and downregulation of the RAA [45].

The presence of vitamin D receptors in the vascular wall suggests that it plays a role in vascular physiology and pathophysiology [74]. Among individuals with peripheral arterial disease, low vitamin D status has been associated with a faster decline of functional performance [79]. Severe vitamin D deficiency results in a disrupted adaptive immune response and an inflammatory milieu, promoting vascular dysfunction, insulin resistance, and arteriosclerosis [45, 80].

In addition, recent evidences indicate that severe vitamin D deficiency is strongly associated with sudden cardiac death [81, 82]. Furthermore, Vitamin D has been found to be a useful biomarker for the prediction of mortality when obtained at admission for chest pain [83]. The mechanisms via which vitamin D exert beneficial effects on human myocardium are as follows: vitamin D inhibits the proliferation of cardiomyoblasts by promoting cell-cycle arrest and enhances the formation of cardiomyotubes without inducing apoptosis. Vitamin D also attenuates left ventricular dysfunction in animal models and humans [84]. However, a group of researchers were unable to find any association between vitamin levels and risk of cardiovascular disease [85]. It should be noted though that analyzed data from more than 13,000 adults in the Third National Health and Nutrition Examination Survey (NHANES III) found a strong association between low vitamin D levels and key cardiovascular risk factors (i.e., hypertension, diabetes, overweight, and hypertriglyceridemia, but not hypercholesterolemia) after adjustment for multiple variables [86].

### 6.1. Implications of these data on physical activity and exercise performance

As per the available literature presented above, vitamin D plays an important role in the cardiovascular system [75]. Vitamin D affects several aspects of vascular health, including arterial stiffness and endothelial function which are crucial components of aerobic and anaerobic exercise performance and even the ability to perform daily activities. It should be noted that there is a linear association between vascular health and arterial stiffness to endurance capacity [80]. In addition, it is well documented that the most accurate measurement of aerobic capacity, i.e., that of  $VO_{2max}$ , is regulated by cardiac output, arterial oxygen content, shunting of blood to active muscle, and extraction of oxygen by muscles. Since low serum vitamin D levels may result in pathological myocardial hypertrophy, increased blood pressure, and endothelial dysfunction [87], these converging lines of evidence could support the idea that inadequate vitamin D levels could negatively influence cardiorespiratory fitness and the ability to perform efficiently during exercise. Thus, vitamin D affects aerobic capacity and  $VO_{2max}$ , cardiac output, vascular tone, and supply of oxygen and nutrients to the exercising muscles.

## 7. Effects of vitamin D on the nervous system

Vitamin D affects the central and peripheral nervous systems. Vitamin D receptors are present throughout the brain including the primary motor cortex, the region coordinating movement [88–91]. Specifically, VDRs have been identified in neuronal and glial cells in several brain areas including the cortex, deep gray matter, cerebellum, brainstem nuclei, spinal cord, and the ventricular system [90]. Furthermore, the enzyme  $1\alpha$ -hydroxylase, the activator of vitamin D precursors, is also present in the brain [89–93]. Vitamin D levels are associated with the conductance velocity of motor neurons and neurotransmission mediated by dopamine, serotonin, acetylcholine, GABA, and the catecholamines [92, 93]. Vitamin D also affects neuronal differentiation, maturation, and growth, its levels correlating to the levels of several neurotrophic factors including nerve growth factors (NGF) and that of neurotrophins, which play crucial roles in the maintenance and growth of neurons [94, 95]. In addition, vitamin D exerts direct neuroprotective effects via the synthesis of proteins binding calcium ( $Ca^{2+}$ ) ions which are important in neuronal function including transmission. Proper levels of neuronal calcium are critical since their excess may result in the formation of reactive oxygen species (ROS) which lead to neuronal damage. Indeed, vitamin D levels are inversely associated with oxidative stress which damages the brain leading to neuronal apoptosis or necrosis [96]. Vitamin D also affects neuroplasticity, a process via which neural synapses and pathways are adapted to the needs of environmental and behavioral demands adjusting the brain to noxious stimuli diseases or environmental cues [90, 97, 98]. The VDRs in glial cells play in the uptake and release of neurotransmitters, including that of GABA neurotransmission within the motor cortex [99]. GABAergic function is the principal “brake” within the brain affecting muscle relaxation via the corticospinal neurons [100].

### 7.1. Implications of these data on physical activity and exercise performance

The data described above suggest that vitamin D exerts major effects on several aspects of brain function. The effects of vitamin D on GABAergic tone and on serotonin and dopamine are crucial for muscular coordination and avoidance of central fatigue, a condition associated with the synaptic concentration of several neurotransmitters [101]. A high ratio of serotonin to dopamine affects exercise performance because of its effect on the general feeling of tiredness and the perceptions of effort [102].

Another mechanism via which vitamin D affects the brain and athletic performance may involve the nociceptors or nociceptors, i.e., the sensory nerve cell that responds to noxious stimuli by sending signals to the spinal cord and brain. The nociceptors are replete in VDRs and  $1\alpha$ -hydroxylase [103, 104]. When these receptors transfer pain signals to the brain, an inhibitory physical response takes place [104]. The relevance of this mechanism and vitamin D and physical activity/performance is based on the recent findings in animal studies that vitamin D depletion could result in nociceptive hyperinnervation and hypersensitivity within deep muscle tissue and a loss of balance without affecting muscle strength or the cutaneous nociceptive response [103]. Based on this finding, we could speculate that nociceptive hyperinnervation and hypersensitivity within deep muscle tissue could result to a false onset of myalgia during physical activity which in vitamin D-deficient individuals could result to a reduction in performance. The beneficial effects of vitamin D-mediated effects of the nervous system on physical activity and exercise performance are further supported by the findings of Dhesi et al. [105]. Reaction times deteriorate with age [105] and play an important role in neuroprotective responses [106]. In cross-sectional work, we identified a correlation between vitamin D status and reaction times, suggesting a neuroprotective role of this vitamin [105]. However, it is still unknown whether the latter mechanisms could be attributed also to young healthy individuals or athletic population.

## 8. Effects of vitamin D on pulmonary function

Vitamin D affects pulmonary function and health. Vitamin D insufficiency has been associated with impaired pulmonary function, asthma, and chronic obstructive pulmonary disease (COPD) [107, 108]. It should be noted that vitamin D deficiency may harm the lungs not only via its well-known immune effects. The Third National Health and Nutrition Examination Survey (NHANES III) reported a strong positive association between serum 25(OH)D and forced expiratory volume (FEV1) and forced vital capacity (FVC). These effects of vitamin D may be mediated through surfactant, a substance maintaining alveolar structural integrity [109], lung compliance, and oxygen exchange. It should be stressed that lung compliance, FEV1, FVC, and cardiopulmonary oxygen transfer are linearly related with physical performance [110]. Indeed, vitamin D induces the synthesis of alveolar surfactant and plays a critical role in epithelial-mesenchymal interactions during lung growth [111]. Vitamin D plays a role in COPD and more specifically in smooth muscle proliferation in the airways [112, 113]. Zosky et al. [114] have provided evidence supporting a direct role of vitamin D on lung growth in

vivo, while vitamin D deficiency results in lung volume deficits [115]. Finally, vitamin D insufficiency correlates with multiple indices of compromised lung function and increased airway reactivity [115].

### **8.1. Implication of these data on physical activity and exercise performance**

The data described above provide evidence implicating vitamin D in alveolar structural integrity, lung compliance, vital capacity, cardiopulmonary oxygen transfer, airway smooth muscle proliferation, and lung remodeling. Regarding athletes, exercise performance and aerobic capacity ( $VO_{2max}$ ) depend on all these lung functions [28]. Adequate  $VO_{2max}$  levels are needed in all sporting disciplines. Vitamin D appears to affect all components of  $VO_{2max}$  adequacy.

## **9. Effects of vitamin D on immunity**

Vitamin D affects the innate and adaptive immunity via VDRs [116–118]. Vitamin D enhances monocyte/macrophage activity. Vitamin D directly activates the transcription of several antimicrobial substances including defensin  $\beta 2$  (DEFB) and cathelicidin [118, 119]. Human cathelicidin (hCAP18) causes destabilization of microbial membranes [120]. In college athletes vitamin D insufficiency is associated with higher frequency of illnesses including common colds, influenza, and gastroenteritis [27]. Furthermore, vitamin D affects both T and B cells [121]. Under resting conditions the expressions of VDRs are low in both T and B cells, but they are upregulated in infectious diseases, suggesting a crucial role in adaptive immunity [122–125]. Indeed, vitamin D affects B-cell function, including inhibition of memory- and plasma-cell generation and apoptosis.

It is also known for years that immune cells can activate vitamin D to  $1,25(OH)_2D$  via  $1\alpha$ -hydroxylase, an effect resulting in several beneficial paracrine effects including modulation of T-cell function [126]. Vitamin D upregulates the expression of several antimicrobial peptides (AMP) while at the same time downregulates the expression of inflammatory cytokines including that of tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) and interleukin (IL) 6 [126]. Low levels of vitamin D in the general population and in athletes (especially following intense exercise) result in IL6 and TNF $\alpha$  increase. Vitamin D ameliorates this inflammatory response.

Several studies have also documented a negative association between vitamin D levels and upper respiratory tract infections (URTIs) in young and elderly adults [127–129]. In athletes, the incidence of respiratory illnesses is higher (especially at the elite level) because of the high demands on the lungs during prolonged exercise [130]. This could lead to increased exposure to viral and bacterial pathogens from the external environment, increasing the risk of URTIs. Li and Gleeson [131] have suggested that low vitamin D levels may exaggerate the vulnerability of athletes to URTIs, while individuals with higher vitamin D levels exhibit a lower propensity to URTI [132, 133]. It has been suggested that vitamin D prevents the development of URTIs by affecting the activation of toll-like receptors (TLRs) [134]. Vitamin D interferes with the activation of TLR signaling by microbial antigens via cathelicidin and  $\beta$ -defensin

[134]. Cathelicidin enhances the microbicidal capability of monocytes/macrophages by increasing the oxidative burst of these phagocytic cells, an effect also depended on vitamin D [134].

### **9.1. Implications of these data on physical activity and exercise performance**

It is reasonable to suggest that the well-documented association between optimal vitamin D levels and the well-being and the proper function of the immune system combined with the anti-inflammatory effects of this vitamin D may improve athletic performance. The observation that low vitamin D levels may result in increased pro-inflammatory cytokine levels, such as TNF $\alpha$  and IL6, following periods of intense exercise training may suggest that this pro-inflammatory period may be ameliorated by vitamin D [68]. This effect may accelerate the recovery phase from the stress of intense exercise. High IL6 and TNF $\alpha$  levels may partially explain age-related skeletal and muscular failings in vitamin D-deficient older individuals. Reduction of TNF $\alpha$  levels following vitamin D supplementation may be crucial to combat the “overtraining syndrome” in athletes [135]. Furthermore, the cytokines IL-10 and IL-13 have been shown to promote muscle regeneration and prevent skeletal muscle damage [136, 137]. Regarding the effects of vitamin D on the URTIs, its negative association with URTIs has been addressed by He et al. [138] who reported a higher proportion of subjects who experienced one or more URTI episodes in the vitamin D-deficient state during the 4-month study period than the optimal vitamin D group [138]. In closing, since vitamin D has been shown to affect the inflammatory response and prevent URTIs, its use may be of practical significance in the athletes.

## **10. Effects of vitamin D on the adipose tissue**

Vitamin D plays a crucial role in the physiology of adipocytes. VDRs and the 25-hydroxyvitamin D 1 $\alpha$ -hydroxylase (CYP27B1) genes are all expressed in human adipocytes [139]. Vitamin D levels are inversely related to obesity including body fat percentage (BFP), body mass index (BMI), and waist circumference [140, 141]. Vitamin D supplementation for 16 weeks reduces visceral fat in overweight and obese adults [142]. Drincic et al. [143] identified body weight as the single strongest predictor of 25OHD levels followed by fat mass. The loss of weight results in an elevation of vitamin D levels [144]. Epidemiologically, vitamin D levels are negatively associated with BFP in non-athletes [19, 145, 146]. Interestingly, a high BFP in winter months has been found correlating with indoor athletic activities and low vitamin D levels [146].

### **10.1. Implications for physical activity/exercise performance**

The close association between vitamin D levels and adiposity could affect athletic performance. It should be noted that neuromuscular performance capacity including sprinting ability, jumping performance, agility, acceleration, aerobic capacity, and deceleration is inversely

associated with BFP [147]. These data suggest that high BFP in conjunction with low vitamin D levels exerts a detrimental effect on exercise performance in athletes as well as in the general population. It should be noted that increased BFP is associated with the development of a systemic chronic low-grade inflammation and insulin resistance, a characteristic of obesity, which also negatively affect exercise performance [148].

## **11. Effects of vitamin D on the electron transport system**

Vitamin D has been also suggested to be related with the production of adenosine triphosphate (ATP) via the aerobic pathway. In particular, this speculation has been based on the observations that vitamin D appears to affect electron transport. Vitamin D receptors are present within mitochondria. Silencing of vitamin D receptors in several cell lines results in growth inhibition accompanied by an increase in mitochondrial membrane potential sensitizing cells to oxidative stress [149]. Vitamin D plays a role on mitochondrial respiratory chain activity acting as a facilitator of the diversion of acetyl-CoA from the energy-producing TCA cycle toward biosynthetic pathways that are essential for cellular proliferation. These effects of vitamin D on aerobic energy production during exercise are needless to say crucial for exercise performance.

## **12. Insolation and vitamin D on athletic performance**

Early studies on collegiate athletes and students showed that cardiovascular fitness, muscle endurance, and speed were enhanced following exposure to ultraviolet radiation [150–152]. In particular, in 1938 Russian authors [151] reported that a course of ultraviolet irradiation improved speed in the 100 m dash in four students compared with matched controls. The authors observed that the experimental group (i.e., under UV treatment) had a 7.4 % increase in 100 m dash times compared to only 1.7 % of the control group. Notably both groups were undergoing the same training schedule throughout the study. Supportive evidence came from a study by German researchers in 1944 [152]. A treatment under UV irradiation twice a week for a period of 6 weeks resulted in a 13 % improvement in performance on a bike ergometer. No alteration was evident in the control group. Enhanced performance parameters were also reported by Allen and Cureton [150] as a result to UV treatment in college students. The authors observed that a period of 10 weeks under UV treatment resulted in improvements in both cardiovascular fitness and muscular endurance. UV increased cardiovascular fitness by 19.2 % compared to only 1.5 % improvement in the controls. In 1956, Sigmund [153] examined the effects of UV radiation on reaction times on adolescents and adult individuals. UV treatment improved by 17 % the reaction time compared to controls. In another study in the late 1960s, a single dose of UV irradiation beneficially affected the strength, speed, and endurance of college women [154]. Interestingly, the beneficial effect was evident in white participants but not in African American females, suggesting a different skin response to UV. A follow-up study from the same team [155] observed that the same amount of UV managed



to improve aerobic performance in the same population. Enhanced performance was also observed by another study in a 30-yard dash of 15 college women after 6-minute exposure in UV light [156].

### 13. Vitamin D deficiency in athletes versus non-athletes

It is increasingly apparent that athletes need higher levels of vitamin D because of the higher demands of the daily strenuous training exercise which they undergo [157]. Indeed, experimental findings suggest a major role of vitamin D in muscle mass, strength, and function. Vitamin D affects significantly the athletic ergometrics [158]. Furthermore, athletic performance appears to show a seasonal variability peaking when vitamin D levels peak and declining as they decline reaching nadir when vitamin D levels are at their lowest. It should be noted that vitamin D administration improves athletic performance in vitamin D-deficient athletes [41].

### 14. Vitamin D in non-athletes

Overwhelming evidence demonstrates the association between vitamin D levels and the ability of non-athletes to perform their daily physical activities and/or exercise (**Table 3**). Early on, myopathy and muscle weakness were attributed to the most obvious consequence of vitamin D deficiency, i.e., that of rickets and osteomalacia [159]. To date, several cross-sectional observations and longitudinal studies have reported a close association between vitamin D and several parameters of physical performance. The majority of the studies have been performed on elderly subjects although similar data have been reported in adults below the age of 65 years [160] and in young individuals [161, 162].

Author	Participants	Physical performance parameters tested	Results
Ward et al. [161]	99 adolescent girls, 12–14 y	Jump power, countermovement jump(CMJ) height, and velocity, Esslinger Fitness Index, force	Positive relationship between vitamin D levels and jump velocity, jump height, power, Esslinger Fitness Index, and force
Foo et al. [162]	301 healthy Chinese adolescent girls, age 15.0 ± 04 y	Handgrip strength, physical activity levels	Adequate vitamin D status had greater handgrip muscle strength than deficient
Gerdhem et al. 155	1044 elderly women, 75 y old (range 75.0–75.9 y old)	30 m gait, knee extension, knee flexion, Romberg balance, self-estimated activity level	Positive relationship between vitamin D levels and gait speed, Romberg balance



Author	Participants	Physical performance parameters tested	Results
			test, self-estimated activity level, and thigh muscle strength
Houston et al. [166]	988 elderly males and females, age $85.2 \pm 3.6$ y	SPPB test (standing balance, repeated chair stands, 3 m walk [gait speed]), grip and knee extensor strength	Significantly poorer SPPB in deficient individuals versus those in sufficient state. No relationship between vitamin D levels and absolute grip and knee extensor strength. Older adults with deficient 25(OH)D levels had significantly lower muscle strength indicating poorer muscle quality
Visser et al. [167]	1260 elderly males and females >65 y old	Physical activity (30, 30–59, or 60 min/day), timed walking test, and a repeated chair stand test	Vitamin D-deficient and vitamin D-insufficient individuals had poorer physical performance in all tests
Bischoff-Ferrari et al. [168]	4100 elderly males and females >60 y old	8-foot walk test and sit-to-stand test	Vitamin D levels between 40 and 94 nmol/L were associated with better musculoskeletal function in the lower extremities than concentrations of 40 nmol/L. Significant positive association between vitamin D levels and the 8-foot walk test, the sit-to-stand test
Tieland et al. [169]	127 pre-frail and frail elderly people, $\geq 65$ y old	Short physical performance battery (SPPB) test, maximum strength (one repetition maximum)	Significant positive associations were observed between vitamin D levels and gait speed and chair rise
Houston et al. [174]	1155 elderly individuals, aged 65–102 y old	Short physical performance battery (SPPB), handgrip strength	Significant positive associations were observed between vitamin D levels and SPPB score in men and handgrip strength in men and women
Wicherts et al. [177]	1234 elderly men and women, age $79.8 \pm 5.9$ y	Time taken to walk 3 m, turn 180°, and walk back (walking test); time taken to rise five times from a kitchen chair with arms folded in front of the chest (chair stands); the ability to stand with the heel of the one foot directly in front of, and touching the toes of, the other foot for at least 10 s (tandem stand)	Vitamin D levels were associated with physical performance which was poorer in participants with levels < 10 ng/ml and levels of 10–20 ng/ml
Mastaglia et al. [178]	44 postmenopausal women, age $71 \pm 4$ y	Walking-speed test, standing balance, and sit-to-stand tests.	Women with vitamin D levels $\geq 20$ ng/ml scored higher on the muscle function

Author	Participants	Physical performance parameters tested	Results
		Lower extremity muscle strength was determined using a manual dynamometer	tests and had stronger knee extensor and hip abductor muscles
Ceglia et al. [181]	1219 males, age 4.9 ± 12.8 y	Timed walking test, chair stand test, grip strength	No association was evident between any physical performance test and vitamin D levels
Mathei et al. [182]	367 elderly males and females, age 84.7 ± 3.6 y old	Balance, grip strength, and gait speed	No significant relation between vitamin D levels and balance, gait speed and grip strength, and serum
	y, years		

**Table 3.** Vitamin D levels and physical performance in non-athletes.

In the elderly vitamin D deficiency has been associated with myopathy, reduced muscle mass and strength, low exercise performance, and an increased risk of falling [163]. The rise of vitamin D levels has been associated to improved balance [164], reduced frailty and disability [165, 166], independence [167], and better mobility [168, 169]. Improved balance and mobility independent of muscle function may partly explain the association of vitamin D with the reduced risk of falls in older adults. It is well documented that the maintenance of muscle strength and power in this population is also related with the ability to perform all types of physical activities and therefore well-being and functional performance. In the elderly, mobility limitations have been defined as the self-reported inability to walk a mile, climb stairs, or perform heavy housework [170]. This is dependent on multiple components such as muscles, bones, and the cardiovascular system, and vitamin D has been found to play a crucial role in the functionality of these tissues. The impairment in any of these tissues could lead to reduced mobility that in turn could cause accelerated functional decline and disability. Given that both muscle strength and power are related to the functional activities in older adults [161, 171], it could be argued that assessing the potential parameters that could affect muscle strength or power and even muscle quality would identify ways to avoid impaired mobility and functional performance. In addition, proper cardiovascular function is essential to perform adequately all types of activities. In this context, it could be suggested that vitamin D is involved in the regulation of strength and power (as a product of strength) and affects the proper cardiovascular function and also the quality of the muscles (fat infiltration in vitamin D-deficient state). The latter is related with vitamin D since in the deficient state, it has been reported that there is increased fat infiltration in the muscle. This physiological mechanism is associated with lower levels of muscle strength and physical performance, independent of muscle mass [172]. To date, a number of organizations recognize that vitamin D is directly related with the falls in the elderly, including the International Osteoporosis Foundation, the Endocrine Society, and the US Preventive Services Task Force [173]. In accordance are the observations of Gerdhem et al. [164]. The authors evaluated the association between 25-

hydroxyvitamin D levels (25OHD), fall-associated variables (including tests of functional performance), and fracture in ambulatory women. A low 25OHD was associated with lower physical activity, gait speed, and balance. Vitamin D levels below 20 ng/ml have been associated with an increased risk of fractures. It has been suggested that the minimum level of 25OHD required to maximize the reduction of falls and fractures appears to be in the order of 60 nmol/L. Any increases above this do not appear to have any additional beneficial effect in lowering the risk of falling [173].

Regarding physical performance per se, several cross-sectional studies indicate a positive association between physical performance and serum vitamin D in the elderly. In the INCHIANTI study, a representative sample of 976 persons aged 65 years or older at study baseline was examined [174]. Physical performance was assessed. A significant association between low levels of vitamin D and poor physical performance as assessed by the handgrip strength test and a short physical performance battery (SPPB) test (which included ability to stand from a chair and ability to maintain balance in progressively more challenging positions) was documented. It was found that individuals with serum vitamin D levels <25 nmol/L performed worse than those with levels above 25 nmol/L. Muscle strength using a handgrip test was also significantly greater in subjects with vitamin D levels higher than 50 nmol/L than those with lower levels [174]. The use of grip strength as a tool is relatively straightforward to stratify adults at risk of impaired mobility. It has been shown in elderly individuals that higher levels of vitamin D correlated with better lower extremity muscle performance compared to age-matched individuals with lower levels [169]. Similar data have been shown in the Rancho Bernardo Study cohort [175]. Likewise in a longitudinal survey of community-dwelling Japanese, older women showed that those with higher levels of vitamin D responded better in a 3-month exercise program than those with lower vitamin D levels [176].

Wicherts et al. [177] examined the association of serum 25-hydroxyvitamin D levels to physical performance and its decline with age. The authors reported that after adjustment for age, gender, chronic diseases, degree of urbanization, body mass index, and alcohol consumption, a relationship was evident between vitamin D levels and physical performance indices. Physical performance was inferior in individuals with low vitamin D compared to age-matched individuals with serum 25OHD levels higher than 30 ng/ml. Thus 25OHD levels below 20 ng/ml appear to be associated with poorer physical performance and a greater decline in physical performance in older men and women.

In another study on elderly females aged over 65 years, the authors examined the association between vitamin D levels and muscle function and strength. Performance was evaluated by a walking-speed test, standing balance, and a sit-to-stand test [178]. It was observed that vitamin D levels above 50 nmol/L were related with higher levels of muscle strength. The authors also reported that the individuals with the lowest 25OHD levels (i.e., below 20 nmol/L) had the poorest values in strength. Stockton et al. [179], based on the observations from their meta-analysis, concluded that vitamin D had no significant effect on lower extremity muscle strength except in individuals with starting serum 25OHD levels <25 nmol/L. In support to this suggestion are the findings of Pramyothin et al. [180]. In a study performed in older Hawaiian women of Japanese ancestry, which are known for its very low rate of falls, a high dietary intake

in vitamin D, and a large exposure to sunlight, their mean vitamin D level was 80 nmol/L, while no individuals were in a vitamin D-deficient state. The authors failed to observe any association between vitamin D levels and all the measured performance parameters, i.e., physical strength test (except for the quadriceps), falls, and daily activities. It was concluded that the absence of a relationship was due to the very high level of vitamin D that these women had at baseline. Mastaglia et al. [178] suggest that there is a limit below which vitamin D insufficiency may affect physical performance. In agreement to this hypothesis are the findings of a recent study [181]. The authors reported that in a population-based sample of adult men with a broad age range (mean age is 47 years), there was no association between serum 25(OH)D concentration and lean body mass, muscle strength, and physical function after controlling for multiple lifestyle factors. However, only 20 % of the subjects had a vitamin D level below 50 nmol/L. This observation is giving further support to the cause-effect hypotheses of “sufficient” or “insufficient” vitamin D levels on several indices on human physiology including physical function and athletic performance. However confusing findings have been reported by Matheï et al. [182] in elderly individuals (>80 years) with an 80 % prevalence of vitamin D insufficiency where no correlation was evident between vitamin D levels and physical performance, as assessed by gait speed, handgrip test, and a static balance test. The authors suggest that this absence of association could be at least partly due to age-related downregulation of the VDR in muscles [182]. On the other hand, Ward et al. [161] reported that in vitamin D-insufficient young adolescent girls, there was a correlation between vitamin D levels and muscle power, force, velocity, and jump height. Similarly, Foo et al. [162] found a positive association between vitamin D serum levels and handgrip strength in a population of 301 vitamin D-sufficient adolescent girls. However, studies in young women failed to verify any association between vitamin D status and muscle performance, handgrip strength [183], or any other physical performance parameter [184]. These discrepancies may be partially explained by differences in the experimental protocols, accuracy of vitamin D serum measurements, age of the participants, muscle fitness, health state, gender, and other variables. In summary, most published data suggest an association of vitamin D levels greater than 50 nmol/L with muscle fitness. Vitamin D-associated improvement of balance and mobility are independent variables from muscle function and may partially explain the beneficial effect of vitamin D supplementation on the rate of falls in older individuals [185].

## 15. Vitamin supplementation in non-athletes

Vitamin D levels in the blood are increasingly recognized as an important factor in muscular well-being and functional performance. As a consequence, it was speculated that vitamin D3 supplementation may exert beneficial effects on physical activity. Indeed, several major trials have demonstrated that vitamin D supplementation lower the risk of falling (**Table 4**). Prince et al. [186] in a randomized, controlled trial in Australia evaluated women with at least one fall in a 12-month period and with a plasma 25-hydroxyvitamin D level <24.0 ng/mL. The authors suggested that elderly with a history of falling and vitamin D insufficiency living in sunny climates benefit from ergocalciferol supplementation resulting in a 19 % reduction in

the relative risk of falling, mostly in winter. Pfeifer et al. [187] demonstrated a reduction in falls of 27–39 % in community-dwelling seniors supplemented with 800 IU vitamin D and calcium daily versus calcium alone. This drop in falls was correlated with an improvement in quadriceps strength and in the timed up and go (TUG) test. Similarly, Zhu et al. [188] reported an enhanced muscular strength and TUG test in the individuals within the lowest vitamin D quartile. These results are consistent with another study that showed a 49 % reduction of falls in elderly women from a geriatric ward supplemented with 800 IU per day of vitamin D [189]. Bischoff-Ferrari et al. [189] evaluated several studies involving men and women with an average age of 70 that vitamin D supplementation resulted in a significant 22 % decrease in fall risk.

Study	Quality	Participants	Intervention	Parameters measured	Results
Prince et al. [186]	Double-blind, randomized controlled trial	302 community-dwelling ambulant older women, aged 70–90 y old	1000 mg/day of calcium for 1 year combined with 1000 IU/day of ergocalciferol or identical	Risk of falling	Ergocalciferol supplementation resulted in a 19 % reduction in the relative risk of falling
Zhu et al. [188]	Randomized controlled trial	302 community-dwelling ambulant elderly women; mean age: 76.9 ± 4.5 y	1000 IU/day of vitamin D2 or identical placebo; calcium citrate (1 g calcium/day) in both groups for 1 y	Timed up and go (TUG) test, ankle dorsiflexion, knee flexor, knee extensor, hip abductor, hip flexor, hip extensor, and hip adductor strength were assessed	Significant ↑ in knee flexor strength and all hip muscle strength and TUG in both groups. Ankle dorsiflexion strength significantly ↓ in both groups, no change in knee extensor strength after 12 months
Bunout et al. [191]	Randomized double-blind controlled trial	96 healthy elderly subjects aged > 70 y	Participants were randomized to a resistance training or control group. Trained and control groups were further randomized to receive vitamin D 400 IU plus 800 mg of calcium per day or calcium alone for 9 months	Handgrip strength, Isometric quadriceps maximum voluntary strength, endurance, general physical fitness, measuring the timed up and go (TUG), short physical performance battery (SPPB)	Timed up and go ↑ more in trained subjects supplemented with vitamin D. At the end of the follow-up, gait speed was ↑ among subjects supplemented with vitamin D (whether trained or not)
Bischoff et al.	Randomized double-blind controlled	122 elderly women, mean age = 85.15 ±	Calcium 1200 mg and 800 IU/day vitamin D	TUG test, grip a strength, knee flexor strength, knee	↓ Risk of falls and in 62 women with complete for all strength test,

Study	Quality	Participants	Intervention	Parameters measured	Results
[192]	trial	6.8 y	versus calcium 1200 mg/day for 12 weeks	extensor strength, risk of falls	there was significant ↑ in all measured parameters
Moreira-Pfrimer et al. [193]	Randomized double-blind controlled trial	46 individuals (11 males and 35 females), age = 77.6 8 ± 8.2 y	6-month period: daily calcium + monthly placebo or daily calcium + oral cholecalciferol (150,000 IU once/month the first 2 months, 90,000 IU once/month until the end)	Maximum isometric strength of hip flexors (SHF) and knee extensors (SKE)	SHF ↑ in the calcium/vitamin D group by 16.4 % and SKE ↑ by 24.6 %
Pfeifer et al. [194]	Randomized double-blind controlled trial	242 community-dwelling elderly women and men, mean age 77 ± 4 y	1000 mg of Ca or 1000 mg of Ca plus 800 IU of vitamin D per day for 12 months, followed by an 8-month treatment-free observation period	Timed up and go (TUG) test, maximum isometric leg extensor strength	Vitamin D resulted in a significant ↓ falls of 27 % at month 12 and 39 % at month 20 and in significant ↑ and ↓ in quadriceps strength and time to perform the TUG test, respectively
Janssen et al. [196]	Randomized double-blind placebo-controlled trial	70 female geriatric patients >65 y old	400 IU/day of cholecalciferol + 500 mg/day of Ca or a placebo +500 mg/day of Ca for 6 months	Handgrip strength, leg extension power, get up and go (GUT) test, modified Cooper test	No significant improvements in strength and physical activity measures
Brunner et al. [197]	Randomized, double-blind, placebo-controlled trial	33,067 women aged 50–79 y old	1000 mg calcium carbonate plus 400 IU vitamin D-3 or matching placebo in a regimen of two pills per day	Grip strength, chair stand test, timed walk test, physical activity status, physical function (self-report), physical activity (self-report)	No evidence of treatment effects on measures of any kind of physical functioning, activity, and strength were observed
Gupta et al. [200]	Randomized, double-blind, placebo-controlled trial	40 healthy males and females, age 31.5 ± 5.0 y	60,000 IU of vitamin D3/w for 8 weeks followed by 60,000 IU/m for 4 months with 1 g of Ca daily or dual placebos for 6 months	Handgrip and gastro-soleus dynamometry, pinch-grip strength, respiratory pressures, 6 min walk test	All measured physical performance parameters ↑ significantly in the treatment group

Study	Quality	Participants	Intervention	Parameters measured	Results
Owens et al. [201]	Placebo-controlled trial	29 male participants, mean age = 22.7 ± 3 y	Oral dose of 10,000 IU/day vitamin D3 or a visually identical placebo for 3 months	Lower limb muscle function, isokinetic torque, percutaneous isometric electromyostimulation	No significant changes in any of the measured performance indices

y, years; ↑, increase; ↓, decrease

**Table 4.** Vitamin supplementation in nonathletic population.

Recent studies on elderly populations support the data regarding the beneficial effect of vitamin D supplementation on neuromuscular performance improving strength [9] and force production [190]. Furthermore, a significant linear association between vitamin D levels and muscular strength, body sway, and physical performance is evident [106, 191, 192]. In addition, Moreira-Pfrimer et al. [193] reported that vitamin D supplementation results in improved isometric muscle strength following a 6-month supplementation of daily cholecalciferol resulting in an increase of strength of hip flexors (SHF) and strength of knee extensors (SKE).

In addition to the strength of lower extremities, upper extremity strength also benefits from vitamin D supplementation. However, regarding grip strength, a tool that is relatively straightforward to stratify adults at risk of impaired mobility, the results are confusing. In a recent review of the literature, it was reported that grip strength does not seem to respond to vitamin D supplementation in males [194]. The failure to observe any association between grip strength in the male population is in agreement with others. Effects of vitamin D treatment on handgrip strength have been evaluated in several studies, none of them showing significant effects in contrast to women [191, 194–197]. This could be related with the gender of the participants in the studies. A recent research suggests that there are gender differences in the vitamin D that affect functional performance [198].

In an attempt to examine the effects of vitamin D supplementation concomitantly with resistance training, Agergaard et al. [61] performed a study in young and elderly untrained males. The participants of this study were randomized to a daily supplementation of vitamin D plus calcium or just calcium for a period of 16 weeks. The study took place during a period and at latitude of low sunlight (December–April, 56°N). All individuals followed a progressive resistance training program of the quadriceps muscle during the last 12 weeks of the study. Measurements of muscle hypertrophy (defined as changes in CSA) and quadriceps isometric strength levels were performed. Muscle biopsies were analyzed for fiber type morphology, expression of VDR, and Myostatin [61]. The authors failed to observe any difference between the groups in quadriceps CSA and isometric strength despite the observed increase in strength that was evident in both groups compared to baseline. Vitamin D intake and strength training increased both strength and CSA in elderly individuals compared to the young group. In the young vitamin D group, the authors observed that the experimental period resulted in an increase in fiber type IIa percentage and a lower Myostatin mRNA expression compared to the control group. No additive effect of vitamin D intake during 12 weeks of resistance training



could be detected on either whole muscle hypertrophy or muscle strength. The authors attributed this lack of response to vitamin D supplementation to the fact that the participants in their study were vitamin D sufficient, compared to the findings of studies that have observed enhanced performance to vitamin D supplementation that have employed individuals in a deficient state [191–193, 196, 199, 200]. They suggest that the muscular performance benefits may be relevant only for individuals with vitamin D deficiency. However, a recent study using insufficient young males failed to report improvements to performance via vitamin D treatment [201]. The authors observed that despite the elevation in vitamin D levels ( $>120$  nmol/L), no effect was observed in performance parameters (muscle strength evaluated by isokinetic dynamometry and percutaneous isometric electromyostimulation). The latter study suggests that other parameters apart from vitamin D levels are responsible for the effect on performance such as age, type of vitamin D that was given, period of supplementation, and amount of supplementation.

A recent review of the literature which included controlled and randomized controlled trials that have measured muscle strength and serum vitamin concentration in 18–40-year-old participants reported that the outcome of the available evidence is that vitamin D supplementation increases upper and lower limb strength in young and healthy individuals [200, 200, 202]. However, this increase seems to be dependent on the supplementation treatment employed. According to Tomlinson et al. [202], studies that have reported individual significant increases in muscle strength employed a total dosage of between 60,000 and 14,000 IU vitamin D/week over 6–4 months. Another study prescribed a daily dose of 2000 IU vitamin D as per previously published systematic reviews which suggested that daily dosages are more beneficial than larger, staggered doses. However, adding to the contradiction, Gupta et al. [200], who also reported significant results, used a large, weekly dose of vitamin D, 60,000 IU D3/week for the first 8 weeks followed by 60,000 IU/month for 4 months. The findings of these studies contradict previous evidence that suggested larger weekly, monthly, or one-off doses of vitamin D are not as effective in producing an improvement in muscular strength. According to Tomlinson et al. [202], the inconsistencies in the literature are partly the result of different dosages of vitamin D and different treatment regimens.

## 16. Vitamin D levels, Athletes, and exercise performance

The first evidence regarding the possible association between vitamin D and performance comes from early studies [41]. It has been reported that cardiovascular fitness, muscle endurance, and speed were enhanced after exposure to ultraviolet radiation [41].

To date there is a growing number of evidence (**Table 5**) to support the association between this secosteroid and exercise performance indices in athletic and physically active populations. A recent study from our laboratory [28] reported a linear relationship between vitamin D levels and muscle strength as evaluated by squat jump (SJ) and countermovement jump (CMJ), sprinting ability (10 m and 20 m), and  $VO_{2max}$  in non-supplemented professional soccer players. Our results are comparable with others showing that vitamin D levels are related with

neuromuscular performance capacity and aerobic endurance in young physically active individuals and athletes [202–205]. However, these findings are not universal. Others have failed to find an association between exercise performance and vitamin D levels [206]. Vitamin D status was not associated with grip strength or swimming performance in adolescent swimmers [207] nor with isokinetic peak torque during knee flexion and extension after adjusting for total body mass and lean mass in Qatar soccer players [208]. Hamilton et al. [208] reported that vitamin D levels were not related with lower limb isokinetic muscle function in soccer players. According to the authors, the lack of an association was most probably a result of the different modes of exercise used in their study. They hypothesized that vitamin D could preferentially affect muscle groups that were not evaluated in the study. In this study, soccer players with vitamin D levels <50 nmol/L exhibited significantly lower torque in hamstring and quadriceps muscle groups, suggesting that low vitamin D levels are associated with impaired muscular function during exercise. However, no association was observed between performance and vitamin D levels in hockey players [206]. Indeed, low explosive strength during jumping was not associated with vitamin D levels neither status nor power production during the Wingate anaerobic test [206].

Study	No. of participants	Sport	Study location/period	Parameters measured	Results
Koundourakis et al. [28]	67 male professional players, age 25.6 ± 6.2 y	Soccer	6-week off-season transition period, June–July, 35.9°N latitude	Squat jump (SJ), countermovement jump (CMJ), 10m sprint, 20 m sprint, maximal oxygen consumption (VO <sub>2max</sub> )	Significant associations between vitamin D levels and all the measured performance indices
Fitzgerald et al. [206]	53 junior and collegiate male ice hockey players, age 20.1 ± 1.5 y	Hockey	Off-season period, 44.9°N latitude	Grip strength, Wingate anaerobic test, squat jump (SJ)	Significant associations between vitamin D levels and grip strength, time to half peak force but not Wingate test and SJ
Forney et al. [205]	39 males and females, physically active individuals, age 23.26 ± 0.7 y	Various types of physical activity	July and September, 30.46°N latitude	Maximal oxygen consumption (VO <sub>2max</sub> ), Wingate test, vertical and horizontal jump, sit and reach test, maximum muscle strength of bench press, leg curl, leg extension, upright row, bicep curl, triceps pushdown	Significant positive associations between vitamin D levels and VO <sub>2max</sub> , but not for any measures of strength and power
Dubnov-Raz et al. [207]	80 competitive adolescent swimmers from both sexes, age 14 ± 1.6 y	Swimming	Israel, located between latitudes 32.40°N and 31.55°N	Grip strength, balance, and swimming performance at several speeds	No significant associations between vitamin D levels and any measures of performance

y, years

**Table 5.** Vitamin D levels and athletes.

Regarding  $\text{VO}_{2\text{max}}$  and aerobic capacity, evidence from large epidemiologic cross-sectional investigations indicate that 25(OH)D concentration is associated with cardiovascular fitness. Mowry et al. [204] found a positive correlation between baseline cardiorespiratory fitness  $\text{VO}_{2\text{max}}$  and serum 25(OH)D levels in 16- to 24-year-old healthy women. Similarly, Ardestani et al. [209] reported that there was evident a strong association between 25(OH)D levels and  $\text{VO}_{2\text{max}}$  in men and women (20–73 years old) over a broad range of 25(OH)D levels (10–82 ng/mL). Interestingly, this correlation was also evident in subjects with low levels of physical activity after adjusting for age, gender, BMI, and moderate to vigorous physical activity. Early studies have shown that exposed to sun athletes showed improved aerobic fitness compared to controls [28]. A recent study in professional soccer players reported that  $\text{VO}_{2\text{max}}$  was significantly related to vitamin D status [28]. Similar findings were reported in young physically active individuals [205]. The authors also reported that the individuals who had 25OHD levels above the recommended limit of 35 ng/mL, which was used as a cut-off value indicating vitamin D sufficiently in their study, were found to have significantly higher  $\text{VO}_{2\text{max}}$  values (20 %) than individuals below this cut-off point. On the contrary, it was reported that vitamin D status was not associated with neither the  $\text{VO}_{2\text{peak}}$ , as a measure of cardiorespiratory fitness, nor the end stage completed during a skating treadmill GXT in junior and collegiate male hockey players [206]. The authors suggested that these findings could be the result of the fact that none of their players were vitamin D deficient. According to Wilson et al. [210], worsening in aerobic capacity could result only as in vitamin D-deficient individuals. In the study by Fitzgerald et al. [206], the authors reported that although 37.7 % of their athletes had insufficient 25(OH)D levels, levels at the deficiency range are necessary for a compromise of cardiovascular fitness to ensue.

## 17. Vitamin D supplementation and athletes

Few published studies have reported that vitamin D supplementation benefits neuromuscular and aerobic performance capacity (Table 6). Indeed, a randomized placebo-controlled study examined the effects of vitamin D3 supplementation (5000 IU per day) for 8 weeks on musculoskeletal performance [211] in highly trained male professional soccer players. Approximately 62 % of the athletes on supplementation and 73 % of the controls exhibited serum total 25(OH)D < prior to treatment at 50 nmol/L. Vitamin D supplementation increased serum total 25(OH)D from a mean baseline of 29–103 nmol/L, whereas the placebo group showed no significant change. The authors found that vitamin D supplementation increased 10 m sprint time and vertical jump compared to the placebo group. Similar data were found in another study where athletes living at northerly latitudes (the United Kingdom = 53°N) had low vitamin D levels (<50 nmol/L). Additionally these data suggest that inadequate vitamin D levels are detrimental to musculoskeletal performance in athletes, while supplementation may increase both sprinting capacity and jumping ability of the soccer players. In support to these findings, Wyon et al. [212] reported enhanced neuromuscular performance in a different type of athletic population, i.e., in elite classical ballet dancers in a study which employed a 4-month oral supplementation of vitamin D3. Isometric muscular strength and vertical jump height

were measured pre- and post intervention. A significant increase of isometric strength (18.7 %) and vertical jump (7.1 %) was observed. The intervention group also sustained significantly less injuries than the controls during the study period. In agreement with these data are the findings of another study from the same group [213]. The authors examined the acute effects of vitamin D supplementation on muscle function using isokinetic dynamometry. White adult male national-level judoka athletes who were involved in full-time training participated in the study and were randomly allocated to the treatment (150,000 IU vitamin D<sub>3</sub>) or placebo and given blinded supplements by an independent researcher. Participants were tested twice, 8 days apart, on a Monday morning before the start of judo training and after 2 days of rest. A 5–7 mL of blood sample was collected followed by isokinetic concentric quadriceps and hamstring. The treatment group had a significant increase in serum 25(OH)D<sub>3</sub> levels (34 %) and muscle strength (13 %). However, other studies were unable to document any benefit following vitamin D supplementation in athletes with adequate or moderately deficient levels of vitamin D prior to supplementation. Close et al. [214] examined the effects of vitamin D<sub>3</sub> supplementation on serum 25(OH)D concentrations and on several exercise performance indices in club-level athletes. Participants were randomized into those receiving placebo, 20,000 or 40,000 IU per week of oral vitamin D<sub>3</sub> for a 12-week period. At baseline 57 % of the participants were found to be almost vitamin D deficient (mean 51 nmol/L). In the two supplementation groups, the following 6-week and 12-week period either with 20,000 IU vitamin D levels increased to 79 and 85 nmol/l, respectively, or with 40,000 IU vitamin D<sub>3</sub> to 98 and 91 nmol/l, respectively. Despite the different supplementation dosages, all individuals managed to reach levels greater than the suggested deficiency limit of 50 nmol/l. However, despite the observed increase in vitamin D serum levels, neither group exhibited an improvement of their exercise performance compared to controls [214]. The inability of the authors to document any beneficial effect of vitamin D on athletic performance may be due the fact that the participants of their study were not vitamin D deficient. It has been shown that the beneficial effects of vitamin D supplementation take place in individuals with significant vitamin D deficiency [158]. The lack of any beneficial effect of vitamin D supplementation was also reported in another study involving football players [215]. A placebo group and one receiving 5000 IU/day of vitamin D were examined. Both groups followed the same high-intensity training regime and had total work, sprint performance (5, 10, 20, 30 m), and jumping ability (squat jump and countermovement jump) measured. No significant differences were evident at baseline between the two groups for any of the measured parameters. The authors reported that although all examined performance indices, apart from the 30 m sprint value, were significantly higher as a result of the high-intensity interval training regime, the mean change scores (obtained values at the end of the study minus those of baseline) did not differ significantly between the two groups, an outcome suggesting that vitamin D supplementation did not had any beneficial effect. It should be noted here that the baseline vitamin D levels of the majority soccer players in this study were deficient. Therefore the proposed mechanism by Dahlquist et al. [158] was not applicable in this study. The authors suggest that other factors apart of vitamin D levels play a pivotal role including their initial training status. It is true though that the studies evaluating the effects of vitamin D supplementation on aerobic capacity in athletes are few and between. Jastrzebski et al. [216] have examined the effects of vitamin

D supplementation on  $VO_{2max}$ . The treatment group was composed of 14 elite lightweight rowers having sufficient 25(OH)D concentrations ( $>30$  ng/mL). They were put under vitamin D supplementation of 6000 IU per day for an 8-week period. The authors found a significant increase in  $VO_{2max}$  following vitamin D supplementation. Thus, it was concluded that 8 weeks of vitamin D supplementation during the training cycle of the same period of time resulted in enhanced aerobic metabolism in this type of athletes. However, since this is the only study that has examined vitamin D supplementation in athletes, these results cannot be generalized to all sporting disciplines. Furthermore, the low sample size of the study is another limiting factor regarding these findings.

Study	Quality	Sports	Participants	Intervention	Parameters measured	Results
Close et al. [211]	Placebo-controlled trial	Soccer	11 male soccer players, age = $18 \pm 5$ y	8 weeks, supplementation of 5000 IU/day of vitamin D3 or a cellulose placebo	1RM bench press and back squat, 10 m and 30m sprint, Illinois agility test, vertical jump	Significant $\uparrow$ in vertical jump height, 10m sprint times, and a trend for $\uparrow$ bench press and back squat 1RM
Wyon et al. [212]	Controlled trial	Ballet	24 elite male and female classical ballet dancers, age $28 \pm 3.85$ y	4-month supplementation of vitamin D3 (2000 IU per day)	Isometric muscular strength and vertical jump height	Significant $\uparrow$ in isometric muscular strength and vertical jump height
Wyon et al. [213]	Randomized controlled double-blind trial	Judoka	22 white male national-level judoka athletes. $27.5 \pm 4$ y	Acute effects, 1 week, one single dose of 150 000 IU vitamin D3 or placebo	Isokinetic concentric quadriceps, hamstring muscle strength	Significant $\uparrow$ in all measured strength-related indices
Close et al. [214]	Randomized controlled trial	Various sports	30 club-level athletes	Three groups receiving a placebo (PLB) either 20,000 or 40,000 IU/week of oral vitamin D3 for 12 weeks	RM bench press, leg press, vertical jump height	Neither dose given for 12 weeks improved the examined measures of physical performance
Jastrzebski, [216]	Randomized controlled trial	Rowing	14 elite lightweight rowers	8 weeks of 6000 IU/day of vitamin D3 in sufficient athletes ( $>30$ ng/mL)	Maximal oxygen consumption ( $VO_{2max}$ )	Significant $\uparrow$ in $VO_{2max}$
Jastrzebska et al. [215]	Controlled trial	Soccer	36 young soccer players, age: $17.5 \pm 0.6$ y	Treatment: around 5000 IU/day placebo— sunflower oil	Wingate test, 5, 10, 20, and 30 m sprint, squat jump (SJ), countermovement jump (CMJ)	No significant difference in any of the measured exercise performance parameters

y, years;  $\uparrow$ , increase

**Table 6.** Vitamin D supplementation and athletes.

## 18. Conclusion

This chapter describes the role of vitamin D in athletic performance. It describes the effect of vitamin D in several systems involved in exercise performance and physical activity in both athletes and non-athletes.

Vitamin D undoubtedly has a key role in the osseous health of both athletes and non-athletes. A growing body of evidence suggests that vitamin D is strongly related with the cardiovascular, immune, and neuromuscular systems. It is now clear that vitamin D plays an important role in the ability to perform efficiently during the normal daily activities, while its deficiency may result in poorer exercise performance. It should be mentioned that the vast majority of the evidence regarding vitamin D and exercise come from non-athletes. A growing number of studies have suggested that vitamin D exerts a beneficial effect on exercise performance, particularly in older adults and those with lower levels of circulating vitamin D. Regarding the optimum levels of vitamin D, the current evidence suggests that value levels above 30 ng/mL are needed. The majority of the vitamin D supplementation studies indicate that it may beneficially affect physical activity, frailty, muscular tone, stability, mobility in the elderly, and performance in physically active individuals or athletes only when pretreatment levels are within the insufficiency range (<30 ng/ml). Specifically for athletes, although the results of performance trials are not yet convincing enough to support vitamin D as a direct performance enhancer, obtaining optimal 25(OH)D levels can reduce the risk of debilitating musculoskeletal injury, and due to the active role in muscle, resolution of vitamin D insufficiency has the potential to impact performance.

## Author details

Nikolaos E. Koundourakis\* and Andrew N. Margioris

\*Address all correspondence to: [nkoyndoy@hotmail.com](mailto:nkoyndoy@hotmail.com)

Department of Clinical Chemistry, School of Medicine, University of Crete, Heraklion, Greece

## References

- [1] Holick MF: Vitamin D: a millennium perspective. *J Cell Biochem.* 2003;88:296–307. doi: 10.1002/jcb.10338
- [2] Muir SW, Montero-Odasso M: Effect of vitamin D supplementation on muscle strength, gait and balance in older adults: a systematic review and meta-analysis. *J Am Geriatr Soc.* 2011;59:2291–300.

- [3] Holick M: Sunlight and vitamin D. *J Gen Intern Med.* 2002;17:733–5. doi: 10.1046/j.1525-1497.2002.20731.x
- [4] Todd JJ, Pourshahidi LK, McSorley EM, Madigan SM, Magee PJ: Vitamin D: recent advances and implications for athletes. *Sports Med.* 2015;45:213–29. doi: 10.1007/s40279-014-0266-7.
- [5] Pojednic RM, Ceglia L: The emerging biomolecular role of vitamin D in skeletal muscle. *Exerc Sport Sci Rev.* 2014;42(2):76–81. doi: 10.1249/JES.0000000000000013.
- [6] Artaza JN, Mehrotra R, Norris KC: Vitamin D and the cardiovascular system. *Clin J Am Soc Nephrol.* 2009;4(9):1515–22. doi: 10.2215/CJN.02260409.
- [7] Moran DS, McClung JP, Kohen T, Lieberman HR: Vitamin D and physical performance. *Sports Med.* 2013;43(7):601–11. doi: 10.1007/s40279-013-0036-y.
- [8] Holick MF: Vitamin D deficiency. *N Engl J Med.* 2007;357(3):266–81. doi: 10.1056/NEJMra070553
- [9] Hamilton B: Vitamin D and human skeletal muscle. *Scand J Med Sci Sports.* 2010;20:182–90. doi: 10.1111/j.1600-0838.2009.01016.x
- [10] Cashman KD: Calcium and vitamin D. *Novartis Found Symp.* 2007;282:123–38.
- [11] Tavera-Mendoza LE, White JH. Cell defenses and the sunshine vitamin. *Sci Am.* 2007;297(5):62–5, 68–70, 72.
- [12] Udowenko M, Trojan T: Vitamin D: extent of deficiency, effect on muscle function, bone health, performance, and injury prevention. *Conn Med.* 2010;74(8):477–80.
- [13] Baker AR, McDonnell DP, Hughes M, Crisp TM, Mangelsdorf DJ, Haussler MR, Pike JW, Shine J, O'Malley W: Cloning and expression of full-length cDNA encoding human vitamin D receptor. *Proc Natl Acad Sci U S A.* 1988;85:3294–8.
- [14] Brumbaugh PF, Haussler DH, Bressler R, Haussler MR: Radioreceptor assay for 1 alpha, 25-dihydroxyvitamin D<sub>3</sub>. *Science.* 1974;183:1089–91.
- [15] Brumbaugh PF, Haussler MR: Specific binding of 1 $\alpha$ ,25-dihydroxycholecalciferol to nuclear components of chick intestine. *J Biol Chem.* 1975;250:1588–94.
- [16] Saccone D, Asani F, Bornman L: Regulation of the vitamin D receptor gene by environment, genetics and epigenetics. *Gene.* 2015;561(2):171–80. doi: 10.1016/j.gene.2015.02.024.
- [17] Christakos S, Hewison M, Gardner DG, Wagner CL, Sergeev IN, Rutten E, Pittas AG, Boland R, Ferrucci L, Bikle DD: Vitamin D: beyond bone. *Ann N Y Acad Sci.* 2013;1287:45–58. doi: 10.1111/nyas.12129.
- [18] Soni M, Kos K, Lang IA, Jones K, Melzer D, Llewellyn DJ: Vitamin D and cognitive function. *Scand J Clin Lab Invest Suppl.* 2012;243:79–82. doi: 10.3109/00365513.2012.681969.



- [19] Halliday TM, Peterson NJ, Thomas JJ, Kleppinger K, Hollis BW, Larson-Meyer DE: Vitamin D status relative to diet, lifestyle, injury, and illness in college athletes. *Med Sci Sports Exerc.* 2011;43(2):335–43. doi: 10.1249/MSS.0b013e3181eb9d4d.
- [20] Holick MF: Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr.* 2004;80 (6 suppl):1678S–88S.
- [21] Meyer C. Scientists probe role of vitamin D: deficiency a significant problem, experts say. *JAMA.* 2004;292(12):1416–8.
- [22] Zittermann A. Vitamin D in preventive medicine: are we ignoring the evidence? *Br J Nutr.* 2003;89(5):552–72.
- [23] Farrokhyar F, Tabasinejad R, Dao D, Peterson D, Ayeni OR, Hadioonzadeh R, Bhandari M: Prevalence of vitamin D inadequacy in athletes: a systematic-review and meta-analysis. *Sports Med.* 2015;45(3):365–78. doi: 10.1007/s40279-014-0267-6.
- [24] Wacker M, Holick MF: Vitamin D—effects on skeletal and extraskeletal health and the need for supplementation. *Nutrients.* 2013;5(1):111-48. doi: 10.3390/nu5010111.
- [25] Golden NH, Abrams SA: Committee on Nutrition. Optimizing bone health in children and adolescents. *Pediatrics.* 2014;134(4):e1229–43. doi: 10.1542/peds.2014-2173.
- [26] Morton JP, Iqbal Z, Drust B, Burgess D, Close GL, Brukner PD: Seasonal variation in vitamin D status in professional soccer players of the English Premier League. *Appl Physiol Nutr Metab.* 2012;37:798–802. doi: 10.1139/h2012-037.
- [27] Larson-Meyer DE, Willis KS: Vitamin D and athletes. *Curr Sports Med Rep.* 2010;9(4): 220–6. doi: 10.1249/JSR.0b013e3181e7dd45.
- [28] Koundourakis NE, Androulakis NE, Malliaraki N, Margioris AN: Vitamin D and exercise performance in professional soccer players. *PLoS ONE.* 2014;9(7):e101659. doi: 10.1371/journal.pone.0101659.
- [29] Angeline ME, Gee AO, Shindle M, Warren RF, Rodeo SA: The effects of vitamin D deficiency in athletes. *Am J Sports Med.* 2013;41:461–4. doi: 10.1177/0363546513475787.
- [30] Lovell G: Vitamin D status of females in an elite gymnastics program. *Clin J Sport Med.* 2008;18(2):159–61. doi: 10.1097/JSM.0b013e3181650eee.
- [31] Thacher TD, Clarke BL: Vitamin D insufficiency. *Mayo Clin Proc.* 2011;86:50–60. doi: 10.4065/mcp.2010.0567.
- [32] Kopeć A, Solarz K, Majda F, Słowińska-Lisowska M, Mędraś M: An evaluation of the levels of vitamin D and bone turnover markers after the summer and winter periods in polish professional soccer players. *J Hum Kinet.* 2013;38:135–40. doi: 10.2478/hukin-2013-0053.

- [33] Willis KS, Smith DT, Broughton KS, Larson-Meyer DE. Vitamin D status and biomarkers of inflammation in runners. *Open Access J Sports Med.* 2012;3:35–42. doi: 10.2147/OAJSM.S31022.
- [34] Halliday TM, Peterson NJ, Thomas JJ, Kleppinger K, Hollis BW, Larson-Meyer DE: Vitamin D status relative to diet, lifestyle, injury, and illness in college athletes. *Med Sci Sports Exerc.* 2011;43(2):335–43. doi: 10.1249/MSS.0b013e3181eb9d4d.
- [35] Constantini NW, Arieli R, Chodick G, Dubnov-Raz G: High prevalence of vitamin D insufficiency in athletes and dancers. *Clin J Sport Med.* 2010;20(5):368–71. doi: 10.1097/JSM.0b013e3181f207f2.
- [36] Elsammak MY, Al-Wosaibi AA, Al-Howeish A, Alsaeed J: Vitamin d deficiency in Saudi Arabs. *Horm Metab Res.* 2010;42:364–8. doi: 10.1055/s-0030-1248296.
- [37] Babu US, Calvo MS: Modern India and the vitamin D dilemma: evidence for the need of a national food fortification program. *Mol Nutr Food Res.* 2010;54:1134–47. doi: 10.1002/mnfr.200900480.
- [38] Maxwell JD. Seasonal variation in vitamin D. *Proc Nutr Soc* 1994;53:533–43.
- [39] Storlie DM, Pritchett K, Pritchett R, Cashman L. 12-Week vitamin D supplementation trial does not significantly influence seasonal 25(OH)D status in male collegiate athletes. *Int J Health Nutr.* 2011;2:8–13.
- [40] Galan F, Ribas J, Sánchez-Martinez PM, Calero T, Sánchez AB, Muñoz A. Serum 25-hydroxyvitamin D in early autumn to ensure vitamin D sufficiency in mid-winter in professional football players. *Clin Nutr.* 2012;31:132–6. doi: 10.1016/j.clnu.2011.07.008.
- [41] Cannell JJ, Hollis BW, Sorenson MB, Taft TN, Anderson JJ. Athletic performance and vitamin D. *Med Sci Sport Exer.* 2009;41:1102–10.
- [42] Koch H, Raschka C: Circannual period of physical performance analysed by means of standard cosinor analysis: a case report. *Rom J Physiol.* 2000;37(1–4):51–8.
- [43] Erikssen J, Rodahl K. Seasonal variation in work performance and heart rate response to exercise. A study of 1,835 middle-aged men. *Eur J Appl Physiol Occup Physiol.* 1979;42(2):133–40.
- [44] Svedenhag J, Sjodin B. Physiological characteristics of elite male runners in and off-season. *Can J Appl Sport Sci.* 1985;10(3):127–33.
- [45] Mozos I, Marginean O: Links between Vitamin D deficiency and cardiovascular diseases. *Biomed Res Int.* 2015;2015:109275. doi: 10.1155/2015/109275.
- [46] Bonjour JP: Calcium and phosphate: a duet of ions playing for bone health. *Am Coll Nutr.* 2011;30(5 Suppl 1):438S–48S
- [47] Holick, MF: Vitamin D deficiency. *N Engl J Med.* 2007, 357, 266–81.

- [48] Christakos S, Dhawan P, Porta A, Mady LJ, Seth T: Vitamin D and intestinal calcium absorption. *Mol Cell Endocrinol.* 2011;347:25–29. doi: 10.1016/j.mce.2011.05.038.
- [49] Kulie T, Groff A, Redmer J, Hounshell J, Schragger S: Vitamin D: an evidence-based review. *J Am Board Fam Med.* 2009;22(6):698–706. doi: 10.3122/jabfm.2009.06.090037.
- [50] Kota S, Jammula S, Kota S, Meher L, Modi K: Correlation of vitamin D, bone mineral density and parathyroid hormone levels in adults with low bone density. *Indian J Orthop.* 2013;47(4):402–7. doi: 10.4103/0019-5413.114932.
- [51] Ethgen O, Hiligsmann M, Burlet N, Reginster JY: Public health impact and cost-effectiveness of dairy products supplemented with vitamin D in prevention of osteoporotic fractures. *Arch Public Health.* 2015;14:48. doi: 10.1186/s13690-015-0099-3.
- [52] Buchebner D, McGuigan F, Gerdhem P, Ridderstråle M, Akesson K: Association between hypovitaminosis D in elderly women and long- and short-term mortality-results from the osteoporotic prospective risk assessment cohort. *J Am Geriatr Soc.* 2016;64:990–7. doi: 10.1111/jgs.14087.
- [53] Sundh D, Mellström D, Ljunggren Ö, Karlsson MK, Ohlsson C, Nilsson M, Nilsson AG, Lorentzon M: Low serum vitamin D is associated with higher cortical porosity in elderly men. *J Intern Med.* 2016;15. doi: 10.1111/joim.12514. [Epub ahead of print].
- [54] Scino Y, Ishizuka S, Shima M, Tanaka H: Vitamin D in bone formation. *Osteoporosis Int.* 1993;Suppl. 1:S196–198.
- [55] Jones BH, Thacker SB, Gilchrist J, Kimsey CD Jr, Sosin DM: Prevention of lower extremity stress fractures in athletes and soldiers: a systematic review. *Epidemiol Rev.* 2002;24:228–47.
- [56] McClung JP, Karl JP: Vitamin D and stress fracture: the contribution of vitamin D receptor gene polymorphisms. *Nutr Rev.* 2010;68(6):365–9. doi: 10.1111/j.1753-4887.2010.00295.x.
- [57] Friedl KE, Evans RK, Moran DS: Stress fracture and military medical readiness: bridging basic and applied research. *Med Sci Sports Exerc.* 2008;40 (11 Suppl):S609–22. doi: 10.1249/MSS.0b013e3181892d53.
- [58] Willis KS: Vitamin D. Status and immune system biomarkers in athletes. Laramie: University of Wyoming; 2008. p. 85.
- [59] Prineas JW, Mason AS, Henson RA: Myopathy in metabolic bone disease. *Br Med J.* 1965;1:1034–6.
- [60] Wang Y, Zhu J, DeLuca HF: Where is the vitamin D receptor? *Arch Biochem Biophys.* 2012;523:123–33. doi: 10.1016/j.abb.2012.04.001.
- [61] Agergaard J, Trøstrup J, Uth J, Iversen JV, Boesen A, Andersen JL, Schjerling P, Langberg H: Does vitamin-D intake during resistance training improve the skeletal muscle

- hypertrophic and strength response in young and elderly men?—a randomized controlled trial. *Nutr Metab (Lond)*. 2015;12:32. doi: 10.1186/s12986-015-0029-y.
- [62] Garay E, Jankowski P, Lizano P, Marczak S, Garay E, Jankowski P, Lizano P, Marczak S, Maehr H, Adorini L, Uskokovic MR, Studzinski GP: Calcitriol derivatives with two different side-chains at C-20. Part 4: further chain modifications that alter VDR-dependent monocytic differentiation potency in human leukemia cells. *Bioorg Med Chem*. 2007;15(13):4444–55.
- [63] Garcia LA, King KK, Ferrini MG, Norris KC, Artaza JN: 1,25(OH)<sub>2</sub>vitamin D<sub>3</sub> stimulates myogenic differentiation by inhibiting cell proliferation and modulating the expression of promyogenic growth factors and myostatin in C2C12 skeletal muscle cells. *Endocrinology*. 2011;152(8):2976–86. doi: 10.1210/en.2011-0159.
- [64] Lee SJ, Lee YS, Zimmers TA, Soleimani A, Matzuk MM, Tsuchida K, Cohn RD, Barton ER: Regulation of muscle mass by follistatin and activins. *Mol Endocrinol*. 2010;24:1998–2008. doi: 10.1210/me.2010-0127.
- [65] Angeline ME, Gee AO, Shindle M, Warren RF, Rodeo SA: The effects of vitamin D deficiency in athletes. *Am J Sports Med*. 2013;41:461–4. doi: 10.1177/0363546513475787.
- [66] Black PN and, Scragg R: Relationship between serum 25-hydroxyvitamin D and pulmonary function in the third national health and nutrition examination survey. *Chest*. 2005;128:3792–8.
- [67] Deschenes MR, Kraemer WJ: Performance and physiologic adaptations to resistance training. *Am J Phys Med Rehabil*. 2002;81(11), S3–S16.
- [68] Shuler FD, Wingate MK, Moore GH, Giangarra C: Sports health benefits of vitamin d. *Sports Health*. 2012;4(6):496–501. doi: 10.1177/1941738112461621.
- [69] Beavers KM, Beavers DP, Houston DK, Harris TB, Hue TF, Koster A, Newman AB, Simonsick EM, Studenski SA, Nicklas BJ, Kritchevsky SB: Associations between body composition and gait-speed decline: results from the Health, Aging, and Body Composition study. *Am J Clin Nutr*. 2013;97(3):552–60. doi: 10.3945/ajcn.112.047860.
- [70] Birge SJ, Haddad JG: 25-hydroxycholecalciferol stimulation of muscle metabolism. *J Clin Invest*. 1975;56(5):1100–7.
- [71] Oh J, Weng S, Felton SK, Bhandare S, Riek A, Butler B, Proctor BM, Petty M, Chen Z, Schechtman KB, Bernal-Mizrachi L, Bernal-Mizrachi C: 1,25(OH)<sub>2</sub> vitamin d inhibits foam cell formation and suppresses macrophage cholesterol uptake in patients with type 2 diabetes mellitus. *Circulation*. 2009;120(8):687–98. doi: 10.1161/CIRCULATIONAHA.109.856070.
- [72] Rigby WF, Denome S, Fanger MW: Regulation of lymphokine production and human T lymphocyte activation by 1,25-dihydroxyvitamin D<sub>3</sub>. Specific inhibition at the level of messenger RNA. *J Clin Invest*. 1987;79(6):1659–64.

- [73] Simpson RU, Hershey SH, Nibelink KA: Characterization of heart size and blood pressure in the vitamin D receptor knockout mouse. *J Steroid Biochem Mol Biol.* 2007;103:521–4.
- [74] Saleh FN, Schirmer H, Sundsfjord J, Jorde E: Parathyroid hormone and left ventricular hypertrophy. *Eur Heart J.* 2003;24:2054–60.
- [75] Al Mheid I, Patel R, Murrow J, Morris A, Rahman A, Fike L, Nino Al Mheid I, Patel R, Murrow J, Morris A, Rahman A, Fike L, Kavtaradze N, Uphoff I, Hooper C, Tangpricha V, Alexander RW, Brigham K, Quyyumi AA. Vitamin D status is associated with arterial stiffness and vascular dysfunction in healthy humans. *J Am Coll Cardiol.* 2011;58(2): 186–92. doi 10.1016/j.jacc.2011.02.051.
- [76] Tomaschitz A, Pilz S, Ritz E, Grammer T, Drechsler C, Boehm BO, März W: Independent association between 1,25-dihydroxyvitamin D, 25-hydroxyvitamin D and the renin-angiotensin system: The Ludwigshafen Risk and Cardiovascular Health (LURIC) study. *Clin Chim Acta.* 2010;411:1354–60. doi: 10.1016/j.cca.2010.05.037.
- [77] Wang TJ, Pencina MJ, Booth SL, Jacques PF, Ingelsson E, Lanier K, Benjamin EJ, D'Agostino RB, Wolf M, Vasani RS: Vitamin D deficiency and risk of cardiovascular disease. *Circulation.* 2008;117(4):503–11. doi: 10.1161/CIRCULATIONAHA.107.706127.
- [78] Lee JH, O'Keefe JH, Bell D, Hensrud DD, Holick MF: Vitamin D deficiency an important, common, and easily treatable cardiovascular risk factor? *J Am Coll Cardiol.* 2008;52(24): 1949–56. doi: 10.1016/j.jacc.2008.08.050.
- [79] McDermott MM, Liu K, Ferrucci L, Tian L, Guralnik J, Kopp P, Van Horn L, Liao Y, Green D, Kibbe M, Sufit R, Zhao L, Criqui MH. Vitamin D status, functional decline, and mortality in peripheral artery disease. *Vasc Med.* 2014;19(1):18–26. doi: 10.1177/1358863X13518364.
- [80] Al Mheid I, Patel RS, Tangpricha V, Quyyumi AA: Vitamin D and cardiovascular disease: is the evidence solid? *Eur Heart J.* 2013;34(48):3691–8. doi: 10.1093/eurheartj/eh166.
- [81] Drechsler C, Pilz S, Obermayer-Pietsch B, Verduijn M, Tomaschitz A, Krane V, Espe K, Dekker F, Brandenburg V, März W, Ritz E, Wanner C: Vitamin D deficiency is associated with sudden cardiac death, combined cardiovascular events, and mortality in haemodialysis patients. *Eur Heart J.* 2010;31:2253–61. doi: 10.1093/eurheartj/ehq246.
- [82] Deo R, Katz R, Shlipak MG, Sotoodehnia N, Psaty BM, Sarnak MJ, Fried LF, Chonchol M, de Boer IH, Enquobahrie D, Siscovick D, Kestenbaum B: Vitamin D, parathyroid hormone, and sudden cardiac death: results from the Cardiovascular Health Study. *Hypertension.* 2011;58:1021–8. doi: 10.1161/HYPERTENSIONAHA.111.179135.
- [83] Naesgaard PA, León De La Fuente RA, Nilsen ST, Woie L, Aarsland T, Brede C, Staines H, Nilsen DW: Serum 25(OH)D is a 2-year predictor of all-cause mortality, cardiac death

- and sudden cardiac death in chest pain patients from Northern Argentina. *PLoS One*. 2012;7:e43228. doi: 10.1371/journal.pone.0043228.
- [84] Artaza JN, Mehrotra R, Norris KC: Vitamin D and the cardiovascular system. *Clin J Am Soc Nephrol*. 2009;4:1515–22. doi: 10.2215/CJN.02260409.
- [85] Bajaj A, Stone KL, Peters K, Parimi N, Barrett-Connor E, Bauer D, Cawthon PM, Ensrud KE, Hoffman AR, Orwoll E, Schernhammer ES: Circulating vitamin D, supplement use, and cardiovascular disease risk: the MrOS Sleep Study. *J Clin Endocrinol Metab*. 2014;99:3256–62. doi: 10.1210/jc.2013-4178.
- [86] Martins D, Wolf M, Pan D, Zadshir A, Tareen N, Thadhani R, Felsenfeld A, Levine B, Mehrotra R, Norris K: Prevalence of cardiovascular risk factors and the serum levels of 25-hydroxyvitamin D in the United States: data from the Third National Health and Nutrition Examination Survey. *Arch Intern Med*. 2007;167:1159–65.
- [87] Gouni-Berthold I, Krone W, Berthold HK: Vitamin D and cardiovascular disease. *Curr Vasc Pharmacol*. 2009;7:414–22.
- [88] Mpandzou G, Aït Ben Haddou E, Regragui W, Benomar A, Yahyaoui M: Vitamin D deficiency and its role in neurological conditions: a review. *Rev Neurol (Paris)*. 2016;172(2):109–22. doi: 10.1016/j.neurol.2015.11.005.
- [89] Wrzosek M, Łukaszkiwicz J, Wrzosek M, Jakubczyk A, Matsumoto H, Piątkiewicz P, Radziwoń-Zaleska M, Wojnar M, Nowicka G: Vitamin D and the central nervous system. *Pharmacol Rep*. 2013;65(2):271–8.
- [90] DeLuca GC, Kimball SM, Kolasinski J, Ramagopalan SV, Ebers GC: Review: the role of vitamin D in nervous system health and disease. *Neuropathol Appl Neurobiol*. 2013;39(5):458–84. doi: 10.1111/nan.12020.
- [91] Mpandzou G, Aït Ben Haddou E, Regragui W, Benomar A, Yahyaoui M: Vitamin D deficiency and its role in neurological conditions: a review. *Rev Neurol (Paris)*. 2016;172:109–22. doi: 10.1016/j.neurol.2015.11.005.
- [92] Buell JS, Dawson-Hughes B: Vitamin D and neurocognitive dysfunction: preventing “D” ecline? *Mol Aspects Med*. 2008;29:415–22. doi: 10.1016/j.mam.2008.05.001.
- [93] Zehnder D, Bland R, Williams MC, McNinch RW, Howie AJ, Stewart PM, Hewison M: Extrarenal expression of 25-hydroxyvitamin d(3)-1 alpha-hydroxylase. *J Clin Endocrinol Metab*. 2001;86:888–94.
- [94] Cai Q, Tapper DN, Gilmour RF Jr, deTalamoni N, Aloia RC, Wasserman RH: Modulation of the excitability of avian peripheral nerves by vitamin D: relation to calbindin-D28k, calcium status and lipid composition. *Cell Calcium*. 1994;15:401–10.
- [95] Cantorna MT, Hayes CE, and DeLuca HF: 1,25-Dihydroxyvitamin D3 reversibly blocks the progression of relapsing encephalomyelitis, a model of multiple sclerosis. *Proc Natl Acad Sci USA*. 1996;93:7861–64.

- [96] Radak Z, Taylor AW, Ohno H, Goto S: Adaptation to exercise induced oxidative stress: from muscle to brain. *Exerc Immunol Rev.* 2001;7:90–107.
- [97] Pirotta S, Kidgell DJ, Daly RM: Effects of vitamin D supplementation on neuroplasticity in older adults: a double-blinded, placebo-controlled randomised trial. *Osteoporos Int.* 2015;26(1):131–40. doi: 10.1007/s00198-014-2855-6.
- [98] Taghizadeh M, Talaei SA, Djazayeri A, Salami M : Vitamin D supplementation restores suppressed synaptic plasticity in Alzheimer's disease. *Nutr Neurosci.* 2014;17(4):172–7. doi: 10.1179/1476830513Y.0000000080.
- [99] Nutt D: GABA-A receptors: subtypes, regional distribution, and function. *J Clin Sleep Med.* 2006;2:S7-S11.
- [100] Manini TM, Clark BC: Dynapenia and aging: an update. *J Gerontol Ser A: Biol Sci Med Sci.* 2012;67A(1):28–40. doi: 10.1093/gerona/qlr010.
- [101] Acworth I, Nicholass J, Morgan B, Newsholme EA: Effect of sustained exercise on concentrations of plasma aromatic and branched-chain amino acids and brain amines. *Biochem Biophys Res Commun.* 1986;137(1):149–53.
- [102] Meeusen R, Watson P, Hasegawa H, Roelands B, Piacentini MF: Central fatigue: the serotonin hypothesis and beyond. *Sports Med.* 2006;36(10):881–909.
- [103] Tague SE, Smith PG: Vitamin D receptor and enzyme expression in dorsal root ganglia of adult female rats: modulation by ovarian hormones. *J Chem Neuroanat.* 2011;41(1):1–12. doi: 10.1016/j.jchemneu.2010.10.001.
- [104] Dubin AE, Patapoutian A: Nociceptors: the sensors of the pain pathway. *J Clin Invest.* 2010;120(11):3760–72. doi: 10.1172/JCI42843.
- [105] Dhesi JK, Jackson SH, Bearne LM, Moniz C, Hurley MV, Swift CG, Allain TJ: Vitamin D supplementation improves neuromuscular function in older people who fall. *Age Ageing.* 2004;33(6):589–95.
- [106] Lord SR, Clark RD, Webster IW: Physiological factors associated with falls in an elderly population. *J Am Geriatr Soc.* 1991;39:1194–200.
- [107] Shi YY, Liu TJ, Fu JH, Xu W, Wu LL, Hou AN, Xue XD: Vitamin D/VDR signaling attenuates lipopolysaccharide-induced acute lung injury by maintaining the integrity of the pulmonary epithelial barrier. *Mol Med Rep.* 2016;13(2):1186–94. doi: 10.3892/mmr.2015.4685.
- [108] Rehan VK, Torday JS. Vitamin D and Lung Development in Early Life. In: Litonjua AA, editor. *Vitamin D and the Lung: Mechanisms and Disease Associations.* New York; Humana Press London. P. 46. doi: 10.1007/978-1-61779-888-7.
- [109] Rehan VK, Torday JS, Peleg S, Gennaro L, Vouros P, Padbury J, Rao DS, Reddy GS: 1 $\alpha$ ,25-dihydroxy-3-epi-vitamin D<sub>3</sub>, a natural metabolite of 1 $\alpha$ ,25-dihydroxy



- vitamin D3: production and biological activity studies in pulmonary alveolar type II cells. *Mol Genet Metab.* 2002;76(1):46–56.
- [110] Vedala SR, Paul N, Mane AB: Difference in pulmonary function test among the athletic and sedentary population. *Natl J Physiol Pharm Pharmacol.* 2013;3(2):118–23. doi: 10.5455/njppp.2013.3.109-114.
- [111] Nguyen M, Trubert CL, Rizk-Rabin M, Rehan VK, Besancon F, Cayre YE, Garabedian M: 1,25-Dihydroxyvitamin D3 and fetal lung maturation: immunogold detection of VDR expression in pneumocytes type II cells and effect on fructose 1,6 bisphosphatase. *J Steroid Biochem Mol Biol.* 2004;89:93–97.
- [112] Janssens W, Bouillon R, Claes B, Carremans C, Lehouck A, Buysschaert I, Coolen J, Mathieu C, Decramer M, Lambrechts D: Vitamin D deficiency is highly prevalent in COPD and correlates with variants in the vitamin D-binding gene. *Thorax.* 2010;65:215–20. doi: 10.1136/thx.2009.120659.
- [113] Damera G, Fogle HW, Lim P, Goncharova EA, Zhao H, Banerjee A, Tliba O, Krymskaya VP, Panettieri RA: Vitamin D inhibits growth of human airway smooth muscle cells through growth factor-induced phosphorylation of retinoblastoma protein and checkpoint kinase 1. *Br J Pharmacol.* 2009;158:1429–41. doi: 10.1111/j.1476-5381.2009.00428.x.
- [114] Zosky GR, Berry LJ, Elliot JG, James AL, Gorman S, Hart PH: Vitamin D deficiency causes deficits in lung function and alters lung structure. *Am J Respir Crit Care Med.* 2011;183(10):1336–43. doi: 10.1164/rccm.201010-1596OC.
- [115] Litonju A, Weiss S: Is vitamin D deficiency to blame for the asthma epidemic? *J Allergy Clin Immunol.* 2007;120:1031–5.
- [116] Trochoutsou AI, Kloukina V, Samitas K, Xanthou G: Vitamin-D in the immune system: genomic and non-genomic actions. *Mini Rev Med Chem.* 2015;15(11):953–63.
- [117] Aranow C: Vitamin D and the Immune System. *J Invest Med.* 2011;59(6):881–6. doi: 10.231/JIM.0b013e31821b8755.
- [118] Prietl B, Treiber G, Pieber TR, Amrein K: Vitamin D and immune function. *Nutrients.* 2013;5(7):2502–21. doi: 10.3390/nu5072502.
- [119] Wang TT, Nestel FP, Bourdeau V, Nagai Y, Wang Q, Liao J, Tavera-Mendoza L, Lin R, Hanrahan JW, Mader S, White JH: Cutting edge: 1,25-dihydroxyvitamin D3 is a direct inducer of antimicrobial peptide gene expression. *J Immunol.* 2004;173(5):2909–12.
- [120] Ramanathan B, Davis EG, Ross CR, Blecha F: Cathelicidins: microbicidal activity, mechanisms of action, and roles in innate immunity. *Microbes Infect.* 2002;4(3):361–72.
- [121] Provvedini DM, Tsoukas CD, Deftos LJ, Manolagas SC: 1,25-Dihydroxyvitamin D3 receptors in human leukocytes. *Science.* 1983;221:1181–3.

- [122] Lemire JM, Adams JS, Sakai R, Jordan SC: 1 $\alpha$ ,25-dihydroxyvitamin D3 suppresses proliferation and immunoglobulin production by normal human peripheral blood mononuclear cells. *J Clin Investig.* 1984;74:657–61.
- [123] Chen S, Sims GP, Chen XX, Gu YY, Chen S, Lipsky PE: Modulatory effects of 1,25-dihydroxyvitamin D3 on human B cell differentiation. *J Immunol.* 2007;179:1634–47.
- [124] Mahon BD, Wittke A, Weaver V, Cantorna MT: The targets of vitamin D depend on the differentiation and activation status of CD4 positive T cells. *J Cell Biochem.* 2003;89:922–32.
- [125] Hewison M: Vitamin D and immune function: an overview. *Proc Nutr Soc.* 2012;71(1):50–61.
- [126] Boonstra A, Barrat FJ, Crain C, Heath VL, Savelkoul HF, O'Garra A: 1 $\alpha$ ,25-Dihydroxyvitamin d3 has a direct effect on naive CD4(+) T cells to enhance the development of Th2 cells. *J Immunol.* 2001;167(9):4974–80.
- [127] Berry DJ, Hesketh K, Power C, Hyppönen E: Vitamin D status has a linear association with seasonal infections and lung function in British adults. *Br J Nutr.* 2011;106:1433–40. doi: 10.1017/S0007114511001991.
- [128] Ginde AA, Mansbach JM, and Camargo CA Jr: Association between serum 25-hydroxyvitamin D level and upper respiratory tract infection in the Third National Health and Nutrition Examination Survey. *Arch Intern Med.* 2009;169:384–90. doi: 10.1001/archinternmed.2008.560.
- [129] Laaksi I, Ruohola JP, Tuohimaa P, Auvinen A, Haataja R, Pihlajamäki H, and Ylikomi T: An association of serum vitamin D concentrations < 40 nmol/L with acute respiratory tract infection in young Finnish men. *Am J Clin Nutr.* 2007;86(3):714–7.
- [130] Gleeson M (Editor). *Immune function in sport and exercise.* Edinburgh: Elsevier, 2005.
- [131] Li TL, Gleeson M: The effect of single and repeated bouts of prolonged cycling and circadian variation on saliva flow rate, immunoglobulin A and alpha-amylase responses. *J Sports Sci.* 2004;22(11–12):1015–24.
- [132] de Sa Del Fiol F, Barberato-Filho S, Lopes LC, de Cassia Bergamaschi C: Vitamin D and respiratory infections. *J Infect Dev Ctries.* 2015;9(4):355–61. doi: 10.3855/jidc.5711.
- [133] Bryson KJ, Nash AA, Norval M: Does vitamin D protect against respiratory viral infections? *Epidemiol Infect.* 2014;142(9):1789–801. doi: 10.1017/S0950268814000193.
- [134] Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, Ochoa MT, Schaubert J, Wu K, Meinken C, Kamen DL, Wagner M, Bals R, Steinmeyer A, Zügel U, Gallo RL, Eisenberg D, Hewison M, Hollis BW, Adams JS, Bloom BR, Modlin RL: Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science.* 2006;311:1770–3.
- [135] Larson-Meyer DE, Willis KS: Vitamin D and athletes. *Curr Sports Med Rep.* 2010;9(4):220–6. doi: 10.1249/JSR.0b013e3181e7dd45.

- [136] Villalta SA, Rinaldi C, Deng B, Liu G, Fedor B, Tidball JG: Interleukin-10 reduces the pathology of mdx muscular dystrophy by deactivating M1 macrophages and modulating macrophage phenotype. *Hum Mol Genet.* 2011;20:790–805. doi: 10.1093/hmg/ddq523.
- [137] Takahashi Y, Katayose D, Shindoh C: Interleukin-13 prevents diaphragm muscle deterioration in a septic animal model. *Tohoku J Exp Med.* 1999;189:191–202.
- [138] He C, Handzlik M, Fraser WD, Muhamad A, Preston H, Richardson A, Gleeson M: Influence of vitamin D status on respiratory infection incidence and immune function during 4 months of winter training in endurance sport athletes. *Exerc Immunol Rev.* 2013;19:86–101
- [139] Li J, Byrne ME, Chang E, Jiang Y, Donkin SS, Buhman KK, Burgess JR, Teegarden D:  $1\alpha,25$ -dihydroxyvitamin D hydroxylase in adipocytes. *J Steroid Biochem Mol Biol.* 2008;112(1–3):122–6. doi: 10.1016/j.jsbmb.2008.09.006.
- [140] Koszowska AU, Nowak J, Dittfeld A, Brończyk-Puzoń A, Kulpok A, Zubelewicz-Szkodzińska B: Obesity, adipose tissue function and the role of vitamin D. *Cent Eur J Immunol.* 2014;39(2):260–4. doi: 10.5114/ceji.2014.43732.
- [141] Pannu PK, Zhao Y, Soares MJ : Reductions in body weight and percent fat mass increase the vitamin D status of obese subjects: a systematic review and meta-regression analysis. *Nutr Res.* 2016;36(3):201–13. doi: 10.1016/j.nutres.2015.11.013.
- [142] Rosenblum JL, Castro VM, Moore CE, Kaplan LM: Calcium and vitamin D supplementation is associated with decreased abdominal visceral adipose tissue in overweight and obese adults. *Am J Clin Nutr.* 2012;95(1):101–8. doi: 10.3945/ajcn.111.019489.
- [143] Drincic AT, Armas LA, Van Diest EE, Heaney RP: Volumetric dilution, rather than sequestration best explains the low vitamin D status of obesity. *Obesity (Silver Spring).* 2012;20:1444–48. doi: 10.1038/oby.2011.404.
- [144] Earthman CP, Beckman LM, Masodkar K, Sibley SD: The link between obesity and low circulating 25-hydroxyvitamin D concentrations: considerations and implications. *Int J Obes.* 2012;36(3):387–96. doi: 10.1038/ijo.2011.119.
- [145] Datta S, Pal M, De A. The dependency of vitamin D status on anthropometric data. *Malay J Med Sci.* 2014;21(3):54–61.
- [146] Forney LA, Earnest CP, Henagan TM, Johnson LE, Castleberry TJ, Stewart LK: Vitamin D status, body composition, and fitness measures in college-aged students. *J Strength Cond Res.* 2014;28(3):814–24. doi: 10.1519/JSC.0b013e3182a35ed0.
- [147] Koundourakis NE, Androulakis NE, Malliaraki N, Tsatsanis C, Venihaki M, Margioris AN: Discrepancy between exercise performance, body composition, and sex steroid response after a six-week detraining period in professional soccer players. *PLoS ONE.* 2014;9(2):e87803. doi: 10.1371/journal.pone.0087803.

- [148] Lira FS, Rosa JC, Cunha CA, Ribeiro EB, do Nascimento CO, Oyama LM, Mota JF: Supplementing alpha-tocopherol (vitamin E) and vitamin D3 in high fat diet decrease IL-6 production in murine epididymal adipose tissue and 3T3-L1 adipocytes following LPS stimulation. *Lipids Health Dis.* 2011;10:37. doi: 10.1186/1476-511X-10-37.
- [149] Consiglio M, Destefanis M, Morena D, Foglizzo V, Forneris M, Pescarmona G, Silvagno F: The vitamin D receptor inhibits the respiratory chain, contributing to the metabolic switch that is essential for cancer cell proliferation. *PLoS One.* 2014;9:e115816. doi: 10.1371/journal.pone.0115816.
- [150] Allen R, Cureton T: Effects of ultraviolet radiation on physical fitness. *Arch Phys Med.* 1945;10:641–4.
- [151] Gorkin Z, Gorkin MJ, Teslenko NE: [The effect of ultraviolet irradiation upon training for 100m sprint.] *Fiziol Zh USSR.* 1938;25:695–701. [Article in Russian].
- [152] Lehmann G, Mueller EA: [Ultraviolet irradiation and altitude fitness.] *Luftfahrtmedizin.* 1944;9:37–43. [Article in German].
- [153] Sigmund R: [Effect of ultraviolet rays on reaction time in man.] *Strahlentherapie.* 1956;101(4):623–9. [Article in German].
- [154] Rosentsweig J: The effect of a single suberythemic biodose of ultraviolet radiation upon the strength of college women. *J Assoc Phys Ment Rehabil.* 1967;21(4):131–3.
- [155] Rosentsweig J: The effect of a single suberythemic biodose of ultraviolet radiation upon the endurance of college women. *J Sports Med Phys Fitness.* 1969;9(2):104–6.
- [156] Cheatum BA: Effects of a single biodose of ultraviolet radiation upon the speed of college women. *Res Q.* 1968;39(3):482–5.
- [157] Macdougall JD, Hicks AL, Macdonald JR, Mckelvie RS, Green HJ, Smith KM: Muscle performance and enzymatic adaptations to sprint interval training. *J Appl Physiol.* 1998;84:2138–142.
- [158] Dahlquist DT, Dieter BP, Koehle MS: Plausible ergogenic effects of vitamin D on athletic performance and recovery. *J Int Soc Sports Nutr.* 2015;12:33. doi: 10.1186/s12970-015-0093-8.
- [159] Bouillon R, Verstuyf A: Vitamin D, mitochondria, and muscle. *J Clin Endocrinol Metabol.* 2013;98(3):961–3. doi: 10.1210/jc.2013-1352.
- [160] Redzic M, Lewis RM, Thomas DT: Relationship between 25-hydroxyvitamin D, muscle strength, and incidence of injury in healthy adults: a systematic review. *Nutr Res.* 2013;33(4):251–8. doi: 10.1016/j.nutres.2013.02.007.
- [161] Ward KA, Das G, Berry JL, Roberts SA, Rawer R, Adams JE, Mughal Z: Vitamin D status and muscle function in post-menarchal adolescent girls. *J Clin Endocrinol Metab.* 2009;94(2):559–63. doi: 10.1210/jc.2008-1284.

- [162] Foo LH, Zhang Q, Zhu K, Ma G, Hu X, Greenfield H, Fraser DR. Low vitamin D status has an adverse influence on bone mass, bone turnover, and muscle strength in Chinese adolescent girls. *J Nutr*. 2009;139(5):1002–7. doi: 10.3945/jn.108.102053.
- [163] Pojednic RM, Ceglia L: The emerging biomolecular role of vitamin D in skeletal muscle. *Exerc Sport Sci Rev*. 2014;42(2):76–81. doi: 10.1249/JES.0000000000000013.
- [164] Gerdhem P, Ringsberg KAM, Obrant KJ, Akesson K: Association between 25-hydroxy vitamin D levels, physical activity, muscle strength and fractures in the prospective population-based OPRA Study of Elderly Women. *Osteoporos Int*. 2005;16(11):1425–31.
- [165] Shardell M, Hicks GE, Miller RR, Kritchevsky S, Andersen D, Bandinelli S, Cherubini A, Ferrucci L. Association of low vitamin D levels with the frailty syndrome in men and women. *J Gerontol (A Biol Sci Med Sci)*. 2009;64(1):69–75. doi: 10.1093/gerona/gln007.
- [166] Houston DK, Toozé JA, Davis CC, Chaves PH, Hirsch CH, Robbins JA, Arnold AM, Newman AB, Kritchevsky SB: Serum 25-hydroxyvitamin D and physical function in adults of advanced age: the CHS All Stars. *J Am Geriatr Soc*. 2011;59(10):1793. doi: 10.1111/j.1532-5415.2011.03601.x.
- [167] Visser M, Deeg DJH, Puts MTE, Seidell JC, Lips P. Low serum concentrations of 25-hydroxyvitamin D in older persons and the risk of nursing home admission. *Am J Clin Nutr*. 2006;84(3):616–22.
- [168] Bischoff-Ferrari HA, Dietrich T, Orav EJ, Hu FB, Zhang Y, Karlson EW, et al. Higher 25-hydroxyvitamin D concentrations are associated with better lower-extremity function in both active and inactive persons aged  $\geq 60$  years. *Am J Clin Nutr*. 2004;80(3):752–8.
- [169] Tieland M, Brouwer-Brolsma E, Nienaber-Rousseau C, van Loon L, De Groot L: Low vitamin D status is associated with reduced muscle mass and impaired physical performance in frail elderly people. *Eur J Clin Nutr*. 2013;67(10):1050–5. doi: 10.1038/ejcn.2013.144.
- [170] Dufour AB, Hannan MT, Murabito JM, Kiel DP, McLean RR: Sarcopenia definitions considering body size and fat mass are associated with mobility limitations: the Framingham Study. *J Gerontol A Biol Sci Med Sci*. 2013;68:168–74. doi: 10.1093/gerona/gls109.
- [171] Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, Martin FC, Michel J-P, Rolland Y, Schneider SM, Topinková E, Vandewoude M, Zamboni M: Sarcopenia: European consensus on definition and diagnosis: report of the European Working Group on Sarcopenia in Older People. *Age Ageing*. 2010;39:412–23. doi: 10.1093/ageing/afq034.
- [172] Gilsanz V, Kremer A, Mo AO, Wren TA, Kremer R: Vitamin D status and its relation to muscle mass and muscle fat in young women. *Clin Endocrinol Metab*. 2010;95(4):1595–601. doi: 10.1210/jc.2009-2309.

- [173] Dawson-Hughes B: 70th Anniversary Conference on Vitamins in early development and healthy aging: impact on infectious and chronic disease' symposium 1: vitamins and cognitive development and performance Serum 25-hydroxyvitamin D and muscle atrophy in the elderly. *Proc Nutr Soc.* 2012;71:46–9. doi:10.1017/S0029665111003260
- [174] Houston DK, Cesari M, Ferrucci L, Cherubini A, Maggio D, Bartali B, Johnson MA, Schwartz GG, Kritchevsky SB: Association between vitamin D status and physical performance: the InCHIANTI study. *J Gerontol A Biol Sci Med Sci.* 2007;62(4):440–6.
- [175] Dam TT, von Mühlen D, Barrett-Connor EL: Sex-specific association of serum vitamin D levels with physical function in older adults. *Osteoporos Int.* 2009;20(5):751–60.
- [176] Okuno J, Tomura S, Yabushita N, Kim MJ, Okura T, Tanaka K, Yanagi H: Effects of serum 25-hydroxyvitamin D(3) levels on physical fitness in community-dwelling frail women. *Arch Gerontol Geriatr.* 2010;50(2):121–6. doi: 10.1016/j.archger.2009.02.011.
- [177] Wicherts IS, van Schoor NM, Boeke AJ, Visser M, Deeg DJ, Smit J, Knol DL, Lips P: Vitamin D status predicts physical performance and its decline in older persons. *J Clin Endocrinol Metab.* 2007;92(6):2058–65.
- [178] Mastaglia SR, Seijo M, Muzio D, Somoza J, Nuñez M, Oliveri B: Effect of vitamin D nutritional status on muscle function and strength in healthy women aged over sixty-five years. *Nutr Health Aging.* 2011;15(5):349–54.
- [179] Stockton KA, Mengersen K, Paratz JD, Kandiah D, Bennell KL: Effect of vitamin D supplementation on muscle strength: a systematic review and meta-analysis. *Osteoporos Int.* 2011 22(3):859–71. doi: 10.1007/s00198-010-1407-y.
- [180] Pramyothin P, Techasurungkul S, Lin J, Wang H, Shah A, Ross PD, Puapong R, Wasnich RD: Vitamin D status and falls, frailty, and fractures among postmenopausal Japanese women living in Hawaii. *Osteoporos Int.* 2009;20(11):1955–62. doi: 10.1007/s00198-009-0910-5.
- [181] Ceglia L, Chiu GR, Harris SS, Araujo AB: Serum 25-hydroxyvitamin D concentration and physical function in adult men. *Clin Endocrinol (Oxf).* 2011;74(3):370–6. doi: 10.1111/j.1365-2265.2010.03926.x.
- [182] Mathei C, Van Pottelbergh G, Vaes B, Adriaensen W, Gruson D, Degryse JM: No relation between vitamin D status and physical performance in the oldest old: results from the Belfrail study. *Age Ageing.* 2013;42(2):186–90. doi: 10.1093/ageing/afs186.
- [183] Allali F, El Aichaoui S, Khazani H, Benyahia B, Saoud B, El Kabbaj S, Bahiri R, Abouqal R, Hajjaj-Hassouni N: High prevalence of hypovitaminosis D in Morocco: relationship to lifestyle, physical performance, bone markers, and bone mineral density. *Semin Arthritis Rheum.* 2009;38:444–51. doi: 10.1016/j.semarthrit.2008.01.009.
- [184] Gamero P, Munoz F, Sornay-Rendu E, Delmas PD: Associations of vitamin D status with bone mineral density, bone turnover, bone loss and fracture risk in healthy postmenopausal women. The OFELY study. *Bone.* 2007;40:716–22.

- [185] Ceglia L: Vitamin D and its role in skeletal muscle. *Curr Opin Clin Nutr Metab Care*. 2009;12(6):628–33. doi: 10.1097/MCO.0b013e328331c707.
- [186] Prince RL, Austin N, Devine A, Dick IM, Bruce D, Zhu K: Effects of ergocalciferol added to calcium on the risk of falls in elderly high-risk women. *Arch Intern Med*. 2008;168(1):103–8. doi: 10.1001/archinternmed.2007.31.
- [187] Pfeifer M, Begerow B, Minne H, Suppan K, Fahrleitner-Pammer A, Dobnig H: Effects of a long-term vitamin D and calcium supplementation on falls and parameters of muscle function in community-dwelling older individuals. *Osteoporos Int*. 2009;20:315–22.
- [188] Zhu K, Austin N, Devine A, Bruce D, Prince R: A randomized controlled trial of the effects of vitamin D on muscle strength and mobility in older women with vitamin D insufficiency. *J Am Geriatr Soc*. 2010;58(11):2063–8. doi: 10.1111/j.1532-5415.2010.03142.x
- [189] Bischoff-Ferrari HA, Dawson-Hughes B, Willett WC, Staehelin HB, Bazemore MG, Zee RY, Wong JB: Effect of Vitamin D on falls: a meta-analysis. *JAMA*. 2004;291(16):1999–2006.
- [190] Tracy BL, Ivey FM, Hurlbut D, Martel GF, Lemmer JT, Siegel EL, Metter EJ, Fozard JL, Fleg JL, Hurley BF: Muscle quality. II. Effects of strength training in 65- to 75-yr-old men and women. *J Appl Physiol*. 1999;86(1):195–201.
- [191] Bunout D, Barrera G, Leiva L, Gattas V, de la Maza MP, Avendaño M, Hirsch S: Effects of vitamin D supplementation and exercise training on physical performance in Chilean vitamin D deficient elderly subjects. *Exp Gerontol*. 2006;41:746–52.
- [192] Bischoff HA, Stähelin HB, Dick W, Akos R, Knecht M, Salis C, Nebiker M, Theiler R, Pfeifer M, Begerow B, Lew RA, Conzelmann M.: Effects of vitamin D and calcium supplementation on falls: a randomized controlled trial. *J Bone Miner Res*. 2003;18:343–51.
- [193] Moreira-Pfrimer LD, Pedrosa MA, Teixeira L, Lazaretti-Castro M: Treatment of vitamin D deficiency increases lower limb muscle strength in institutionalized older people independently of regular physical activity: a randomized double-blind controlled trial. *Ann Nutr Metab*. 2009;54(4):291–300. doi: 10.1159/000235874.
- [194] Pfeifer M, Begerow B, Minne H, Suppan K, Fahrleitner-Pammer A, Dobnig H: Effects of a long-term vitamin D and calcium supplementation on falls and parameters of muscle function in community-dwelling older individuals. *Osteoporos Int*. 2009;20:315–22.
- [195] Rejnmark R: Effects of vitamin D on muscle function and performance: a review of evidence from randomized controlled trials. *Ther Adv Chronic Dis*. 2011;2(1):25–37. doi: 10.1177/2040622310381934.



- [196] Janssen HC, Samson MM, Verhaar HJ: Muscle strength and mobility in vitamin D-insufficient female geriatric patients: a randomized controlled trial on vitamin D and calcium supplementation. *Aging Clin Exp Res.* 2010;22(1):78–84.
- [197] Brunner RL, Cochrane B, Jackson RD, Larson J, Lewis C, Limacher M, Rosal M, Shumaker S, Wallace R; Women's Health Initiative Investigators. Calcium, vitamin D supplementation, and physical function in the Women's Health Initiative. *J Am Diet Assoc.* 2008;108(9):1472–9. doi: 10.1016/j.jada.2008.06.432.
- [198] Fuleihan G, Nabulsi M, Tamim H, Maalouf J, Salamoun M, Khalife H, Choucair M, Arabi A, Vieth R: Effect of vitamin D replacement on musculoskeletal parameters in school children: a randomized controlled trial. *J Clin Endocrinol Metab.* 2006;91(2):405–12.
- [199] Lim JS, Kim KM, Rhee Y, Lim SK. Gender-dependent skeletal effects of vitamin D deficiency in a younger generation. *J Clin Endocrinol Metab.* 2012 97(6):1995–2004. doi: 10.1210/jc.2011-3098.
- [200] Gupta R, Sharma U, Gupta N, Kalaivani M, Singh U, Guleria R, Jagannathan NR, Goswami R: Effect of cholecalciferol and calcium supplementation on muscle strength and energy metabolism in vitamin D-deficient Asian Indians: a randomized, controlled trial. *Clin Endocrinol (Oxf).* 2010;73:445–51. doi: 10.1111/j.13 65-2265.2010.03816.x.
- [201] Owens DJ, Webber D, Impey SG, Tang J, Donovan TF, Fraser WD, Morton JP, Close GL: Vitamin D supplementation does not improve human skeletal muscle contractile properties in insufficient young males. *Eur J Appl Physiol.* 2014;114(6):1309–20. doi: 10.1007/s00421-014-2865-2.
- [202] Tomlinson PB, Joseph C, Angioi M: Effects of vitamin D supplementation on upper and lower body muscle strength levels in healthy individuals. A systematic review with meta-analysis. *J Sci Med Sport.* 2015;18:575–80. doi: 10.1016/j.jsams.2014.07.022.
- [203] Barker T, Henriksen VT, Martins TB, Hill HR, Kjeldsberg CR, Schneider ED, Dixon BM, Weaver LK: Higher serum 25-hydroxyvitamin D concentrations associate with a faster recovery of skeletal muscle strength after muscular injury. *Nutrients.* 2013;5(4):1253–75. doi: 10.3390/nu5041253.
- [204] Mowry DA, Costello MM, Heelan KA: Association among cardiorespiratory fitness, body fat, and bone marker measurements in healthy young females. *J Am Osteopath Assoc.* 2009;109(10):534–9.
- [205] Forney LA, Earnest CP, Henagan TM, Johnson LE, Castleberry TJ, Stewart LK: Vitamin D status, body composition, and fitness measures in college-aged students. *J Strength Cond Res.* 2014;28(3):814–24. doi: 10.1519/JSC.0b013e3182a35ed0.
- [206] Fitzgerald JS, Peterson BJ, Warpeha JM, Wilson PB, Rhodes GS, Ingraham SJ: Vitamin D status and V[combining dot above]O<sub>2</sub>peak during a skate treadmill graded exercise

- test in competitive ice hockey players. *J Strength Cond Res.* 2014;28(11):3200–5. doi: 10.1519/JSC.0000000000000523.
- [207] Dubnov-Raz G, Livne N, Raz R, Rogel D, Cohen AH, Constantini NW: Vitamin D concentrations and physical performance in competitive adolescent swimmers. *Pediatr Exerc Sci.* 2014;26:64–70. doi: 10.1123/pes.2013-0034.
- [208] Hamilton B, Whiteley R, Farooq A, Chalabi H: Vitamin D concentration in 342 professional football players and association with lower limb isokinetic function. *J Sci Med Sport.* 2014;17:139–43. doi: 10.1016/j.jsams.2013.03.006.
- [209] Ardestani A, Parker B, Mathur S, Clarkson P, Pescatello LS, Hoffman HJ, Polk DM, Thompson PD: Relation of vitamin D level to maximal oxygen uptake in adults. *Am J Cardiol.* 2011;107(8):1246–9. doi: 10.1016/j.amjcard.2010.12.022.
- [210] Wilson PB, Fitzgerald JS, Ingraham SJ: Relationship between serum 25-hydroxyvitamin D status and cardiorespiratory fitness: findings from the National Health and Nutrition Examination Survey. *Int J Cardiol.* 2013;168:3112–3. doi: 10.1016/j.ijcard.2013.04.069.
- [211] Close GL, Russell J, Cobley JN, Owens DJ, Wilson G, Gregson W, Fraser WD, Morton JP: Assessment of vitamin D concentration in non-supplemented professional athletes and healthy adults during the winter months in the UK: implications for skeletal muscle function. *J Sports Sci.* 2013;31(4):344–53. doi: 10.1080/02640414.2012.733822.
- [212] Wyon MA, Koutedakis Y, Wolman R, Nevill AM, Allen N: The influence of winter vitamin D supplementation on muscle function and injury occurrence in elite ballet dancers: a controlled study. *J Sci Med Sport.* 2014;17(1):8–12. doi: 10.1016/j.jsams.2013.03.007.
- [213] Wyon MA, Wolman R, Nevill AM, Cloak R, Metsios GS, Gould D, Ingham A, Koutedakis Y: Acute effects of vitamin D3 supplementation on muscle strength in Judoka athletes: a randomized placebo-controlled, double-blind trial. *Clin J Sport Med.* 2015. [Epub ahead of print]
- [214] Close GL, Leckey J, Patterson M, Bradley W, Owens DJ, Fraser WD, Morton JP: The effects of vitamin D(3) supplementation on serum total 25[OH]D concentration and physical performance: a randomised dose–response study. *Br J Sports Med.* 2013;47:692–6. doi: 10.1136/bjsports-2012-091735.
- [215] Jastrzębski Z: Effect of vitamin D supplementation on the level of physical fitness and blood parameters of rowers during the 8-week high intensity training. *Facila Educ Fiz și Sport.* 2014;2:57–67.
- [216] Jastrzębska M, Kaczmarczyk M, Jastrzębski Z: The effect of vitamin d supplementation on training adaptation in well trained soccer players. *Strength Cond Res.* 2016; [Epub ahead of print].