

PUBLISHED BY

# INTECH

open science | open minds

World's largest Science,  
Technology & Medicine  
Open Access book publisher



**3,100+**  
OPEN ACCESS BOOKS



**103,000+**  
INTERNATIONAL  
AUTHORS AND EDITORS



**102+ MILLION**  
DOWNLOADS



**BOOKS**  
DELIVERED TO  
151 COUNTRIES

AUTHORS AMONG

**TOP 1%**  
MOST CITED SCIENTIST



**12.2%**  
AUTHORS AND EDITORS  
FROM TOP 500 UNIVERSITIES



Selection of our books indexed in the  
Book Citation Index in Web of Science™  
Core Collection (BKCI)

**WEB OF SCIENCE™**

Chapter from the book *A Critical Evaluation of Vitamin D - Basic Overview*

Downloaded from: <http://www.intechopen.com/books/a-critical-evaluation-of-vitamin-d-basic-overview>

Interested in publishing with InTechOpen?  
Contact us at [book.department@intechopen.com](mailto:book.department@intechopen.com)

# Vitamin D and Sphingolipids: Role in Bone and Neural System

Alessia Frati, Mercedes Garcia Gil,  
Federica Pierucci and Elisabetta Meacci

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/66648>

## Abstract

1-Alpha,25-dihydroxyvitamin D<sub>3</sub> (1,25(OH)<sub>2</sub>D<sub>3</sub>) is known to play an important physiological role on growth and differentiation in a variety of nonmalignant and malignant cell types through classical actions, mediated by its specific receptor (VDR), and nongenomic actions resulting in the activation of specific signalling pathways. Due to the broad distribution of Vitamin D Receptor (VDR) in many tissues and the ability of 1,25(OH)<sub>2</sub>D<sub>3</sub> to regulate fundamental processes, such as cell proliferation and differentiation, this steroid hormone has been suggested in the treatment of different diseases, from cancer to neurodegenerative diseases. In fact, structural 1,25(OH)<sub>2</sub>D<sub>3</sub> analogues, with weaker collateral effects, have recently entered in clinical trials. Other interesting molecules due to their pleiotropic actions are the bioactive sphingolipids (SLs), in particular ceramide (Cer) and sphingosine 1-phosphate (SIP). Cells maintain a dynamic balance of these metabolites since Cer and sphingoid bases mediate cell death, while SIP exerts mitogenic effects and promotes differentiation of several cell types including osteogenic and neural cells. The biological actions of 1,25(OH)<sub>2</sub>D<sub>3</sub> and SLs, in particular SIP, share many common effectors, including calcium regulation, growth factor expression, inflammatory cytokines, etc., but whether they could act synergistically is still unknown and deserves further investigation.

**Keywords:** vitamin D, sphingosine 1-phosphate, ceramide, neurodegeneration, bone

## 1. Introduction

1-Alpha,25-dihydroxyvitamin D<sub>3</sub> (1,25(OH)<sub>2</sub>D<sub>3</sub>), a known regulator of calcium and phosphorus homeostasis, has also important physiological effects on growth and differentiation in a variety of nonmalignant and malignant cell types [1–4] and a central role in host defense

against infections [5]. The classical actions of  $1,25(\text{OH})_2\text{D}_3$  start with the hormone binding to the  $1,25(\text{OH})_2\text{D}_3$  receptor/retinoic X receptor (VDR/RXR) heterodimeric complex to specific DNA sequences, whereas the rapid nongenomic actions result in the activation of specific signal transduction pathways [6, 7]. The broad distribution of VDR in the human body and the ability of  $1,25(\text{OH})_2\text{D}_3$  to control cell growth and differentiation make this hormone a potentially useful agent in the treatment of diseases, including cancer and neurodegenerative diseases. However, the systemic application of  $1,25(\text{OH})_2\text{D}_3$  is limited because of its hypercalcemic side effects [8]. Therefore,  $1,25(\text{OH})_2\text{D}_3$  analogues, with potent cell regulatory effects, but with weaker effects on calcium metabolism, have recently been obtained and some of them have entered in clinical trials [9, 10]. Sphingolipids (SLs) constitute a biologically active lipid class that is significantly important from both structural and regulatory aspects [11–14]. Indeed, they regulate fundamental cellular processes that are important in determining cellular fate, such as proliferation, apoptosis senescence, and inflammation [15–19]. Cells maintain a dynamic balance of distinct SL metabolites [20, 21], with ceramide (Cer) and sphingoid bases acting in opposite manner with respect to sphingosine 1-phosphate (S1P) that exerts mitogenic effects and promote differentiation of several cell types including skeletal muscle cells [22, 23] and neural cells [24–27]. The manipulation of SL metabolism is currently being studied as a novel strategy to regulate cell proliferation/inflammation [18, 19, 28, 29]. Interestingly, S1P can be released from many cell types including neuronal cells and osteoblasts, thus, acting as ligand of specific S1P receptors, triggers paracrine and autocrine signalling [13, 19, 30]. In the present chapter, we review the potential contribution of the biological effects of sphingolipids,  $1,25(\text{OH})_2\text{D}_3$  and its structural analogues in bone and neural disorders.

## 2. Vitamin D in bone physiology, osteogenesis, and osteoporosis

### 2.1. Bone remodelling

Bone remodelling consists in the balance between the elimination of bone due to osteoclast death and the formation of new bone by osteoblast proliferation [31]. The migration of mesenchymal stem cells (MSCs) to areas of new bone formation is also a fundamental process for skeleton maintenance. In fact, in these functional locations, MSCs, under the influence of bone morphogenetic protein, differentiate into osteoblasts. The majority of osteoblasts become osteocytes, the fully differentiated cells, within the bone matrix, where they also help in tissue repair [32]. Osteoclasts are multinucleated cells, abundant in mitochondria, vacuoles, and lysosomes [33, 34]. These cells derive from the fusion of preosteoclasts through a mechanism that is regulated by the dendritic cell-specific transmembrane protein (DC-STAMP) and the osteoclast-stimulatory transmembrane protein (OC-STAMP) [35, 36]. Osteoclasts are characterized by a ruffled border in contact with the bone surface, where the vacuolar  $\text{H}^+$ -ATPase, responsible of the maintenance of acid pH that favors the dissolution of the bone minerals, is localized. The zone beneath the ruffled borders is called resorption lacunae and is isolated from the surrounding by the sealing zone of osteoclasts that attaches the cells to the bone surface. Cathepsin K, MMP9, and tartrate resistant-acid phosphatase (TRAP), the main enzymes responsible for the degradation of bone matrix, are released in the resorption

lacunae. Matrix degradation products are endocytosed from the central portion of the ruffled border, packaged into transcytotic vesicles and secreted from the functional secretory domain [37]. The bone matrix represents an important storage of factors secreted by the osteoblasts during bone formation. Among them are the transforming growth factor  $\beta$  (TGF $\beta$ ), the bone morphogenetic proteins (BMPs), fibroblast growth factors (FGFs), etc. They are stored in the bone matrix and serve as osteoblast-promoting components once liberated and activated by the osteoclasts.

In 1981, Rodan and Martin [38] observed that osteoblasts, but not osteoclasts, express the receptors of bone-resorbing factors, such as Parathyroid hormone (PTH) and prostaglandin E<sub>2</sub>, and proposed that osteoblasts may interfere in the process of osteoclastic resorption. Experimental evidence of the role of the microenvironment provided by osteoblasts for cell differentiation of splenic precursors, confirms this hypothesis [39]. In response to bone-resorption stimulating factors, osteoblasts produce Receptor Activator of Nuclear Factor kappa-B Ligand (RANKL), a membrane associate factor that, by binding to its receptor constitutively expressed on the surface of osteoclast precursors, stimulates cell differentiation and the activation of bone resorption. Another potential physiological regulator of bone mass is the prolin/arginine-rich end leucine-rich repeat protein (PRELP), a heparin/heparan sulfate-binding protein expressed in developing bone, cartilage, and basement membranes. PRELP inhibits osteoclast formation with a mechanism that affects the RANKL-dependent late stage of osteoclastogenesis, and its administration is reported to reduce bone loss in ovariectomized and tumor-bearing mice [40].

## 2.2. Effect of vitamin D on calcium homeostasis and bone remodeling

1,25(OH)<sub>2</sub>D<sub>3</sub> was historically discovered as an anti-rachitic agent due to its effects on the demineralization process carried out by osteoclasts and intestinal calcium absorption [41]. The hormonally active form is the dihydroxylated metabolite 1,25-dihydroxyvitamin D<sub>3</sub>, or 1,25(OH)<sub>2</sub>D<sub>3</sub>. It is generated by two enzymatic hydroxylation reactions, which occur first in the liver to produce 25-hydroxyvitamin D<sub>3</sub> and second in the kidney, where both 1,25(OH)<sub>2</sub>D<sub>3</sub> and its sister metabolite 24,25-dihydroxyvitamin D<sub>3</sub> (24,25[OH]<sub>2</sub>D<sub>3</sub>) are produced. Thus, most tissues have the ability to convert 1,25(OH)<sub>2</sub>D<sub>3</sub> into its active form, which, in turn, will bind to hormone nuclear receptor (VDR) [42]. The presence of the VDR in many tissues that are not involved in mineral metabolism indicates a wider physiological role for 1,25(OH)<sub>2</sub>D<sub>3</sub> able to positively or negatively influence target gene expression via binding of the hormone/VDR complex to specific receptor response elements, i.e., RANKL gene [43]. 1,25(OH)<sub>2</sub>D<sub>3</sub> ability to maintain serum calcium homeostasis is due to VDR-mediated signaling affecting bone physiology and intestinal calcium absorption.

In the intestine, 1,25(OH)<sub>2</sub>D<sub>3</sub>-mediated processes can occur by two different mechanisms: (a) the *paracellular pathway* predominates when dietary levels of calcium are high, and it is a passive, nonsaturable diffusion process [44], calcium is transported through tight junctions, and involves several claudins proteins upregulated by 1,25(OH)<sub>2</sub>D [45]; (b) the *transcellular pathway* occurs in the jejunum and duodenum in circumstances of low calcium dietary intake. It requires an active transport of calcium through the thickness of the enterocytes, and it is regulated by the subsequent involvement of transient receptor potential vanilloid

6 (TRPV6), calcium-binding protein calbindin  $D_{9k}$  (CaBP-9K), and a calcium ATPase. The transcription of these proteins is stimulated by the  $1,25(\text{OH})_2\text{D}$  signal [46]. When intestinal calcium absorption decreases, the transcellular transport in the kidney, a process similar to the active transport through the enterocytes, increases calcium reabsorption. This pathway is also stimulated by  $1,25(\text{OH})_2\text{D}_3$  [44].

In addition, in hypocalcemia conditions, parathyroid glands release PTH which, in the kidney, stimulate the production of the active form of  $1,25(\text{OH})_2\text{D}_3$ . Whenever intestinal and renal calcium fluxes are insufficient to maintain the correct calcium levels in the serum, the bone represents an additional pool of calcium and the bone-resorbing process permits to adjust serum calcium concentration.

$1,25(\text{OH})_2\text{D}_3$  activates multiple signaling pathways in bone precursor cells [42]—*nongenomic pathways* that involve: (a) the activation of voltage-sensitive calcium channels located at plasma membrane; (b) the release of calcium from intracellular stores; (c) a shift in the charge state of the matrix protein osteopontin (OPN)—*a classical nuclear receptor-mediated event* that leads to the upregulation of the OPN gene at 48 h after hormone addition. Nanomolar concentration of  $1,25(\text{OH})_2\text{D}_3$  is necessary for the nonclassical effects, while physiological  $1,25(\text{OH})_2\text{D}_3$  serum concentrations are necessary in the picomolar range [42].

Calcium balance and cell differentiation stage affect VDR action in osteogenic cells. In fact during a positive calcium balance,  $1,25(\text{OH})_2\text{D}_3$  signaling acts in a different way in osteoblasts at diverse differentiation stage: in immature osteoblasts, the hormone causes an increase in RANKL expression leading to a catabolic function on bone mass [47], whereas in mature osteoblasts, VDR stimulation produces a decrease in RANKL expression and an increase in the production of the osteoclastogenic inhibitor osteoprotegerin (OPG), which inhibits RANKL-RANK interaction [48].

During a negative calcium balance, the high RANKL/OPG ratio and the increased levels of mineralization inhibitors, such as OPN, are the two events that permit the mobilization of calcium from the bone to the serum in response to  $1,25(\text{OH})_2\text{D}_3$  and PTH signaling. By increasing calcium absorption, hormone signaling indirectly preserves bone mass mineralization. In osteocytes, VDR signaling upregulates the transcription of the bone-derived fibroblast growth factor-23 (FGF23) that stimulates TRPV5 expression, thus, increasing renal phosphate excretion [49]. Notably, FGF23 also decreases renal CYP27B1 activity, the enzyme that catalyzes the hydroxylation of  $25(\text{OH})_2\text{D}_3$  to  $1,25(\text{OH})_2\text{D}_3$  [42], avoiding overstimulation of the hormone pathway.

Osteoporosis is a systemic skeletal disease characterized by a reduction of bone mass and microarchitectural deterioration that leads to increase in bone fragility and susceptibility to fracture [50]. Osteoporosis results from the imbalance between bone resorption and bone formation, and  $1,25(\text{OH})_2\text{D}_3$  can regulate both aspects of bone turnover. Nearly in all studies, the treatment with  $1,25(\text{OH})_2\text{D}_3$  or its precursor,  $1\alpha(\text{OH})\text{D}_3$ , is found to increase bone mineral density [51, 52]. More recently, the combination of  $1\alpha(\text{OH})\text{D}_3$  with an anti-resorptive bisphosphonate (alendronate) enhanced bone mass with fewer falls and fractures [53].

### 2.3. Vitamin D deficiency and supraphysiological dose

The lack of  $1,25(\text{OH})_2\text{D}_3$  has obvious implications for human diseases. Severe  $1,25(\text{OH})_2\text{D}_3$  deficiency causes nutritional rickets in children [54] that can easily be prevented and cured by  $1,25(\text{OH})_2\text{D}_3$  supplementation [55]. Moreover, deficiency of  $1,25(\text{OH})_2\text{D}_3$  or reduced level of VDR can lead to osteomalacia disease and the recovery of a normal hormone level can resolve the disease [56]. Less severe  $1,25(\text{OH})_2\text{D}_3$  deficiency produces an increased bone turnover and an accelerated bone loss and it is associated with osteoporosis. A correlation between the sun exposure and bone physiology in regions insufficiently exposed to sunlight may involve  $1,25(\text{OH})_2\text{D}_3$  deficiency [57]. However, since  $1,25(\text{OH})_2\text{D}_3$  insufficiency is widespread, it is difficult to provide evidence regarding the direct role of sunlight on osteogenesis and osteoporosis without taking into account many other variables, such as personal bone physiology, age, weight, clothing habits, medication, and others.

Supraphysiological doses of the  $1,25(\text{OH})_2\text{D}_3$  induce calcemic side effects. In order to preserve or augment the beneficial effects of the hormone and to minimize its collateral consequences, structural analogues of  $1,25(\text{OH})_2\text{D}_3$  have been synthesized by introducing chemical modifications in the A-ring, central CD-ring region, or side chain of the hormone [58].

Some of these analogues have tissue-specific actions, exert prodifferentiating and antiproliferative effects on keratinocytes, and also possess important anti-inflammatory properties. The recently approved eldecalcitol ( $1\alpha,25[\text{OH}]_2\text{-}2\text{b-(3-hydroxypropyloxy)vitamin D}_3$ ; ED-71; Ediol®) is an orally administered analogue of calcitriol that binds to VDR [59]. Ediol® is available for the treatment of osteoporosis [60]. The effects of this compound on bone metabolism have been reported in a randomized, open-label study in postmenopausal women, in which reductions in the markers of bone reabsorption were observed [61]. Similarly, in a randomized, noncomparative study of patients with osteoporosis, the structural analogue suppresses the biochemical markers of bone turnover in a dose-dependent manner.

Another  $1,25(\text{OH})_2\text{D}_3$  analogue with some effects on bone is ZK191784, a compound characterized by 22,23-double bond, 24R-hydroxy group, 25-cyclopropyl ring, and 5-butyloxazole-group [10]. It has been shown to exert therapeutic potential in T cell-mediated immune disorders and to significantly counteract acute and chronic intestinal inflammation [62]. While the  $1,25(\text{OH})_2\text{D}_3$  agonist effect of ZK191784 in kidney and antagonistic effect in intestine were clear, its effect on bone is still Ongoing: preliminary analyses appeared to suggest a tendency toward restoration of the reduced bone thickness in mice lacking the renal epithelial calcium channel TRPV ( $\text{Trpv5}^{-/-}$ ) [37].

Another  $1,25(\text{OH})_2\text{D}_3$  analogue, Seocalcitol, is able to reduce the number and growth of metastasis originating from various types of cancer cells, such as bone metastasis originating from intracardially injected breast cancer cells [63]. Since  $1,25(\text{OH})_2\text{D}_3$  and its analogues possess cytostatic properties, many in vivo studies have focused on hormone analog cancer treatment combined with radiotherapy and/or chemotherapy. However, while the combination of Seocalcitol, with radiotherapy in a xenograft model for breast cancer, lead to more effective anti-cancer effects [64, 65], the combination of the analogues with chemotherapy does not always result in additive or synergistic effects. In view of the promising results that certain



hormone analogs show against cancer in vitro and in vivo animal models, some of them have been tested in cancer patients, such as Seocalcitol, which however has given rather insufficient results in clinical trials.

#### 2.4. Animal model for bone remodeling study

Several animal models have been developed and used to understand the pathogenesis of osteoporosis and osteogenesis for the preclinical testing of new treatment options [66]. Only few of them were used in combination with  $1,25(\text{OH})_2\text{D}_3$  treatment.

The senescence accelerated mouse (SAM/P6) is a mouse model for severe osteoporosis that has low level of bone mass and develops fractures in old age [67]. It is a unique model for the study of age-related osteopenia and severe osteoporosis mimicking many aspects of the age-related changes seen in bones of humans and offering the opportunity to study relevant genes that contribute to this process.

Other studies have been performed in ovariectomy (OVX) on rats for either osteoporotic induction or fracture healing. Notably, an additive effect on bone loss was reported by combined OVX-deficient calcium or OVX-deficient  $1,25(\text{OH})_2\text{D}_3$ . In addition, the rat model subjected to ovariectomy and multid deficiency diet (depletion of  $1,25(\text{OH})_2\text{D}_3$ , calcium, vitamin K, and phosphorus), and, thus, characterized by increased bone turnover could contribute to the study of bone- and energy metabolism in early and late stages of osteoporosis. Several ovariectomized large animals might be also used as models of osteoporosis, such as the dog, the pig, the sheep, and the nonhuman primates [68]. In particular, the sheep is also well established as a model for human bone loss/osteoporosis in orthopedic research [69]. Some limits can be the largeness in size and the difficulty to manage relatively expensive experiments.

Mouse can be a reliable animal model of glucocorticoid-induced osteopenia/osteoporosis and mimic the changes seen in humans [70]. Mice receiving glucocorticoid for a week showed an early increase in bone resorption, decreased bone mineral density, and bone mass.

Transgenic mice showing bone alterations have been also developed. Klotho mouse is a transgenic mouse model obtained by an insertion mutation that disrupts the Klotho gene locus. Klotho is a gene encoding a transmembrane protein that forms a complex with multiple fibroblast growth factor receptors and functions as coreceptor for FGF23, an osteocyte-derived hormone that induces negative phosphate balance. Defects in either Klotho or FGF23 gene expression result in osteopenia [71]. In these animals, similarly to what occurs in human senile osteoporosis, the reduction in bone formation occurs faster than bone resorption. Tg $\text{huRANKL}$  (Tg5519) is a transgenic mouse overexpressing human RANKL resulting in the spontaneous development of osteoporosis similar to human pathology [71, 72]. The overexpression of huRANKL results in the spontaneous development of early onset osteoporosis characterized by lack of trabecular bone, increased osteoclastogenesis, increased bone remodeling, and decreased bone strength. Another model, already mentioned in this paragraph is TRPV5 $^{-/-}$  mice that display hypercalciuria due to a primary renal failure to reabsorb calcium and hypervitaminosis D, leading to calcium hyperabsorption in the intestine and reduced bone mass [37, 73]. Other models are useful to evaluate osteocyte functions. They have been obtained,

for example, by osteocyte-specific disruption of gap junction protein, *Gja1*. Osteocyte-specific *Gja1* conditional knockout mice show an increase in apoptosis of osteocytes [74].

Animal models for disused osteoporosis were also developed. Methods to reduce skeletal biomechanical loading include nerve, spinal cord, or tendon resections, casting, bandaging of one limb or suspension of both hindlimbs in rats. Immobilization-induced osteopenia/osteoporosis in rat skeletal model with the highly predictable pattern of bone loss and the hormone in plasma is significantly decreased [75].

Regarding osteogenesis, most of the studies have focused on osteogenesis imperfecta (OI), an autosomal dominant disorder caused by mutations in type I collagen, the most abundant protein of bone, skin, and tendon extracellular matrices. OI is characterized by increased bone fragility and low bone mass. Transgenic mice expressing a premature stop codon or glycine substitution in the COL1A1 gene encoding the chains of type I collagen are good model for OI [76]. Very recently, the effect of high dose of hormones on bone density in OI patients has been reported [77]. Notably, new genes implicated in autosomal recessive forms of OI have been described and one of each is sphingomyelin phosphodiesterase (SMPD3), the gene encoding for neutral sphingomyelinase [78].

Some conflicting results are reported in literature, but the animal models developed so far have given valuable information on the pathogenesis of osteoporosis as well as on other pathological conditions of the skeleton and bone. The development of new animal models will help to better understand what have been poorly investigated in the past.

### 3. Vitamin D in nervous system physiology, neuroprotection, and neurogenesis

VDR is expressed in both neurons and glial cells (i.e., microglia, astrocytes, and oligodendrocytes) in different regions of the nervous system [3]. Vitamin D Response Element (VRE) response elements modulate gene expression. For example, it increases the expression genes codifying growth factors such as Nerve growth factor (NGF), glial cell line-derived neurotrophic factor (GDNF), NT3, and enzymes involved in the synthesis of neurotransmitters (tyrosine hydroxylase, tryptophan hydroxylase 2, and glutamate decarboxylase), whereas it decreases expression of voltage-dependent calcium channel [3, 79]. VDR is also expressed in the caveolae and induces nongenomic effects that include activation of cAMP-dependent protein kinase (PKA),  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase, phosphatidylinositol 3-kinase, and mitogen-activated protein kinase p38 leading to phosphorylation of neurofilaments, modulation of chloride, potassium, and voltage-dependent calcium channel in rat cortical neurons [80]. In addition, 25-hydroxylase and 1- $\alpha$ -hydroxylase activity are also found in the nervous tissue.

The combination of in vitro, ex vivo, and animal model data provides compelling evidence that  $1,25(\text{OH})_2\text{D}_3$  has a crucial role in neuronal proliferation, differentiation, neurotransmission, neuroplasticity, and neuroprotection. Increasing evidence derived from studies of  $1,25(\text{OH})_2\text{D}_3$  deficiency and from VDR polymorphisms implicates  $1,25(\text{OH})_2\text{D}_3$  as a candidate



in influencing susceptibility to a number of psychiatric and neurological diseases, such as schizophrenia, autism, Parkinson disease (PD), amyotrophic lateral sclerosis, epilepsy, Alzheimer disease (AD), and is especially strong for multiple sclerosis (MS) [8, 81, 82]. In epigenetic studies, maternal dietary deprivation of  $1,25(\text{OH})_2\text{D}_3$  has induced vitamin D deficiency (VDD) in rats prior to mating and maintained it during pregnancy. The  $1,25(\text{OH})_2\text{D}_3$ -deficient rats showed modifications in brain morphology including increased overall brain size and larger lateral ventricles. Interestingly, some changes persist despite the addition of  $1,25(\text{OH})_2\text{D}_3$  to the diet of the pups. In adult life, these rats tend to demonstrate subtle alterations in learning and memory and impaired attentional processing [81]. Prenatal VDD induces similar alterations in fetal mouse brain morphology and mouse behavior [83, 84]. Notably, it has been reported that maternal  $1,25(\text{OH})_2\text{D}_3$  insufficiency during pregnancy in humans is also significantly associated with offspring's language impairment [85]. Interestingly, prenatal  $1,25(\text{OH})_2\text{D}_3$ -depleted rats showed a significant impairment of latent inhibition, a feature often associated with schizophrenia [81].

The neuroprotective effect of  $1,25(\text{OH})_2\text{D}_3$  has been recently reported in cognitive decline of aging rats [86], and it has been extensively studied in the animal model of MS and the experimental allergic encephalomyelitis (EAE). The hormone prevents onset and reversibly blocks progression of clinical signs, but such a protective effect is absent in VDR knockout mice [81]. The effect of  $1,25(\text{OH})_2\text{D}_3$  might not be due exclusively to its neuroimmunomodulatory properties [81] since recently it has been reported that the hormone enhances neural stem cell proliferation and differentiation into neurons and oligodendrocytes, the myelinating cells of central nervous system [4, 87]. Neural stem cells constitutively express VDR, which can be upregulated by  $1,25(\text{OH})_2\text{D}_3$  [4].

$1,25(\text{OH})_2\text{D}_3$  regulates the expression of many AD-related genes. It attenuates A $\beta$  peptide accumulation by stimulating phagocytosis of A $\beta$  peptide probably by modulating transcription of Toll-like receptors and cytokines together with enhancing brain-to-blood efflux transport by increasing P-glycoprotein expression [88].

Adult neurogenesis is limited to specific brain regions in the mammalian brain, such as the hippocampal dentate gyrus and the subventricular zone. Alterations in adult neurogenesis appear to be a common hallmark in different neurodegenerative diseases including PD and AD [89]. Therefore, factors that stimulate neurogenesis have been indicated as possible treatments of neurodegenerative disorders. The antiproliferative and prodifferentiating effects of  $1,25(\text{OH})_2\text{D}_3$  in neural cells were described more than 10 years ago [90, 91].  $1,25(\text{OH})_2\text{D}_3$  decreases the expression of G1/S and G2/M cellular gatekeeper components, such as cyclins D1 and B1 [91, 92] and decreases the percentage of cultured hippocampal cells undergoing mitosis in conjunction with increases in both neurite outgrowth and NGF production [90]. Recently, ceramide kinase signalling pathway has been involved in the antiproliferative action of  $1,25(\text{OH})_2\text{D}_3$  human neuroblastoma cells [18]. Accumulated evidence indicates that  $1,25(\text{OH})_2\text{D}_3$  has complex effects on neurogenesis of neural stem cells. Cui et al. [93] have investigated the effect of Developmental vitamin D (DVD) deficiency on neuroprogenitor formation in the neonatal brain, and they have shown an increase in the number of neurospheres formed in cultures from the neonatal subventricular zone. Exogenous  $1,25(\text{OH})_2\text{D}_3$  added to

the culture medium reduced neurosphere number in control (in agreement with the putative antiproliferative effect of  $1,25(\text{OH})_2\text{D}_3$ ), but not in cultures from the deprived pups [93]. In contrast, neurogenesis in adult subgranular zone of the hippocampus is decreased [94]. In another model of  $1,25(\text{OH})_2\text{D}_3$  deficiency, Zhu et al. [95] have reported increased proliferation, but decreased survival of newborn neurons in the dentate gyrus of adult mice lacking  $1,25(\text{OH})_2\text{D}_3$ , the  $1\alpha$ -hydroxylase knockout mice [95]. The different effects probably depend on the time window of exposition and/or the different sensibility to the hormone of distinct neurogenic niches.

The neuroprotective effect of high intake of  $1,25(\text{OH})_2\text{D}_3$  has been confirmed in some AD trials, but not in others [88]. It is not clear whether hypovitaminosis D triggers AD or it removes protection in the Central nervous system (CNS) against AD. However, the combination of antineurodegenerative drugs with  $1,25(\text{OH})_2\text{D}_3$  supplementation might be useful. Indeed, the supplementation of the combination nemantidine plus  $1,25(\text{OH})_2\text{D}_3$  has been shown to prevent cognitive decline more efficiently than that of the single compounds [96].


#### 4. Sphingolipids

Sphingolipids (SLs) have long been regarded as inactive and stable structural components of the membrane, but they are biologically active molecules [11–14, 17, 19]. They are formed by the attachment of different polar headgroups at the primary alcohol group of a ceramide molecule. As reported in **Figure 1A** and **B**, there are two main pathways of Cer production: (1) *de novo* biosynthesis and (2) hydrolysis of sphingolipids, such as Sphingomyelin (SM). Which of these pathways dominates depends on the cell type, stimulus, and developmental stage of the cell [97].

SLs are the main components of specific membrane platforms that function in membrane signaling and trafficking, named lipid rafts [98, 99]. The most studied SLs are the bioactive lipid S1P and Cer [14], even if in the last few years relevant role has been demonstrated also for ceramide 1-phosphate (C1P) [18, 99, 100] and Cer species [101].

S1P plays a crucial physiological function. In fact, it is present at different concentrations in plasma as well in tissues, constituting a gradient that drives the trafficking of various immune cells [102]. The S1P gradient is due to the constitutive activity of Sphingosine-1-phosphate lyase (SPL) in the cells, leading to a low concentration of S1P in most tissues (0.5–75 pmol/mg) and low-micromolar range in plasma (0.2–0.9  $\mu\text{M}$ ) where S1P, produced and released by platelets and erythrocytes, is complexed with albumin and lipoproteins, particularly High-density lipoprotein (HDL) [103]. Of note, it has been proposed that cell fate is regulated by the ratio between S1P and Cer/Sphingosine (Sph) described the first time by Cuivillier et al. [20]. (**Figure 1C**) S1P enhances cell growth and survival, whereas its precursors, Cer and Sph, are generally associated with cell growth arrest and death. However, recent results indicate that Cer species (short or long chain Cer) have different functions [104].

S1P has a peculiar mechanism of action. In fact, S1P, produced inside the cell, can act as intracellular mediator or can be exported outside by the putative transporter Spnster 2, spns2 [105, 106]. Outside the cells, S1P can act as ligand of specific G-protein-coupled receptors. Five



**Figure 1.** Schematic representation of sphingolipid metabolism pathways and rheostat model. (A) Cers are produced from Sphingomyelin (SM) by activation of sphingomyelinase (SMase/SMPD) that produces Cer and phosphocholine. The reverse reaction is catalyzed by SM synthase (SMS). Ceramides can also be phosphorylated by ceramide kinase (CerK) to form ceramide-1-phosphate (C1P). S1P is derived by Cer deacylation by ceramidases (CDase) to Sph followed by its phosphorylation by Sph kinases (SphK). The degradation of S1P requires S1P phosphatases activity that produces Sph, and by S1P lyase (Spl) that produces hexadecenal and phosphoethanolamine. (B) *De novo* synthesis of SLs. The *de novo* synthesis is initiated at the cytosolic membranes of the endoplasmic reticulum and in mitochondria, via the condensation of L-serine with palmitoyl-CoA to form 3-keto-dihydrospingosine, and it is catalyzed by serine palmitoyltransferase (SPT). The product of SPT is reduced by a hydroxyl by 3-keto-dihydrospingosine reductase in a NADPH-dependent manner producing dihydrospingosine that is subsequently N-acylated to dihydroceramide (dhCer) by ceramide synthases (CerS). In mammals, CerS is encoded by six distinct genes, and each enzyme has a distinct, but overlapping acyl CoA preference. Dihydroceramide desaturase converts the DHSph backbone within Cer into Sph. (C) Sphingolipids rheostat model.

S1P specific named EDG/S1PR<sub>1-5</sub> have been described to mediate S1P signaling [11, 13, 14, 19]. S1P<sub>1</sub>, S1P<sub>2</sub>, and S1P<sub>3</sub> show broad tissue gene expression, while S1P<sub>4</sub> shows gene expression primarily in immune system cells, and S1P<sub>5</sub> is primarily expressed in the spleen (natural killer cells and other lymphocytes) and central nervous system [107]. Regarding the intracellular S1P effects, the bioactive lipid induces calcium release from the ER, alters the function of intracellular proteins, such as E3 ligase activity of TNF receptor associated factor 2 (TRAF2), binds the mitochondrial protein, and regulates mitochondrial assembly and function. When produced in the nuclei, S1P modulates gene expression inhibiting histone deacetylases [107].

#### 4.1. Sphingolipids in bone physiology

Various SLs play a crucial role in the development of normal skeletogenesis acting in three different skeletal cell types: chondrocytes in cartilage and osteoblasts and osteoclasts in bone. Abnormal tissue development of genetically modified animal models, such as mouse lacking

sphingomyelin phosphatase 3 (SMP3) also named sphingomyelinase 2 (SMase2), have been reported [108]. As described in the first part of this chapter, SMase are a family of different isoenzymes. Among them, the nSMSase2/SMPD3 is largely expressed in bone and cartilage, and it is involved in Cer-mediated signaling events [109]. In other studies, the role of SLs in bone have been performed by using synthetic analogues of ceramide (i.e., C2-Cer), specific inhibitors of the rate limiting enzyme (i.e., SphK), and agonists and antagonists that mimic or inhibit, respectively, the function of S1P, as ligand, of specific receptor subtypes. In this paragraph, we examine the involvement of Sph/S1P and Cer/C1P axis in the remodelling and physiology of bone.

#### 4.1.1. S1P—sphingosine

Several studies demonstrate that osteoblasts at different stages of differentiation may respond differently to SLs [110]. S1P has been shown to promote the proliferation of rat primary chondrocytes, whereas in preosteoblast MC3T3-E1, it negatively regulates the synthesis of the osteoblast marker osteocalcin [111, 112]. Notably, S1P can be released by osteoclasts [113] and secreted S1P can promote bone formation by enhancing the differentiation of osteoblast precursors to functional osteoblasts and by the recruitment of MSCs leading to high bone mass phenotype [114]. Recently, Keller et al. [115] report that the release of S1P by osteoclasts is regulated by calcitonin. S1P induces the upregulation of osteopontin and osteoblast differentiation markers in two osteoblast-like cell lines promoting the translocation of  $\beta$ -catenin [116].

In the same cell type, Sph and S1P induce intracellular calcium release [117, 118] and in rat osteoblasts and in human osteosarcoma cells; both sphingoids prevent the apoptotic process elicited by serum deprivation [119]. A functional cross talk exists between S1P and platelet-derived growth factor (PDGF) signaling. In fact, S1P limits, whereas PDGF promotes the migration of preosteoblasts, and it is the balance between these two bioactive molecules that allow only the differentiated osteoblasts to reach the site of bone formation. Very recently, it has been confirmed that in humans the detrimental effects of S1P on bone metabolism depend on the S1P gradient between blood and bone marrow cavity and on S1P receptor subtypes, mainly S1P1 and S1P2, play a crucial role in this control [112, 120]. In particular, S1P1 exerts positive chemotaxis action on an S1P gradient, whereas S1P2 counteracts this positive effect [121]. The deletion of S1P2 led to moderate osteopetrosis by affecting the homing of osteoclast precursors into bone [112, 120]. Higher circulating S1P levels are associated with lower bone mineral density, higher levels of bone resorption markers, and higher prevalence and severity of osteoporotic vertebral fracture in Koreans [122]. Therefore, S1P appears to be an important osteoclast-derived anabolic factor that couples bone resorption to bone formation, but it preferentially influences bone resorption rather than bone formation in humans.

Notably, increase in the secretion of BMP-2, OPN, and osteocalcin as well as highest extracellular matrix mineralization and osteonodules formation were observed when MSCs are cultured on thin titania dioxide coatings ( $\text{TiO}_2$ ) on stainless steel substrate doped with S1P [ $\text{TiO}_2/\text{S1P}(\text{CH}_3)$ ] [123]. Therefore, S1P, also in an appropriate combination with other sphingoid, such as C1P, may find wide application in regenerative medicine, particularly in bone regeneration with the use of MSCs [123].

#### 4.1.2. C1P—ceramide

Abnormal cartilage development and bone mineralization defects are observed in animal models in which the enzyme inactivation is generated by gene targeting (Smpd3<sup>-/-</sup> model) or chemically induced deletion in the Smpd3 locus (fro/fro model) [124, 125]. Several evidence underlines the importance of phosphocholine in bone mineralization [126]; however, at present no experimental data directly links Cer or choline to Extracellular matrix (ECM) mineralization. Studies in animal models are in some way in conflict with studies performed in cell culture: the inhibition of nSMase2/SMPD3 by GW4869 accelerates the mineralization of chondrogenic ATDC5 cultures [127], whereas either the fro/fro mice or mice lacking choline kinase presents an expansion of hypertrophic zone likely due to a delay in apoptosis of hypertrophic chondrocytes [128]. The role of Cer in bone tissue has been examined by other approaches leading also to conflicting results, such as short-chain synthetic cell-permeable forms of Cer, preferentially C-2 Cer [129]. In fact, it is not clear whether exogenous Cer has the same cellular targets as the endogenous species and the dose used in cell culture may not relate to the endogenous levels found under physiological or pathological conditions. Therefore, it is reported that Cer may be able to promote cell apoptosis or cell survival, and it may depend on the dose and cell type used. The treatment of cells with high-dose of C-2 Cer (1–10  $\mu$ M) leads to apoptosis of mouse primary osteoblasts. Contrary, the low-dose treatment (10 time less or more) of C-2 Cer promotes an antiapoptotic effect in other cells [130]. Regarding cell types, C-2 Cer treatment leads to increased apoptosis in both osteoblasts and chondrocytes, while there was no effect on the apoptosis of rabbit osteoclasts [131]. Recently, it has been demonstrated that short-chain C-6 Cer induces anti-osteosarcoma activity in vitro and in vivo [132]. In other studies, Cer has been shown to be mitogenic in preosteoblast MC3T3-E1 cells [133]. Recently, Cer species, in particular long-chain Cer (C-22 and C-24 Cer), have been demonstrated to mediate the proapoptotic effect of sodium nitroprusside, a nitric oxide donor, in MC3T3-E1 cells [134], suggesting that the controversial results may also be due to different levels of Cer species. Detectable increase of endogenous Cer levels is observed when cells are induced to apoptosis by TNF- $\alpha$  [135]. This upregulation leads to the modulation of NF- $\kappa$ B localization and function [136]. Different intracellular signaling pathways are involved in osteoblast death and survival: Cer induces osteoblast apoptosis through protein phosphatase 1 and protein kinase C (PKC)  $\delta$  [137, 138], whereas the ability of Cer to promote osteoblast survival is prevented by PKC  $\zeta$  inhibitor. At present, it is not clear whether Cer acts up- or downstream to or independently of caspases [139]. Interestingly, in mice lacking the enzyme that converts dihydroceramide to ceramide (dihydroceramide desaturase 1) Cer reduction does not lead to bone defects, indicating that the maintenance of Cer levels is not essential for normal bone mineralization. Recently, it has been found a role also for C1P in the osteogenesis of multipotent stromal cells derived from MSCs. C1P can affect the growth and expanded intercellular connections, thereby, increasing the proliferative activity, acting in opposite manner of S1P. Therefore, it has been suggested that an appropriate combination of C1P and S1P may be a useful strategy in bone regeneration with the use of MSCs [123].

#### 4.2. Sphingolipids in nervous system physiology

SLs are particularly abundant in the central nervous system. Mutations in genes coding for enzymes involved in their metabolism cause sphingolipidosis many of which show alterations



in the nervous system (reviewed by Sabourdy et al. [140]). In addition, modifications in SL metabolism are found in neurodegenerative diseases, such as AD, PD, Huntington disease (HD), MS, and major depression [141–146]. Various SLs are crucial in regulating neural physiological functions, including cell survival, apoptosis, differentiation, inflammation, excitability, and neurotransmitter release [15, 137–149].

#### 4.2.1. S1P—sphingosine

Like Cer, Sph acts as a proapoptotic signal as well as an inhibitor of several enzymes such as protein kinase C (PKC), phospholipase D, and of the transcription factor SF-1 [15, 150]. Sph also directly modulates voltage-activated calcium channels in pituitary cells and several components of the melastatin-like transient receptor potential channel subfamily, such as Transient receptor potential cation channel subfamily M member 3 (TRPM3) [151, 152]. Sph alters the integrity of membranes and induces the release of lysosomal cysteine proteases, such as cathepsins and of cytochrome *c*, which in turn activates the intrinsic pathway of apoptosis. Increasing evidence indicates that Sph regulates vesicle fusion and trafficking and, therefore, the strength and reliability of synaptic transmission [153, 154]. In addition, Sph is a competitive antagonist of the type 1 cannabinoid receptors (CB1Rs) [155] and possibly it contributes to SL regulation of nociception [156]. S1P modulates survival, proliferation, differentiation, cell migration, calcium homeostasis, neurite retraction, angiogenic vascular maturation, and cytoskeleton dynamics [13, 107, 149]. The bioactive lipid can also induce neuroprotection through many mechanisms that include production of growth factors, decrease of oxidative stress, increase of Mitogen-activated protein kinase (MAPK) activation, activation of PI3K/AKT pathway, modulation of antiapoptotic proteins, pigenomic effects by direct inhibition of deacetylases, and affecting mitochondrial functionality (for a recent review see Ref. [149].

S1P receptors are expressed in CNS cells (neurons, oligodendrocytes, astrocytes, and microglia) and their expression levels change during development. The role of S1PR depends on cell types and on their different expression and localization during development or following stimulation. The studies of knockout mice with deficits in both Sphk1 and Sphk2 highlight the importance of S1P in the development of the nervous system. These mice show severely disturbed neurogenesis, including neural tube closure and angiogenesis, and they die at an early embryonic stage [157]. Migration of neural stem progenitor cells toward injury sites is promoted by S1P via S1P1 [158] and inhibited on the oligodendrocyte progenitor cells via S1P5 [159]. S1P receptors are expressed in CNS cells (neurons, oligodendrocytes, astrocytes, and microglia) and their expression levels change during development. Analogously to the double SphK knockout, the S1P1 knockout mouse shows altered neurogenesis and angiogenesis, while the knockout of the other receptors have less severe consequences. MacLennan et al. [160] reported that mice lacking S1P2 show significant increase in excitatory postsynaptic potentiation, resulting in spontaneous seizures. The role of S1PR depends on their different expression and localization during development or following stimulation. For example, activation of S1P5 causes activation of Rho and retraction of processes in immature oligodendrocytes and survival in mature cells [161]. NGF induces differentiation in PC12 cells through a relocation of S1P receptors; S1P1 that induces neurite growth is expressed at the plasma membrane, while S1P2 is internalized [162] becoming unable to cause cell rounding and loss of neurites. S1P and neurotrophic factors have mutual effects on expression of each other:



NGF and GDNF are able to stimulate S1P generation, and S1P increases GDNF production by astrocytes [144]. S1P plays different roles in synaptic transmission. It increases glutamate release in hippocampus via S1P3 [163], it enhances excitability in rat sensory neurons through S1P1 and S1P3 [155, 158, 164], and it is involved in the recruitment of vesicles in the presynaptic membrane [165] and also in endocytic membrane trafficking [166].

#### 4.2.2. C1P—ceramide

Cer regulates cell growth, differentiation, apoptosis, inflammation, exosome release, and neural excitability [15, 157, 158, 167, 168]. Accumulating evidence indicates that different Cer species might have different functions [169]. Cer 18:0 is synthesized by CerS1, an enzyme abundant in the brain, and appears to have a protective role [167, 168]. Serum deprivation that increases apoptosis in embryonic hippocampal cells increases Cer 16:0 and decreases Cer 24:0 content [27]. Cer has been involved in synaptic regulation and plasticity [170, 171]. For example, Cer is able to increase dopamine release and uptake [172] and to modulate excitatory postsynaptic currents by controlling the insertion and clustering of NMDA receptors [173]. Moreover, SMase2 inhibition delays formation of spatial memory in mice [174]. Regarding inflammation, astrocytes display increased Cer following ischemia/reperfusion leading to generation of pro-inflammatory cytokines [175]. Recently, it has been reported that Cer induces ciliogenesis, a critical step in differentiation in embryonic stem cells and neural progenitors [176]. CerK, the enzyme generating C1P, was first observed in brain synaptic vesicles [177] and found to be highly expressed in brain [178] suggesting a role of C1P in neurotransmitter release. C1P induces proliferation or survival in several types of cells including macrophages and fibroblasts [179] while inhibition or downregulation of CerK decreases proliferation in human neuroblastoma cells [18]. C1P plays a regulatory role in inflammation since it directly binds and activates  $\alpha$ -type cytosolic phospholipase A2 stimulating arachidonic acid release [180]. Recently, it has been demonstrated that C1P plays an important role in recruitment of stem/progenitor cells to damaged organs [181]. Whether C1P is also released from the injured nervous system or whether it induces migration in nervous stem cells is unknown.

## 5. Cross talk between vitamin D and sphingolipid metabolism: a potential role in Alzheimer's disease

1,25(OH)<sub>2</sub>D<sub>3</sub> and SL metabolism cross talk at different levels. For example, 1,25(OH)<sub>2</sub>D<sub>3</sub> modulates the expression of genes involved in S1P degradation, such as Sphingosine-1-phosphate phosphatase 2 (SGPP2) [182] and of growth factors involved in differentiation and neuroprotection, such as NT-3, Brain-derived neurotrophic factor (BDNF) and ciliary neurotrophic factor (CNTF) [3, 4]. On the other hand, many neuroprotective actions of 1,25(OH)<sub>2</sub>D<sub>3</sub> have been reported to be due to stimulation of SphK and increased levels of S1P [183, 184]. In addition, 1,25(OH)<sub>2</sub>D<sub>3</sub> also is able to decrease the expression of CerK, the enzyme that generates C1P [18]. C1P activates directly Secretory PLA2 enzymes (sPLA2) [180], producing arachidonic acid that can be further metabolized to proinflammatory mediators. C1P and S1P have a crucial role in migration. C1P is released from damaged cells and chemoattracts bone marrow-derived

multipotent stromal cells, endothelial progenitor cells, and very small embryonic-like stem cell. The migration of osteoclast precursors is controlled by S1P and, recently, it has been found that  $1,25(\text{OH})_2\text{D}_3$  reduces the expression of the chemorepulsive receptor S1P2 on circulating precursors [185]. AD is a neurodegenerative disorder of the central nervous system and the most common form of dementia [186]. The pathogenic hallmarks of AD include extracellular amyloid-containing plaques, intracellular neurofibrillary tangles consisting of hyperphosphorylated tau protein and death of cholinergic neurons of the basal forebrain. Amyloid plaques are mainly formed by aggregated amyloid beta peptide ( $\text{A}\beta$ ). Alterations in the enzymes involved in SL metabolism and content have been observed in brains and cerebrospinal fluid of AD patients [187–192], leading to an increase of Cer and loss of S1P. Many studies in culture cells and animal models have demonstrated that  $\text{A}\beta$  affects SL metabolism. For example,  $\text{A}\beta_{42}$  directly binds and activates nSMase in vitro decreasing SM content [193]. In addition,  $\text{A}\beta$  can activate both neutral and acidic SMases through increased ROS accumulation via Nicotinamide adenine dinucleotide phosphate (NADPH) oxidase activation and glutathione (GSH) depletion [194]. Increased Cer could then induce apoptosis [186, 195, 196]. In addition, Cer increases stability of  $\beta$ -amyloid precursor protein cleaving enzyme (BACE1) activity [197], while S1P binds to and increases proteolytic activity of BACE1 [189] and SM decreases  $\text{A}\beta$  production by inhibition of the  $\gamma$ -secretase [193]. Summarizing, some SLs might be protective by enhancing  $\text{A}\beta$  clearance or decreasing  $\text{A}\beta$  production, others increase  $\text{A}\beta$  toxicity or  $\text{A}\beta$  oligomerization and at the same time, amyloid precursor protein processing also affects lipid metabolism, resulting in complex regulatory feed-back cycles, which appear to be dysregulated in AD. It is worth noting that recently, a novel mechanism of Cer-enriched exosomes released by  $\text{A}\beta$ -treated astrocytes has been proposed to be responsible for  $\text{A}\beta$ -induced apoptosis [167].

The neuroprotective action of SL analogues, such as FTY720, has been tested in some neurodegenerative diseases. FTY720 is a prodrug that is converted to an analogue of S1P when phosphorylated by SphK. It has been approved for the treatment of multiple sclerosis (MS) and acts both as an immunomodulatory drug and on different cells of the CNS (neurons, astrocytes, oligodendrocytes, and microglia), all of which express S1P receptors [198]. Administration of FTY720 in a rat model of AD obtained by injection of  $\text{A}\beta$  decreases death in hippocampus and cortex and increases memory compared with control rats [199, 200]. FTY720 decreases production of  $\text{A}\beta$  in cultured neuronal cells [189]. Increasing evidence derived from epidemiological studies indicate that  $1,25(\text{OH})_2\text{D}_3$  deficiency and VDR polymorphisms influence susceptibility to AD [201], whereas  $\text{A}\beta$  may disrupt the hormone-VDR pathway and cause defective utilization of  $1,25(\text{OH})_2\text{D}_3$  by suppressing the level of the VDR and elevating the level of 24OHase [202]. In addition to neuroprotective effects involving calcium, Reactive Oxygen Species (ROS), and inflammation,  $1,25(\text{OH})_2\text{D}_3$  is able to exert other specific effects important for AD, by regulating the expression of many AD-related genes. It attenuates  $\text{A}\beta$  peptide accumulation by stimulating phagocytosis of  $\text{A}\beta$  peptide probably by modulating transcription of Toll-like receptors and cytokines together with enhancing brain-to-blood efflux transport by increasing P-glycoprotein expression and likely by altering Amyloid Precursor Protein (APP) processing [88] and prevents the acetylcholine defect by increasing the activity of choline acetyltransferase (thus acetylcholine synthesis) in the brain [203]. Part of VDR is also located in lipid microdomains in the nuclear membrane [2], and

this localization is modified by altering SL metabolism and has been associated with embryonic hippocampal cell differentiation. Neuroprotective actions of SLs, in particular S1P, and  $1,25(\text{OH})_2\text{D}_3$  include many common effectors such as calcium regulation, synaptic modulation, growth factor expression, regulation of inflammation, etc., but whether  $1,25(\text{OH})_2\text{D}_3$  and SLs, in particular S1P, could act synergistically on neuroprotection and/or neurogenesis in AD is still unknown and deserves further investigation. A study in our lab indicates that the cross talk between SLs and  $1,25(\text{OH})_2\text{D}_3$  leads to a specific balance between neurodegeneration/neuroprotection in neuronal cells [204]. The neuroprotective effect of high intake of  $1,25(\text{OH})_2\text{D}_3$  has been found in some AD trials, but not in others [88]. It is not clear whether hypovitaminosis D triggers AD or it removes protection in the CNS against AD. However, the combination of antineurodegenerative drugs with  $1,25(\text{OH})_2\text{D}_3$  supplementation might be useful. In fact, the supplementation with nemantidine plus  $1,25(\text{OH})_2\text{D}_3$  has been shown to prevent cognitive decline more efficiently than that with the single compounds [96]. Regarding MS, very limited evidence suggests a potential benefit of  $1,25(\text{OH})_2\text{D}_3$  supplementation for the prevention of MS and this needs to be further verified by future studies [3].

## Acknowledgements

This work was supported by grants from Fondazione Cassa di Risparmio di Pistoia e Pescia, Regione Toscana to E.M., MIUR (ex 60%) to EM and MM and from A.I.L. for contributing to AF fellowship. The authors are grateful to Dr. Maria Martinesi and Dr. Elena Bruzzone from the Department of Experimental and Clinical Biomedical Sciences for their contribution.

## Author details

Alessia Frati<sup>1,2</sup>, Mercedes Garcia Gil<sup>3,4</sup>, Federica Pierucci<sup>1,2</sup> and Elisabetta Meacci<sup>1,2\*</sup>

\*Address all correspondence to: [elisabetta.meacci@unifi.it](mailto:elisabetta.meacci@unifi.it)

1 Department of Experimental and Clinical Biomedical Sciences, Molecular and Applied Biology Research Unit, University of Firenze, Firenze, Italy

2 Interuniversity Institute of Myology, Italy

3 Department of Biology, University of Pisa, Pisa, Italy

4 Interdepartmental Research Center Nutrafood "Nutraceuticals and Food for Health," University of Pisa, Pisa, Italy

## References

- [1] Marini F, Bartoccini E, Cascianelli G, Voccoli V, Bavaglia MG, Magni MV, Garcia-Gil M, Albi E. Effect of  $1\alpha,25$ -dihydroxyvitamin D<sub>3</sub> in embryonic hippocampal cells. *Hippocampus*. 2010 Jun;20(6):696–705. doi: 10.1002/hipo.20670.

- [2] Bartoccini E, Marini F, Damaskopoulou E, Lazzarini R, Cataldi S, Cascianelli G, Gil Garcia M, Albi E. Nuclear lipid microdomains regulate nuclear vitamin D3 uptake and influence embryonic hippocampal cell differentiation. *Mol Biol Cell*. 2011 Sep;22(17):3022–31. doi: 10.1091/mbc.E11-03-0196.
- [3] DeLuca GC, Kimball SM, Kolasinski J, Ramagopalan SV, Ebers GC. Review: the role of vitamin D in nervous system health and disease. *Neuropathol Appl Neurobiol*. 2013 Aug;39(5):458–84. doi: 10.1111/nan.12020.
- [4] Shirazi HA, Rasouli J, Ciric B, Rostami A, Zhang GX. 1,25-Dihydroxyvitamin D3 enhances neural stem cell proliferation and oligodendrocyte differentiation. *Exp Mol Pathol*. 2015 Apr;98(2):240–5. doi: 10.1016/j.yexmp.2015.02.004.
- [5] Chun RF et al. Vitamin D supplementation and antibacterial immune responses in adolescents and young adults with HIV/AIDS. *J Steroid Biochem Mol Biol*. 2015 Apr;148:290–7. doi: 10.1016/j.jsbmb.2014.07.013. Epub 2014 Aug 1.
- [6] DeLuca HF. Overview of general physiologic features and functions of vitamin D. *Am J Clin Nutr*. 2004 Dec;80(6 Suppl):1689S–96S.
- [7] Christakos S, Dhawan P, Verstuyf A, Verlinden L, Carmeliet G. Vitamin D: metabolism, molecular mechanism of action, and pleiotropic effects. *Physiol Rev*. 2016 Jan;96(1):365–408. doi: 10.1152/physrev.00014.2015.
- [8] Trump DL, Deeb KK, Johnson CS. Vitamin D: considerations in the continued development as an agent for cancer prevention and therapy. *Cancer J*. 2010 Jan-Feb;16(1):1–9. doi: 10.1097/PPO.0b013e3181c51ee6.
- [9] Masuda S, Jones G. Promise of vitamin D analogues in the treatment of hyperproliferative conditions. *Mol Cancer Ther*. 2006 Apr;5(4):797–808.
- [10] Zügel U, Steinmeyer A, Giesen C, Asadullah K. A novel immunosuppressive 1 $\alpha$ ,25-dihydroxyvitamin D3 analog with reduced hypercalcemic activity. *J Invest Dermatol*. 2002 Dec;119(6):1434–42.
- [11] Spiegel S, Milstien S. Sphingosine 1-phosphate, a key cell signaling molecule. *J Biol Chem*. 2002 Jul 19;277(29):25851–4. doi: 10.1038/nrm1103.
- [12] Gault CR, Obeid LM, Hannun YA. An overview of sphingolipid metabolism: from synthesis to breakdown. *Adv Exp Med Biol*. 2010;688:1–23.
- [13] Maceyka M, Harikumar KB, Milstien S, Spiegel S. Sphingosine-1-phosphate signaling and its role in disease. *Trends Cell Biol*. 2012 Jan;22(1):50–60. doi: 10.1016/j.tcb.2011.09.003.
- [14] Newton J, Lima S, Maceyka M, Spiegel S. Revisiting the sphingolipid rheostat: evolving concepts in cancer therapy. *Exp Cell Res*. 2015 May 1;333(2):195–200. doi: 10.1016/j.yexcr.2015.02.025.
- [15] Colombaioni L, Garcia-Gil M. Sphingolipid metabolites in neural signalling and function. *Brain Res Brain Res Rev*. 2004 Nov;46(3):328–55.

- [16] Bartke N, Hannun YA. Bioactive sphingolipids: metabolism and function. *J Lipid Res.* 2009 Apr;50 Suppl:S91–6. doi: 10.1194/jlr.R800080-JLR200.
- [17] Fyrst H, Saba JD. An update on sphingosine-1-phosphate and other sphingolipid mediators. *Nat Chem Biol.* 2010 Jul;6(7):489–97. doi: 10.1038/nchembio.392.
- [18] Bini F, Frati A, Garcia-Gil M, Caleo M, Peretto P, Gomez-Muñoz A, Meacci E. New signalling pathway involved in the anti-proliferative action of vitamin D3 and its analogues in human neuroblastoma cells. A role for ceramide kinase. *Neuropharmacology.* 2012 Sep;63(4):524–37. doi: 10.1016/j.neuropharm.2012.04.026.
- [19] Pyne NJ et al. Role of sphingosine 1-phosphate receptors, sphingosine kinases and sphingosine in cancer and inflammation. *Adv Biol Regul.* 2016 Jan;60:151–9. doi: 10.1016/j.jbior.2015.09.001.
- [20] Cuvillier O, Pirianov G, Kleuser B, Vanek PG, Coso OA, Gutkind S, Spiegel S. Suppression of ceramide-mediated programmed cell death by sphingosine-1-phosphate. *Nature.* 1996 Jun 27;381(6585):800–3. doi: 10.1038/381800a0.
- [21] Catarzi S, Giannoni E, Meacci E, Iantomasi T, Vincenzini MT. Sphingosine 1-phosphate stimulation of NADPH oxidase activity: relationship with platelet-derived growth factor receptor and c-Src kinase. *Biochim Biophys Acta.* 2007 Jun;1770(6):872–83.
- [22] Meacci E, Bini F, Sassoli C, Martinesi M, Squecco R, Chellini F, Zecchi S, Francini F, Formigli L. Functional interaction between TRPC1 channel and connexin-43 protein: a novel pathway underlying S1P action on skeletal myogenesis. *Cell Mol Life Sci.* 2010 Dec;67(24):4269–85. doi: 10.1007/s00018-010-0442-3.
- [23] Sassoli C, Formigli L, Bini F, Squecco R, Battistini C, Zecchi S, Francini F, Meacci E. Effects of S1P on skeletal muscle repair/regeneration during eccentric contraction. *J Cell Mol Med.* 2011 Nov;15(11):2498–511. doi: 10.1111/j.1582-4934.2010.01250.x.
- [24] Tavarini S, Colombaioni L, Garcia-Gil M. Sphingomyelinase metabolites control survival and apoptotic death in SH-SY5Y neuroblastoma cells. *Neurosci Lett.* 2000 May 19;285:185–8.
- [25] Colombaioni L, Frago LM, Varela-Nieto I, Pesi R, Garcia-Gil M. Serum deprivation increases ceramide levels and induces apoptosis in undifferentiated HN9.10e cells. *Neurochem Int.* 2002 Apr;40(4):327–36.
- [26] Albi E et al. Nuclear sphingomyelin pathway in serum deprivation-induced apoptosis of embryonic hippocampal cells. *J Cell Physiol.* 2006 Jan;206(1):189–95.
- [27] Garcia-Gil M, Lazzarini A, Lazzarini R, Floridi E, Cataldi S, Floridi A, Albi E. Serum deprivation alters lipid profile in HN9.10e embryonic hippocampal cells. *Neurosci Lett.* 2015 Mar 4;589:83–7. doi: 10.1016/j.neulet.2014.12.059.
- [28] Bornancin F. Ceramide kinase: the first decade. *Cell Signal.* 2011 Jun;23(6):999–1008. doi: 10.1016/j.cellsig.2010.11.012. Epub 2010 Nov 25.

- [29] Frati A, Ricci B, Pierucci F, Nistri S, Bani D, Meacci E. Role of sphingosine kinase/S1P axis in ECM remodeling of cardiac cells elicited by relaxin. *Mol Endocrinol*. 2015 Jan;29(1):53–67. doi: 10.1210/me.2014-1201.
- [30] Sassoli C, Frati A, Tani A, Anderloni G, Pierucci F, Zecchi Orlandini S, Formigli L, Meacci E. Mesenchymal stromal cell secreted sphingosine 1-phosphate (S1P) exerts a stimulatory effect on skeletal myoblast proliferation. *PLoS One*. 2014 Sep 29;9(9):e108662. doi: 10.1371/journal.pone.0108662.
- [31] Jilka RL. Cytokines, bone remodeling, and estrogen deficiency: a 1998 update. *Bone*. 1998 Aug;23(2):75–81.
- [32] Weinstein RS, Manolagas SC. Apoptosis and osteoporosis. *Am J Med*. 2000;108:153–64.
- [33] Nakamura I, Takahashi N, Jimi E, Udagawa N, Suda T. Regulation of osteoclast function. *Mod Rheumatol*. 2012 Apr;22(2):167–77. doi: 10.1007/s10165-011-0530-8.
- [34] Väänänen HK, Laitala-Leinonen T. Osteoclast lineage and function. *Arch Biochem Biophys*. 2008 May 15;473(2):132–8. doi: 10.1016/j.abb.2008.03.037. Epub 2008 Apr 6.
- [35] Yagi M et al. DC-STAMP is essential for cell-cell fusion in osteoclasts and foreign body giant cells. *J Exp Med*. 2005 Aug 1;202(3):345–51. PubMed PMID: 16061724; PubMed Central PMCID: PMC2213087.
- [36] Miyamoto H et al. Osteoclast stimulatory transmembrane protein and dendritic cell-specific transmembrane protein cooperatively modulate cell–cell fusion to form osteoclasts and foreign body giant cells. *J Bone Miner Res*. 2012 Jun;27(6):1289–97. doi: 10.1002/jbmr.1575. PubMed PMID: 22337159.
- [37] Nijenhuis T, van der Eerden BC, Zügel U, Steinmeyer A, Weinans H, Hoenderop JG, van Leeuwen JP, Bindels RJ. The novel vitamin D analog ZK191784 as an intestine-specific vitamin D antagonist. *FASEB J*. 2006 Oct;20(12):2171–3. PubMed PMID: 17012263.
- [38] Rodan GA, Martin TJ. Role of osteoblasts in hormonal control of bone resorption—a hypothesis. *Calcif Tissue Int*. 1982 May;34(3):311. PubMed PMID: 6809295.
- [39] Takahashi N, Akatsu T, Udagawa N, Sasaki T, Yamaguchi A, Moseley JM, Martin TJ, Suda T. Osteoblastic cells are involved in osteoclast formation. *Endocrinology*. 1988 Nov;123(5):2600–2. PubMed PMID: 2844518.
- [40] Rucci N, Teti A. The “love–hate” relationship between osteoclasts and bone matrix. *Matrix Biol*. 2016 May–Jul;52–54:176–90. doi: 10.1016/j.matbio.2016.02.009. Review.
- [41] Roodman GD, Ibbotson KJ, MacDonald BR, Kuehl TJ, Mundy GR. 1,25-Dihydroxyvitamin D3 causes formation of multinucleated cells with several osteoclast characteristics in cultures of primate marrow. *Proc Natl Acad Sci U S A*. 1985 Dec;82(23):8213–7. PubMed PMID: 3865222.
- [42] Miller WL. Genetic disorders of vitamin D biosynthesis and degradation. *J Steroid Biochem Mol Biol*. 2016 April 6;S0960-0760(16):30097–8.



- [43] Pike JW, Lee SM, Meyer MB. Regulation of gene expression by 1,25-dihydroxyvitamin D3 in bone cells: exploiting new approaches and defining new mechanisms. *Bonekey Rep.* 2014 Jan 8;3:482. doi: 10.1038/bonekey.2013.216. eCollection 2014 Jan 8. Review. PubMed PMID: 24466413; PubMed Central PMCID: PMC3899561.
- [44] Bouillon R, Carmeliet G, Verlinden L, van Etten E, Verstuyf A, Luderer HF, Lieben L, Mathieu C, Demay M. Vitamin D and human health: lessons from vitamin D receptor null mice. *Endocr Rev.* 2008 Oct;29(6):726–76. doi: 10.1210/er.2008-0004. Epub 2008 Aug 11. Review. PubMed PMID: 18694980; PubMed Central PMCID: PMC2583388.
- [45] Fujita H et al. Tight junction proteins claudin-2 and -12 are critical for vitamin D-dependent Ca<sup>2+</sup> absorption between enterocytes. *Mol Biol Cell.* 2008 May;19(5):1912–21. doi: 10.1091/mbc.E07-09-0973.
- [46] Van Cromphaut SJ et al. Duodenal calcium absorption in vitamin D receptor-knock-out mice: functional and molecular aspects. *Proc Natl Acad Sci U S A.* 2001 Nov 6;98(23):13324–9. Epub 2001 Oct 30. PubMed PMID: 11687634; PubMed Central PMCID: PMC60869.
- [47] Wong BR et al. TRANCE is a novel ligand of the tumor necrosis factor receptor family that activates c-Jun N-terminal kinase in T cells. *J Biol Chem.* 1997;272(40):25190–4.
- [48] Anderson DM et al. A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function. *Nature.* 1997 Nov 13;390(6656):175–9.
- [49] Lacey DL et al. Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell.* 1998 Apr 17;93(2):165–76. PubMed PMID: 9568710.
- [50] O'Connor KM. Evaluation and treatment of osteoporosis. *Med Clin North Am.* 2016 Jul;100(4):807–26. doi: 10.1016/j.mcna.2016.03.016. Epub 2016 Apr 20. Review.
- [51] Gallagher JC, Jerpbak CM, Jee WS, Johnson KA, DeLuca HF, Riggs BL. 1,25-Dihydroxyvitamin D3: short- and long-term effects on bone and calcium metabolism in patients with postmenopausal osteoporosis. *Proc Natl Acad Sci U S A.* 1982 May;79(10):3325–9.
- [52] Dechant KL, Goa KL. Calcitriol. A review of its use in the treatment of postmenopausal osteoporosis and its potential in corticosteroid-induced osteoporosis. *Drugs Aging.* 1994 Oct;5(4):300–17.
- [53] Gillespie LD, Robertson MC, Gillespie WJ, Sherrington C, Gates S, Clemson LM, Lamb SE. Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev.* 2012 Sep 12;9:1–18.
- [54] Glorieux FH, Pettifor JM. Vitamin D/dietary calcium deficiency rickets and pseudo-vitamin D deficiency rickets. *Bonekey Rep.* 2014 Mar 19;3:524. doi: 10.1038/bonekey.2014.19. Review.
- [55] Lips P, Gielen E, van Schoor NM. Vitamin D supplements with or without calcium to prevent fractures. *Bonekey Rep.* 2014 Mar 5;3:512. doi: 10.1038/bonekey.2014.7.

- [56] Schafer AL et al. Intestinal calcium absorption decreases dramatically after gastric bypass surgery despite optimization of vitamin D status. *J Bone Miner Res.* 2015 Aug;30(8):1377–85. doi: 10.1002/jbmr.2467.
- [57] Iuliano-Burns S, Ayton J, Hillam S, Jones G, King K, Macleod S, Seeman E. Skeletal and hormonal responses to vitamin D supplementation during sunlight deprivation in Antarctic expeditioners. *Osteoporos Int.* 2012 Oct;23(10):2461–7. doi: 10.1007/s00198-011-1858-9.
- [58] Leyssens C, Verlinden L, Verstuyf A. The future of vitamin D analogs. *Front Physiol.* 2014 Apr 3;5:122. doi: 10.3389/fphys.2014.00122. eCollection 2014. Review. PubMed PMID: 24772087; PubMed Central PMCID: PMC3982071.
- [59] Kubodera N, Tsuji N, Uchiyama Y, Endo K. A new active vitamin D analog, ED-71, causes increase in bone mass with preferential effects on bone in osteoporotic patients. *J Cell Biochem.* 2003 Feb 1;88(2):286–9. PubMed PMID: 12520528.
- [60] Uchiyama Y, Higuchi Y, Takeda S, Masaki T, Shira-Ishi A, Sato K, Kubodera N, Ikeda K, Ogata E. ED-71, a vitamin D analog, is a more potent inhibitor of bone reabsorption than alfacalcidol in an estrogen-deficient rat model of osteoporosis. *Bone.* 2002 Apr;30(4):582–8. PubMed PMID: 11934649.
- [61] Matsumoto T, Takano T, Yamakido S, Takahashi F, Tsuji N. Comparison of the effects of eldcalcitol and alfacalcidol on bone and calcium metabolism. *J Steroid Biochem Mol Biol.* 2010 Jul;121(1–2):261–4. doi: 10.1016/j.jsbmb.2010.03.035.
- [62] Strauch UG, Obermeier F, Grunwald N, Dunger N, Rath HC, Schölmerich J, Steinmeyer A, Zügel U, Herfarth HH. Calcitriol analog ZK191784 ameliorates acute and chronic dextran sodium sulfate-induced colitis by modulation of intestinal dendritic cell numbers and phenotype. *World J Gastroenterol.* 2007 Dec 28;13(48):6529–37.
- [63] Park MR, Lee JH, Park MS, Hwang JE, Shim HJ, Cho SH, Chung IJ, Bae WK. Suppressive effect of 19-nor-1 $\alpha$ -25-dihydroxyvitamin D2 on gastric cancer cells and peritoneal metastasis model. *J Korean Med Sci.* 2012 Sep;27(9):1037–43. doi: 10.3346/jkms.2012.27.9.1037. Epub 2012 Aug 22. PubMed PMID: 22969249.
- [64] Ma Y, Yu WD, Hidalgo AA, Luo W, Delansorne R, Johnson CS, Trump DL. Inecalcitol, an analog of 1,25D3, displays enhanced antitumor activity through the induction of apoptosis in a squamous cell carcinoma model system. *Cell Cycle.* 2013 Mar 1;12(5):743–52. doi: 10.4161/cc.23846. Epub 2013 Feb 6. PubMed PMID: 23388458.
- [65] Sundaram S, Sea A, Feldman S, Strawbridge R, Hoopes PJ, Demidenko E, Binderup L, Gewirtz DA. The combination of a potent vitamin D3 analog, EB 1089, with ionizing radiation reduces tumor growth and induces apoptosis of MCF-7 breast tumor xenografts in nude mice. *Clin Cancer Res.* 2003 Jun;9(6):2350–6. PubMed PMID: 12796405.
- [66] Komori T. Animal models for osteoporosis. *Eur J Pharmacol.* 2015 Jul 15;759:287–94. doi: 10.1016/j.ejphar.2015.03.028. Review.

- [67] Matsushita M et al. Age-related changes in bone mass in the senescence-accelerated mouse (SAM). *Am J Pathol.* 1986;125:276–83.
- [68] Aref M, Gallant MA, Organ JM, Wallace JM, Newman CL, Burr DB, Brown DM, Allen MR. In vivo reference point indentation reveals positive effects of raloxifene on mechanical properties following 6 months of treatment in skeletally mature beagle dogs. *Bone.* 2013 Oct;56(2):449–53. doi: 10.1016/j.bone.2013.07.009.
- [69] Shapiro IM, Hickok NJ, Parvizi J, Stewart S, Schaer TP. Molecular engineering of an orthopaedic implant: from bench to bedside. *Eur Cell Mater.* 2012 May 24;23:362–70. Review.
- [70] Weinstein RS, Jilka RL, Parfitt AM, Manolagas SC. Inhibition of osteoblastogenesis and promotion of apoptosis of osteoblasts and osteoclasts by glucocorticoids: potential mechanisms of their deleterious effects on bone. *J Clin Invest.* 1998;102:274–82.
- [71] Klotho and  $\beta$  Klotho. *Adv Exp Med Biol.* 2012;728:25–40. doi: 10.1007/978-1-4614-0887-1\_2
- [72] Rinotas V et al. Novel genetic models of osteoporosis by overexpression of human RANKL in transgenic mice. *J Bone Miner Res.* 2014;29(5):1158–69.
- [73] Renkema KY et al. Hypervitaminosis D mediates compensatory  $\text{Ca}^{2+}$  hyperabsorption in TRPV5 knockout mice. *J Am Soc Nephrol.* 2005 Nov;16(11):3188–95.
- [74] Bivi N, Condon KW, Allen MR, Farlow N, Passeri G, Brun LR, Rhee Y, Bellido T, Plotkin LI. Cell autonomous requirement of connexin 43 for osteocyte survival: consequences for endocortical resorption and periosteal bone formation. *J Bone Miner Res.* 2012 Feb;27(2):374–89. doi: 10.1002/jbmr.548.
- [75] Thierry-Palmer M, Cephas S. Plasma 25-hydroxycholecalciferol and 1,25-dihydroxycholecalciferol concentrations are decreased in hind limb unloaded Dahl salt-sensitive female rats. *J Steroid Biochem Mol Biol.* 2010 Feb 15;118(3):188–93. doi: 10.1016/j.jsbmb.2009.12.012. Epub 2010 Jan 4.
- [76] Kamoun-Goldrat AS, Le Merrer MF. Animal models of osteogenesis imperfecta and related syndromes. *J Bone Miner Metab.* 2007;25(4):211–8. Epub 2007 Jun 25.
- [77] Plante L, Veilleux LN, Glorieux FH, Weiler H, Rauch F. Effect of high-dose vitamin D supplementation on bone density in youth with osteogenesis imperfecta: a randomized controlled trial. *Bone.* 2016 May;86:36–42. doi: 10.1016/j.bone.2016.02.013.
- [78] Aubin I, Adams CP, Opsahl S, Septier D, Bishop CE, Auge N, Salvayre R, Negre-Salvayre A, Goldberg M, Guénet JL, Poirier C. A deletion in the gene encoding sphingomyelin phosphodiesterase 3 (*Smpd3*) results in osteogenesis and dentinogenesis imperfecta in the mouse. *Nat Genet.* 2005 Aug;37(8):803–5.
- [79] Patrick RP, Ames BN. Vitamin D hormone regulates serotonin synthesis. Part 1: relevance for autism. *FASEB J.* 2014 Jun;28(6):2398–413. doi: 10.1096/fj.13-246546.
- [80] Zanatta L et al.  $1\alpha,25$ -Dihydroxyvitamin D(3) mechanism of action: modulation of L-type calcium channels leading to calcium uptake and intermediate filament phosphorylation in cerebral cortex of young rats. *Biochim Biophys Acta.* 2012 Oct;1823(10):1708–19. doi: 10.1016/j.bbamcr.2012.06.023.

- [81] Eyles DW, Burne TH, McGrath JJ. Vitamin D, effects on brain development, adult brain function and the links between low levels of vitamin D and neuropsychiatric disease. *Front Neuroendocrinol.* 2013 Jan;34(1):47–64.
- [82] Jiang P et al. Association between vitamin D receptor gene polymorphisms with childhood temporal lobe epilepsy. *Int J Environ Res Public Health.* 2015;12(11):13913–22.
- [83] Fernandes de Abreu DA, Eyles D, Féron F. Vitamin D, a neuroimmunomodulator: implications for neurodegenerative and autoimmune diseases. *Psychoneuroendocrinology.* 2009 Dec;34 Suppl 1:S265–77. doi: 10.1016/j.psyneuen.2009.05.023.
- [84] Hawes JE, Tesic D, Whitehouse AJ, Zosky GR, Smith JT, Wyrwoll CS. Maternal vitamin D deficiency alters fetal brain development in the BALB/c mouse. *Behav Brain Res.* 2015 Jun 1;286:192–200.
- [85] Whitehouse AJ, Holt BJ, Serralha M, Holt PG, Kusel MM, Hart PH. Maternal serum vitamin D levels during pregnancy and offspring neurocognitive development. *Pediatrics.* 2012 Mar;129(3):485–93.
- [86] Latimer CS et al. Vitamin D prevents cognitive decline and enhances hippocampal synaptic function in aging rats. *Proc Natl Acad Sci U S A.* 2014 Oct 14;111(41):E4359–66. doi: 10.1073/pnas.1404477111.
- [87] De la Fuente AG et al. Vitamin D receptor-retinoid X receptor heterodimer signaling regulates oligodendrocyte progenitor cell differentiation. *J Cell Biol.* 2015 Dec 7;211(5):975–85. doi: 10.1083/jcb.201505119.
- [88] Landel V, Annweiler C, Millet P, Morello M, Féron F. Vitamin D, cognition, and Alzheimer's disease: the therapeutic benefit is in the D-tails. *J Alzheimer's Dis.* 2016 May 11. doi: 10.3233/JAD-150943. [Epub ahead of print].
- [89] Winner B, Winkler J. Adult neurogenesis in neurodegenerative diseases. *Cold Spring Harb Perspect Biol.* 2015 Apr 1;7(4):a021287. doi: 10.1101/cshperspect.a021287.
- [90] Brown J, Bianco JJ, McGrath JJ, Eyles DW. 1,25-dihydroxyvitamin D3 induces nerve growth factor, promotes neurite outgrowth and inhibits mitosis in embryonic rat hippocampal neurons. *Neurosci Lett.* 2003 Jun 5;343(2):139–43.
- [91] Ko P, Burkert R, McGrath J, Eyles D. Maternal vitamin D3 deprivation and the regulation of apoptosis and cell cycle during rat brain development. *Brain Res Dev Brain Res.* 2004 Oct 15;153(1):61–8.
- [92] Osborne JE, Hutchinson PE. Vitamin D and systemic cancer: is this relevant to malignant melanoma?. *Br J Dermatol.* 2002 Aug;147(2):197–213.
- [93] Cui X, McGrath JJ, Burne TH, Mackay-Sim A, Eyles DW. Maternal vitamin D depletion alters neurogenesis in the developing rat brain. *Int J Dev Neurosci.* 2007 Jun;25(4):227–32.
- [94] Keilhoff G, Grecksch G, Becker A. Haloperidol normalized prenatal vitamin D depletion-induced reduction of hippocampal cell proliferation in adult rats. *Neurosci Lett.* 2010 May 31;476(2):94–8. doi: 10.1016/j.neulet.2010.04.010.

- [95] Zhu Y et al. Abnormal neurogenesis in the dentate gyrus of adult mice lacking 1,25-dihydroxy vitamin D<sub>3</sub> (1,25-(OH)<sub>2</sub>D<sub>3</sub>). *Hippocampus*. 2012 Mar;22(3):421–33. doi: 10.1002/hipo.20908.
- [96] Annweiler C, Karras SN, Anagnostis P, Beauchet O. Vitamin D supplements: a novel therapeutic approach for Alzheimer patients. *Front Pharmacol*. 2014 Jan 28;5:6.
- [97] Andholf BKS, Sandhoff K. My journey into the world of sphingolipids and sphingolipidoses. *Proc Jpn Acad Ser B Phys Biol Sci*. 2012;88(10):554–82. doi: 10.2183/pjab.88.554.
- [98] Lingwood D, Simons K. Lipid rafts as a membrane-organizing principle. *Science*. 2010 Jan 1;327(5961):46–50. doi: 10.1126/science.1174621.
- [99] Zhou K, Blom T. Trafficking and functions of bioactive sphingolipids: lessons from cells and model membranes. *Lipid Insights*. 2015;8(S1):11–20. doi: 10.4137/LPI.S31615.
- [100] Gomez-Muñoz A, Frago LM, Alvarez L, Varela-Nieto I. Stimulation of DNA synthesis by natural ceramide 1-phosphate. *Biochem J*. 1997 Jul 15;325(Pt 2):435–40. doi: 10.1042/bj3250435.
- [101] Saddoughi SA, Ogretmen B. Diverse functions of ceramide in cancer cell death and proliferation. *Adv Cancer Res*. 2013;117:37–58. doi: 10.1016/B978-0-12-394274-6.00002-9.
- [102] Schwab SR, Pereira JP, Matloubian M, Xu Y, Huang Y, Cyster JG. Lymphocyte sequestration through S1P lyase inhibition and disruption of S1P gradients. *Science*. 2005 Sep 9;309(5741):1735–9. doi: 10.1126/science.1113640.
- [103] Podbielska M, Krotkiewski H, Hogan EL. Signaling and regulatory functions of bioactive sphingolipids as therapeutic targets in multiple sclerosis. *Neurochem Res*. 2012 Jun;37(6):1154–69. doi: 10.1007/s11064-012-0728-y.
- [104] Tidhar R, Futerman AH. The complexity of sphingolipid biosynthesis in the endoplasmic reticulum. *Biochim Biophys Acta*. 2013 Nov;1833(11):2511–8. doi: 10.1016/j.bbamcr.
- [105] Sato K et al. Critical role of ABCA1 transporter in sphingosine 1-phosphate release from astrocytes. *J Neurochem*. 2007 Dec;103(6):2610–9. doi: 10.1111/j.1471-4159.2007.04958.x.
- [106] Hisano Y, Kobayashi N, Kawahara A, Yamaguchi A, Nishi T. The sphingosine 1-phosphate transporter, SPNS2, functions as a transporter of the phosphorylated form of the immunomodulating agent FTY720. *J Biol Chem*. 2011;286(3):1758–66. doi: 10.1074/jbc.M110.171116.
- [107] Proia RL, Hla T. Emerging biology of sphingosine-1-phosphate: its role in pathogenesis and therapy. *J Clin Invest*. 2015 Apr;125(4):1379–87. doi: 10.1172/JCI76369.
- [108] Khavandgar Z, Poirier C, Clarke CJ, Li J, Wang N, McKee MD, Hannun YA, Murshed. A cell-autonomous requirement for neutral sphingomyelinase 2 in bone mineralization. *J Cell Biol*. 2011;194(2):277–89. doi: 10.1083/jcb.201102051.
- [109] Stoffel W. Functional analysis of acid and neutral sphingomyelinases in vitro and in vivo. *Chem Phys Lipids*. 1999 Nov;102(1–2):107–21. doi: 10.1016/S0009-3084(99)00079-1.

- [110] Dziak R. The role of sphingosine-1-phosphate (S1P) and lysophosphatidic acid (LPA) in regulation of osteoclastic and osteoblastic cells. *Immunol Invest.* 2013;42(7):510–8. doi: 10.3109/08820139.2013.823804.
- [111] Kozawa O, Niwa M, Matsuno H, Tokuda H, Miwa M, Ito H, Kato K, Uematsu T. Sphingosine 1-phosphate induces heat shock protein 27 via p38 mitogen-activated protein kinase activation in osteoblasts. *J Bone Miner Res.* 1999 Oct;14(10):1761–7. doi: 10.1359/jbmr.1999.14.10.1761.
- [112] Ishii M, Egen JG, Klauschen F, Meier-Schellersheim M, Saeki Y, Vacher J, Proia RL, Germain RN. Sphingosine-1-phosphate mobilizes osteoclast precursors and regulates bone homeostasis. *Nature.* 2009 Mar 26;458(7237):524–8. doi: 10.1038/nature07713.
- [113] Lotinun S et al. Osteoclast-specific cathepsin K deletion stimulates S1P-dependent bone formation. *J Clin Invest.* 2013 Feb;123(2):666–81. doi: 10.1172/JCI64840.
- [114] Quint P, Ruan M, Pederson L, Kassem M, Westendorf JJ, Khosla S, Oursler MJ. Sphingosine 1-phosphate (S1P) receptors 1 and 2 coordinately induce mesenchymal cell migration through S1P activation of complementary kinase pathways. *J Biol Chem.* 2013 Feb 22;288(8):5398–406. doi: 10.1074/jbc.M112.413583.
- [115] Keller J et al. Calcitonin controls bone formation by inhibiting the release of sphingosine 1-phosphate from osteoclasts. *Nat Commun.* 2014 Oct 21;5:5215. doi: 10.1038/ncomms6215.
- [116] Matsuzaki E et al. Sphingosine-1-phosphate promotes the nuclear translocation of  $\beta$ -catenin and thereby induces osteoprotegerin gene expression in osteoblast-like cell lines. *Bone.* 2013 Aug;55(2):315–24. doi: 10.1016/j.bone.2013.04.008.
- [117] Liu R, Farach-Carson MC, Karin NJ. Effects of sphingosine derivatives on MC3T3-E1 pre-osteoblasts: psychosine elicits release of calcium from intracellular stores. *Biochem Biophys Res Commun.* 1995;214(2):676–84. doi: 10.1006/bbrc.1995.2339.
- [118] Lyons JM, Karin NJ. A role for G protein-coupled lysophospholipid receptors in sphingolipid-induced  $\text{Ca}^{2+}$  signaling in MC3T3-E1 osteoblastic cells. *J Bone Miner Res.* 2001;16(11):2035–42. doi: 10.1359/jbmr.2001.16.11.2035.
- [119] Grey A, Chen Q, Callon K, Xu X, Reid IR, Cornish J. The phospholipids sphingosine-1-phosphate and lysophosphatidic acid prevent apoptosis in osteoblastic cells via a signaling pathway involving G(i) proteins and phosphatidylinositol-3 kinase. *Endocrinology.* 2002;143(12):4755–63. doi: 10.1210/en.2002-220347.
- [120] Ishii M, Kikuta J. Sphingosine-1-phosphate signaling controlling osteoclasts and bone homeostasis. *Biochim Biophys Acta.* 2013 Jan;1831(1):223–7. doi: 10.1016/j.bbalip.2012.06.002.
- [121] Okamoto H, Takuwa N, Yokomizo T, Sugimoto N, Sakurada S, Shigematsu H, Takuwa Y. Inhibitory regulation of Rac activation, membrane ruffling, and cell migration by the G protein-coupled sphingosine-1-phosphate receptor EDG5 but not EDG1 or EDG3. *Mol Cell Biol.* 2000 Dec;20(24):9247–61. doi: 10.1128/MCB.20.24.9247-9261.2000.



- [122] Kim BJ et al. Plasma sphingosine 1-phosphate levels and the risk of vertebral fracture in postmenopausal women. *J Clin Endocrinol Metab.* 2012 Oct;97(10):3807–14. doi: 10.1210/jc.2012-2346.
- [123] Marycz K, Krzak J, Marędzia M, Tomaszewski KA, Szczurek A, Moszak K. The influence of metal-based biomaterials functionalized with sphingosine-1-phosphate on the cellular response and osteogenic differentiation potential of human adipose derived mesenchymal stem cells in vitro. *J Biomater Appl.* 2016 Jan 22. doi: 10.1177/0885328216628711.
- [124] Aubin I et al. A deletion in the gene encoding sphingomyelin phosphodiesterase3 results in osteogenesis and dentinogenesis imperfecta in the mouse. *Nat Genet.* 2005;37(8):803–5.
- [125] Stoffel W, Jenke B, Block B, Zumbansen M, Koebke J. Neutral sphingomyelinase 2 (smpd3) in the control of postnatal growth and development. *Proc Natl Acad Sci USA.* 2005;102(12):4554–9. doi: 10.1073/pnas.0406380102.
- [126] Yadav MC, Simao AM, Narisawa S, Huesa C, McKee MD, Farquharson C, Millan JL. Loss of skeletal mineralization by the simultaneous ablation of PHOSPHO1 and alkaline phosphatase function: a unified model of the mechanisms of initiation of skeletal calcification. *J Bone Miner Res.* 2011;26(2):286–97. doi: 10.1002/jbmr.195.
- [127] Kakoi H et al. Bone morphogenic protein (BMP) signaling upregulates neutral sphingomyelinase 2 to suppress chondrocyte maturation via the Akt protein signaling pathway as a negative feedback mechanism. *J Biol Chem.* 2014;289(12):8135–50. doi: 10.1074/jbc.M113.509331.
- [128] Li Z et al. Choline kinase beta is required for normal endochondral bone formation. *Biochim Biophys Acta.* 2014;1840(7):2112–22. doi: 10.1016/j.bbagen.2014.03.008.
- [129] Haimovitz-Friedman A, Kan CC, Ehleiter D, Persaud RS, McLoughlin M, Fuks Z, Kolesnick RN. Ionizing radiation acts on cellular membranes to generate ceramide and initiate apoptosis. *J Exp Med.* 1994 Aug 1;180(2):525–35. non lo trovo.
- [130] Hill PA, Tumber A. Ceramide-induced cell death/survival in murine osteoblasts. *J Endocrinol.* 2010 Aug;206(2):225–33. doi: 10.1677/JOE-10-0068.
- [131] Takeda H, Ozaki K, Yasuda H, Ishida M, Kitano S, Hanazawa S. Sphingomyelinase and ceramide inhibit formation of F-actin ring in and bone resorption by rabbit mature osteoclasts. *FEBS Lett.* 1998 Jan 30;422(2):255–8. doi: 10.1016/S0014-5793(98)00005-2.
- [132] Zhai L, Sun N, Han Z, Jin HC, Zhang B. Liposomal short-chain C6 ceramide induces potent anti-osteosarcoma activity in vitro and in vivo. *Biochem Biophys Res Commun.* 2015 Dec 4;468(1–2):274–80. doi: 10.1016/j.bbrc.2015.10.113.
- [133] Takeshita A, Watanabe A, Takada Y, Hanazawa S. Selective stimulation by ceramide of the expression of the alpha isoform of retinoic acid and retinoid X receptors in osteoblastic cells. A role of sphingosine 1-phosphate-mediated AP-1 in the ligand-dependent

- transcriptional activity of these receptors. *J Biol Chem.* 2000 Oct 13;275(41):32220–6. doi: 10.1074/jbc.M002569200.
- [134] Olivier S, Fillet M, Malaise M, Piette J, Bours V, Merville MP, Franchimont N. Sodium nitroprusside-induced osteoblastapoptosis is mediated by long chain ceramide and is decreased by raloxifene. *Biochem Pharmacol.* 2005 Mar 15;69(6):891–901. doi: 10.1016/j.bcp.2004.11.030.
- [135] Kitajima I, Soejima Y, Takasaki I, Beppu H, Tokioka T, Maruyama I. Ceramide-induced nuclear translocation of NF-kappa B is a potential mediator of the apoptotic response to TNF-alpha in murine clonal osteoblasts. *Bone.* 1996;19(3):263–70. doi: 10.1016/8756-3282(96)00181-0.
- [136] Chae HJ et al. Dexamethasone suppresses tumor necrosis factor-alpha-induced apoptosis in osteoblasts: possible role for ceramide. *Endocrinology.* 2000 Aug;141(8):2904–13. doi: 10.1210/endo.141.8.7604.
- [137] Kolesnick RN, Krönke M. Regulation of ceramide production and apoptosis. *Annu Rev Physiol.* 1998;60:643–65. doi: 10.1146/annurev.physiol.60.1.643.
- [138] Ghosh N, Patel N, Jiang K, Watson JE, Cheng J, Chalfant CE, Cooper DR. Ceramide-activated protein phosphatase involvement in insulin resistance via Akt, serine/arginine-rich protein 40, and ribonucleic acid splicing in L6 skeletal muscle cells. *Endocrinology.* 2007 Mar;148(3):1359–66. doi: 10.1210/en.2006-0750.
- [139] Hannun YA. Functions of ceramide in coordinating cellular responses to stress. *Science.* 1996 Dec;274(5294):1855–9. doi: 10.1126/science.274.5294.1855.
- [140] Sabourdy F et al. Monogenic neurological disorders of sphingolipid metabolism. *Biochim Biophys Acta.* 2015 Aug;1851(8):1040–51. doi: 10.1016/j.bbalip.2015.01.010.
- [141] Haughey NJ, Bandaru VV, Bae M, Mattson MP. Roles for dysfunctional sphingolipid metabolism in Alzheimer's disease neuropathogenesis. *Biochim Biophys Acta.* 2010 Aug;1801(8):878–86. doi: 10.1016/j.bbalip.2010.05.003.
- [142] Grimm MO, Zimmer VC, Lehmann J, Grimm HS, Hartmann T. The impact of cholesterol, DHA, and sphingolipids on Alzheimer's disease. *Biomed Res Int.* 2013;2013:814390. doi: 10.1155/2013/814390.
- [143] Murphy KE, Gysbers AM, Abbott SK, Tayebi N, Kim WS, Sidransky E, Cooper A, Garner B, Halliday GM. Reduced glucocerebrosidase is associated with increased  $\alpha$ -synuclein in sporadic Parkinson's disease. *Brain.* 2014 Mar;137(Pt 3):834–48. doi: 10.1093/brain/awt367.
- [144] Abbott SK, Li H, Muñoz SS, Knoch B, Batterham M, Murphy KE, Halliday GM, Garner B. Altered ceramide acyl chain length and ceramide synthase gene expression in Parkinson's disease. *Mov Disord.* 2014 Apr;29(4):518–26. doi: 10.1002/mds.25729.
- [145] Halmer R, Walter S, Faßbender K. Sphingolipids: important players in multiple sclerosis. *Cell Physiol Biochem.* 2014;34(1):111–8. doi: 10.1159/000362988.

- [146] Gulbins E et al. A central role for the acid sphingomyelinase/ceramide system in neurogenesis and major depression. *J Neurochem.* 2015 Jul;134(2):183–92. doi: 10.1111/jnc.13145.
- [147] Mencarelli C, Martinez-Martinez P. Ceramide function in the brain: when a slight tilt is enough. *Cell Mol Life Sci.* 2013 Jan;70(2):181–203.
- [148] Shamseddine AA, Airola MV, Hannun YA. Roles and regulation of neutral sphingomyelinase-2 in cellular and pathological processes. *Adv Biol Regul.* 2015 Jan;57:24–41. doi: 10.1016/j.jbior.2014.10.002.
- [149] Ghasemi R, Dargahi L, Ahmadiani A. Integrated sphingosine-1 phosphate signaling in the central nervous system: from physiological equilibrium to pathological damage. *Pharmacol Res.* 2016 Jan 7;104:156–64. doi: 10.1016/j.phrs.2015.11.006.
- [150] Lucki NC, Sewer MB. Multiple roles for sphingolipids in steroid hormone biosynthesis. *Subcell Biochem.* 2008;49:387–412. doi: 10.1007/978-1-4020-8831-5\_15.
- [151] Titievsky A, Titievskaya I, Pasternack M, Kaila K, Törnquist K. Sphingosine inhibits voltage-operated calcium channels in GH4C1 cells. *J Biol Chem.* 1998 Jan 2;273(1):242–7.
- [152] Qin X et al. Sphingosine and FTY720 are potent inhibitors of the transient receptor potential melastatin 7 (TRPM7) channels. *Br J Pharmacol.* 2013 Mar;168(6):1294–312. doi: 10.1111/bph.12012.
- [153] Rohrbough J, Rushton E, Palanker L, Woodruff E, Matthies HJ, Acharya U. Ceramidase regulates synaptic vesicle exocytosis and trafficking. *J Neurosci.* 2004 Sep 8;24(36):7789–803.
- [154] Darios F et al. Sphingosine facilitates SNARE complex assembly and activates synaptic vesicle exocytosis. *Neuron.* 2009 Jun 11;62(5):683–94. doi: 10.1016/j.neuron.2009.04.024.
- [155] Paugh SW, Cassidy MP, He H, Milstien S, Sim-Selley LJ, Spiegel S, Selley DE. Sphingosine and its analog, the immunosuppressant 2-amino-2-(2-[4-octylphenyl] ethyl)-1,3-propanediol, interact with the CB1 cannabinoid receptor. *Mol Pharmacol.* 2006 Jul;70(1):41–50.
- [156] Selley DE, Welch SP, Sim-Selley LJ. Sphingosine lysolipids in the CNS: endogenous cannabinoid antagonists or a parallel pain modulatory system? *Life Sci.* 2013 Aug 14;93(5–6):187–93. doi: 10.1016/j.lfs.2013.06.004.
- [157] Mizugishi K, Yamashita T, Olivera A, Miller GF, Spiegel S, Proia RL. Essential role for sphingosine kinases in neural and vascular development. *Mol Cell Biol.* 2005 Dec;25(24):11113–21.
- [158] Kimura A et al. Essential roles of sphingosine1-phosphate/S1P1 receptor axis in the migration of neural stem cells toward a site of spinal cord injury. *Stem Cells.* 2007 Jan;25(1):115–24. doi: 10.1634/stemcells.2006-0223.

- [159] Novgorodov AS, El-Alwani M, Bielawski J, Obeid LM, Gudz TI. Activation of sphingosine-1-phosphate receptor S1P5 inhibits oligodendrocyte progenitor migration. *FASEB J*. 2007 May;21(7):1503–14.
- [160] MacLennan AJ et al. An essential role for the H218/AGR16/Edg-5/LP(B2) sphingosine 1-phosphate receptor in neuronal excitability. *Eur J Neurosci*. 2001 Jul;14(2):203–9.
- [161] Jaillard C et al. Edg8/S1P5: an oligodendroglial receptor with dual function on process retraction and cell survival. *J Neurosci*. 2005 Feb 9;25(6):1459–69.
- [162] Toman RE, Payne SG, Watterson KR, Maceyka M, Lee NH, Milstien S, Bigbee JW, Spiegel S. Differential transactivation of sphingosine-1-phosphate receptors modulates NGF-induced neurite extension. *J Cell Biol*. 2004 Aug 2;166(3):381–92.
- [163] Kanno T, Nishizaki T, Proia RL, Kajimoto T, Jahangeer S, Okada T, Nakamura S. Regulation of synaptic strength by sphingosine 1-phosphate in the hippocampus. *Neuroscience*. 2010 Dec 29;171(4):973–80. doi: 10.1016/j.neuroscience.2010.10.021.
- [164] Li C, Li JN, Kays J, Guerrero M, Nicol GD. Sphingosine 1-phosphate enhances the excitability of rat sensory neurons through activation of sphingosine 1-phosphate receptors 1 and/or 3. *J Neuroinflamm*. 2015 Apr 12;12:70. doi: 10.1186/s12974-015-0286-8.
- [165] Chan JP, Sieburth D. Localized sphingolipid signaling at presynaptic terminals is regulated by calcium influx and promotes recruitment of priming factors. *J Neurosci*. 2012 Dec 5;32(49):17909–20. doi: 10.1523/JNEUROSCI.2808-12.2012.
- [166] Shen H et al. Coupling between endocytosis and sphingosine kinase 1 recruitment. *Nat Cell Biol*. 2014 Jul;16(7):652–62. doi: 10.1038/ncb2987.
- [167] Wang G, Dinkins M, He Q, Zhu G, Poirier C, Campbell A, Mayer-Proschel M, Bieberich E. Astrocytes secrete exosomes enriched with proapoptotic ceramide and prostate apoptosis response 4 (PAR-4): potential mechanism of apoptosis induction in Alzheimer disease (AD). *J Biol Chem*. 2012 Jun 15;287(25):21384–95. doi: 10.1074/jbc.M112.340513.
- [168] Scarlatti F, Bauvy C, Ventruti A, Sala G, Cluzeaud F, Vandewalle A, Ghidoni R, Codogno P. Ceramide-mediated macroautophagy involves inhibition of protein kinase B and up-regulation of beclin 1. *J Biol Chem*. 2004 Apr 30;279(18):18384–91.
- [169] Ben-David O, Futerman AH. The role of the ceramide acyl chain length in neurodegeneration: involvement of ceramide synthases. *Neuromol Med*. 2010 Dec;124:341–50. doi: 10.1007/s12017-010-8114-x.
- [170] Zhao L et al. A deficiency of ceramide biosynthesis causes cerebellar purkinje cell neurodegeneration and lipofuscin accumulation. *PLoS Genet*. 2011 May;7(5):e1002063. doi: 10.1371/journal.pgen.1002063.
- [171] Ginkel C et al. Ablation of neuronal ceramide synthase 1 in mice decreases ganglioside levels and expression of myelin-associated glycoprotein in oligodendrocytes. *J Biol Chem*. 2012 Dec 7;287(50):41888–902. doi: 10.1074/jbc.M112.413500.

- [172] Kim SK et al. Neutral sphingomyelinase 2 induces dopamine uptake through regulation of intracellular calcium. *Cell Signal*. 2010 May;22(5):865–70. doi: 10.1016/j.cellsig.2010.01.012.
- [173] Wheeler D et al. Tumor necrosis factor- $\alpha$ -induced neutral sphingomyelinase-2 modulates synaptic plasticity by controlling the membrane insertion of NMDA receptors. *J Neurochem*. 2009 Jun;109(5):1237–49. doi: 10.1111/j.1471-4159.2009.06038.x.
- [174] Tabatadze N, Savonenko A, Song H, Bandaru VV, Chu M, Haughey NJ. Inhibition of neutral sphingomyelinase-2 perturbs brain sphingolipid balance and spatial memory in mice. *J Neurosci Res*. 2010 Oct;88(13):2940–51. doi: 10.1002/jnr.22438.
- [175] Gu L, Huang B, Shen W, Gao L, Ding Z, Wu H, Guo J. Early activation of nSMase2/ceramide pathway in astrocytes is involved in ischemia-associated neuronal damage via inflammation in rat hippocampi. *J Neuroinflamm*. 2013 Sep 3;10:109. doi: 10.1186/1742-2094-10-109.
- [176] He Q, Wang G, Wakade S, Dasgupta S, Dinkins M, Kong JN, Spassieva SD, Bieberich E. Primary cilia in stem cells and neural progenitors are regulated by neutral sphingomyelinase 2 and ceramide. *Mol Biol Cell*. 2014 Jun;25(11):1715–29. doi: 10.1091/mbc.E13-12-0730.
- [177] Bajjalieh SM, Martin TF, Floor E. Synaptic vesicle ceramide kinase. A calcium-stimulated lipid kinase that co-purifies with brain synaptic vesicles. *J Biol Chem*. 1989 Aug 25;264(24):14354–60.
- [178] Sugiura M, Kono K, Liu H, Shimizugawa T, Minekura H, Spiegel S, Kohama T. Ceramide kinase, a novel lipid kinase. Molecular cloning and functional characterization. *J Biol Chem*. 2002 Jun 28;277(26):23294–300.
- [179] Presa N, Gomez-Larrauri A, Rivera IG, Ordoñez M, Trueba M, Gomez-Muñoz A. Regulation of cell migration and inflammation by ceramide 1-phosphate. *Biochim Biophys Acta*. 2016 Feb 11. pii: S1388-1981(16)30028-2. doi: 10.1016/j.bbalip.2016.02.007.
- [180] Pettus BJ et al. Ceramide 1-phosphate is a direct activator of cytosolic phospholipase A2. *J Biol Chem*. 2004 Mar 19;279(12):11320–6.
- [181] Kim C et al. Ceramide-1-phosphate regulates migration of multipotent stromal cells and endothelial progenitor cells—implications for tissue regeneration. *Stem Cells*. 2013 Mar;313:500–10. doi: 10.1002/stem.1291.
- [182] Reardon BJ et al. Vitamin D-responsive SGPP2 variants associated with lung cell expression and lung function. *BMC Med Genet*. 2013 Nov 25;14:122.
- [183] Kleuser B, Cuvillier O, Spiegel S. 1 $\alpha$ ,25-dihydroxyvitamin D3 inhibits programmed cell death in HL-60 cells by activation of sphingosine kinase. *Cancer Res*. 1998 May 1;58(9):1817–24.

- [184] Manggau M, Kim DS, Ruwisch L, Vogler R, Korting HC, Schäfer-Korting M, Kleuser B. 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> protects human keratinocytes from apoptosis by the formation of sphingosine-1-phosphate. *J Invest Dermatol*. 2001 Nov;117(5):1241–9.
- [185] Kikuta J, Kawamura S, Okiji F, Shirazaki M, Sakai S, Saito H, Ishii M. Sphingosine-1-phosphate-mediated osteoclast precursor monocyte migration is a critical point of control in antbone-resorptive action of active vitamin D. *Proc Natl Acad Sci U S A*. 2013 Apr 23;110(17):7009–13.
- [186] Selkoe DJ. The therapeutics of Alzheimer's disease: where we stand and where we are heading. *Ann Neurol*. 2013 Sep;74(3):328–36. doi: 10.1002/ana.24001.
- [187] Satoi H, Tomimoto H, Ohtani R, Kitano T, Kondo T, Watanabe M, Oka N, Akiguchi I, Furuya S, Hirabayashi Y, Okazaki T. Astroglial expression of ceramide in Alzheimer's disease brains: a role during neuronal apoptosis. *Neuroscience*. 2005;130(3):657–66.
- [188] Katsel P, Li C, Haroutunian V. Gene expression alterations in the sphingolipid metabolism pathways during progression of dementia and Alzheimer's disease: a shift toward ceramide accumulation at the earliest recognizable stages of Alzheimer's disease?. *Neurochem Res*. 2007 Apr–May;32(4–5):845–56.
- [189] Takasugi N et al. BACE1 activity is modulated by cell-associated sphingosine-1-phosphate. *J Neurosci*. 2011 May 4;31(18):6850–7. doi: 10.1523/JNEUROSCI.6467-10.2011.
- [190] Grimm MO, Rothhaar TL, Hartmann T. The role of APP proteolytic processing in lipid metabolism. *Exp Brain Res*. 2012 Apr;217(3–4):365–75. doi: 10.1007/s00221-011-2975-6.
- [191] Couttas TA, Lim XY, Don AS. A three-step assay for ceramide synthase activity using a fluorescent substrate and HPLC. *Lipids*. 2015 Jan;50(1):101–9.
- [192] Ceccom J et al. Reduced sphingosine kinase-1 and enhanced sphingosine 1-phosphate lyase expression demonstrate deregulated sphingosine 1-phosphate signaling in Alzheimer's disease. *Acta Neuropathol Commun*. 2014;2(1):12. doi: 10.1186/2051-5960-2-12.
- [193] Grimm C, Kraft R, Schultz G, Harteneck C. Activation of the melastatin-related cation channel TRPM3 by D-erythro-sphingosine. *Mol Pharmacol*. 2005 Mar;67(3):798–805. Epub 2004 Nov 18.
- [194] Jazvinščak Jembrek M, Hof PR, Šimić G. Ceramides in Alzheimer's disease: key mediators of neuronal apoptosis induced by oxidative stress and A $\beta$  accumulation. *Oxid Med Cell Longev*. 2015;2015:346783. doi: 10.1155/2015/346783.
- [195] Jana A, Pahan K. Fibrillar amyloid-beta peptides kill human primary neurons via NADPH oxidase-mediated activation of neutral sphingomyelinase. Implications for Alzheimer's disease. *J Biol Chem*. 2004 Dec 3;279(49):51451–9.



- [196] Malaplate-Armand C et al. Soluble oligomers of amyloid-beta peptide induce neuronal apoptosis by activating a cPLA2-dependent sphingomyelinase-ceramide pathway. *Neurobiol Dis.* 2006 Jul;23(1):178–89.
- [197] Puglielli L, Ellis BC, Saunders AJ, Kovacs DM. Ceramide stabilizes beta-site amyloid precursor protein-cleaving enzyme 1 and promotes amyloid beta-peptide biogenesis. *J Biol Chem.* 2003 May 30;278(22):19777–83.
- [198] Brunkhorst R, Vutukuri R, Pfeilschifter W. Fingolimod for the treatment of neurological diseases-state of play and future perspectives. *Front Cell Neurosci.* 2014 Sep 12;8:283. doi: 10.3389/fncel.2014.00283.
- [199] Asle-Rousta M, Kolahdooz Z, Oryan S, Ahmadiani A, Dargahi L. FTY720 (fingolimod) attenuates beta-amyloid peptide (A $\beta$ 42)-induced impairment of spatial learning and memory in rats. *J Mol Neurosci.* 2013 Jul;50(3):524–32. doi: 10.1007/s12031-013-9979-6.
- [200] Hemmati F, Dargahi L, Nasoohi S, Omidbakhsh R, Mohamed Z, Chik Z, Naidu M, Ahmadiani A. Neurorestorative effect of FTY720 in a rat model of Alzheimer's disease: comparison with memantine. *Behav Brain Res.* 2013 Sep 1;252:415–21. doi: 10.1016/j.bbr.2013.06.016.
- [201] Gezen-Ak D et al. Vitamin D receptor gene haplotype is associated with late-onset Alzheimer's disease. *Tohoku J Exp Med.* 2012;228(3):189–96.
- [202] Dursun E, Gezen-Ak D, Yilmazer S. Beta amyloid suppresses the expression of the vitamin d receptor gene and induces the expression of the vitamin d catabolic enzyme gene in hippocampal neurons. *Dement Geriatr Cogn Disord.* 2013;36:76–86.
- [203] Durk MR, Han K, Chow EC, Ahrens R, Henderson JT, Fraser PE, Pang KS. 1 $\alpha$ ,25-Dihydroxyvitamin D3 reduces cerebral amyloid- $\beta$  accumulation and improves cognition in mouse models of Alzheimer's disease. *J Neurosci.* 2014 May 21;34(21):7091–101.
- [204] Meacci E, Bini F, Cecchi M, Garcia-Gil M. Role of sphingolipids on the cytotoxicity induced by 1–42 beta amyloid peptide in differentiated SH-SY5Y cells. *Acta Physiol.* 2008;194:665.